



Time Remaining: 18 hours 45 min Rank: 11497 Score: 0

malarkannan.p@gmail.com | [Contest scoreboard](#) | [Sign out](#)

Qualification Round 2016

[A. Counting Sheep](#)[B. Revenge of the Pancakes](#)[C. Coin Jam](#)**D. Fractiles**[Ask a question](#) **1**[View my submissions](#)

Submissions	
Counting Sheep	
7pt	Not attempted 11137/12501 users correct (89%)
8pt	Not attempted 10480 users attempted
Revenge of the Pancakes	
10pt	Not attempted 6926/7394 users correct (94%)
10pt	Not attempted 6566 users attempted
Coin Jam	
10pt	Not attempted 2700/3171 users correct (85%)

Problem D. FractilesConfused? Read the [quick-start guide](#).Small input
10 points

Solve D-small

You may try multiple times, with penalties for wrong submissions.

Large input
25 points

You must solve the small input first.

You have 8 minutes to solve 1 input file. (Judged after contest.)

Problem

Long ago, the Fractal civilization created artwork consisting of linear rows of tiles. They had two types of tile that they could use: gold (G) and lead (L).

Each piece of Fractal artwork is based on two parameters: an original sequence of **K** tiles, and a complexity **C**. For a given original sequence, the artwork with complexity 1 is just that original sequence, and the artwork with complexity $X+1$ consists of the artwork with complexity X , transformed as follows:

- replace each L tile in the complexity X artwork with another copy of the original sequence
- replace each G tile in the complexity X artwork with **K** G tiles

For example, for an original sequence of LGL, the pieces of artwork with complexity 1 through 3 are:

- **C** = 1: LGL (which is just the original sequence)
- **C** = 2: LGLGGGLGL
- **C** = 3: LGLGGGLGLGGGGGGGLGLGGGLGL

20pt Not attempted
1718 users attempted

Fractiles


10pt Not attempted
1451/1665 users
 correct (87%)

25pt Not attempted
820 users attempted

- Top Scores

Lewin	100
Endagorion	100
xiaowuc1	100
xyz111	100
HellKitsune123	100
h4tguy	100
YuryBandarchuk	100
ivan.popelyshev	100
burunduk3	100
Aksenov239	100

Here's an illustration of how the artwork with complexity 2 is generated from the artwork with complexity 1:



You have just discovered a piece of Fractal artwork, but the tiles are too dirty for you to tell what they are made of. Because you are an expert archaeologist familiar with the local Fractal culture, you know the values of **K** and **C** for the artwork, but you do not know the original sequence. Since gold is exciting, you would like to know whether there is at least one G tile in the artwork. Your budget allows you to hire **S** graduate students, each of whom can clean one tile of your choice (out of the K^C tiles in the artwork) to see whether the tile is G or L.

Is it possible for you to choose a set of no more than **S** specific tiles to clean, such that *no matter what* the original pattern was, you will be able to know for sure whether at least one G tile is present in the artwork? If so, which tiles should you clean?

Input

The first line of the input gives the number of test cases, **T**. **T** test cases follow. Each consists of one line with three integers: **K**, **C**, and **S**.

Output

For each test case, output one line containing Case #*x*: *y*, where *x* is the test case number (starting from 1) and *y* is either IMPOSSIBLE if no set of tiles will answer your question, or a list of between 1 and **S** positive integers, which are the positions of the tiles that will answer your question. The tile positions are numbered from 1 for the leftmost tile to K^C for the rightmost tile. Your chosen positions may be in any order, but they must all be different.

If there are multiple valid sets of tiles, you may output any of them. Remember that once you submit a Small and it is accepted, you will not be able to download and submit another Small input. See [the FAQ](#) for a more thorough explanation. **This reminder**

won't appear in problems in later rounds.

Limits

$$1 \leq T \leq 100.$$

$$1 \leq K \leq 100.$$

$$1 \leq C \leq 100.$$

$$K^C \leq 10^{18}.$$

Small dataset

$$S = K.$$

Large dataset

$$1 \leq S \leq K.$$

Sample

Input	Output
5	Case #1: 2
2 3 2	Case #2: 1
1 1 1	Case #3: IMPOSSIBLE
2 1 1	Case #4: 1 2
2 1 2	Case #5: 2 6
3 2 3	

Note: for some of these sample cases, other valid solutions exist.

In sample case #1, there are four possible original sequences: GG, GL, LG, and LL. They would produce the following artwork, respectively:

- Original sequence GG: GGGGGGGG

- Original sequence GL: GGGGGGGL
- Original sequence LG: LGGGGGGG
- Original sequence LL: LLLLLLLL

One valid solution is to just look at tile #2. If tile #2 turns out to be G, then you will know for sure the artwork contains at least one G. (You will not know whether the original sequence is GG, GL, or LG, but that doesn't matter.) If tile #2 turns out to be L, then you will know that the original sequence must be LL, so there are no Gs in the artwork. So 2 is a valid solution.

On the other hand, it would not be valid to just look at tile #1. If it turns out to be L, you will only know that the original sequence could have been either LG or LL. If the original sequence is LG, there is at least one G in the artwork, but if the original sequence is LL, there are no Gs. So 1 would not be a valid solution.

Note that 1 2 is also a valid solution, because tile #2 already provides all the information you need. 1 2 3 is not a valid solution, because it uses too many tiles.

In sample case #2, the artwork must consist of only one tile: either G or L. Looking at that tile will trivially tell you whether or not the artwork has a G in it.

In sample case #3, which would not appear in the Small dataset, the artwork must be either GG, GL, LG, or LL. You can only look at one tile, and neither of them on its own is enough to answer the question. If you see L for tile #1, you will not know whether the artwork is LG or LL, so you will not know whether any Gs are present. If you see L for tile #2, you will not know whether the artwork is GL or LL, so you will not know whether any Gs are present.

Sample case #4 is like sample case #3, but with access to one more tile. Now you can just look at the entire artwork.

In sample case #5, there are eight possible original sequences, and they would produce the following artwork:

- Original sequence GGG: GGGGGGGGG
- Original sequence GGL: GGGGGGGGL
- Original sequence GLG: GGGGLGGGG
- Original sequence GLL: GGGLLGLL

- Original sequence LGG: LGGGGGGGG
- Original sequence LGL: LGLGGGLGL
- Original sequence LLG: LLGLLGGGG
- Original sequence LLL: LLLLLLLLLL

One valid solution is to look at tiles #2 and #6. If they both turn out to be Ls, the artwork must be all Ls. Otherwise, there must at least one G. Note that 1 2 would not be a valid solution, because even if those tiles both turn out to be Ls, that does not rule out an original sequence of LLG. 6 2 would be a valid solution, since the order of the positions in your solution does not matter.

All problem statements, input data and contest analyses are licensed under the [Creative Commons Attribution License](#).

© 2008-2015 Google [Google Home](#) - [Terms and Conditions](#) - [Privacy Policies and Principles](#)

