

Bell Violations Without Entanglement?

A Reproducible Rebuttal of Wang et al. (2025)

Meshal Alawein
UC Berkeley
meshal@berkeley.edu

March 30, 2026

Abstract

A recent *Science Advances* paper reports $S = 2.275 \pm 0.057$ at an effective acceptance or post-selection probability of roughly 10^{-18} and interprets that statistic as Bell-inequality violation without entanglement [17]. The present paper makes a narrower claim, even though three published critiques now converge on the stronger diagnosis that the reported violation is generated by fourfold post-selection and unconventional normalization rather than by a loophole-free demonstration of nonlocality [18, 19, 20]. In such a severely filtered regime, the scalar fact $S > 2$ is not exclusionary evidence against local models unless the selection mechanism is independently constrained. We formalize this point in three steps. First, we write the selected estimator as a conditional expectation on accepted events and identify fair sampling as the assumption that transfers the classical CHSH bound from the emitted ensemble to the retained one. Second, we construct a deterministic local hidden-variable model with an outcome-coupled acceptance rule and derive the exact analytic toy-model expectations $E_{\text{sel}} = \pm 13/14$ and $S_{\text{sel}} = 26/7 \approx 3.714$. Third, we compare those exact expectations with the site’s Monte Carlo benchmark, which yields $S = 3.716$ (95% CI [3.714, 3.717]) at **70.0%** acceptance, and with the Wang et al. report. The evidentiary consequence is limited but decisive: in an ultra-low-efficiency Bell test, a post-selected CHSH value above 2 is suggestive at most until raw correlations, setting-resolved acceptance rates, and fair-sampling diagnostics are reported.

Claim. In the ultra-low-efficiency regime, a post-selected CHSH value above 2 is non-exclusionary evidence until the selection rule itself is constrained.

Published findings.

- The Monte Carlo benchmark remains $S = 3.716$ (95% CI [3.714, 3.717]) at **70.0%** acceptance for a purely local source.
- The selected estimator is a conditional expectation on accepted events, so the naive $S \leq 2$ inference survives only under fair-sampling assumptions.
- At Wang et al.’s reported effective acceptance of about 10^{-18} , diagnostics of the selection rule are more informative than the scalar statistic alone.

1 Introduction

Bell’s theorem excludes local hidden-variable explanations for the full set of quantum correlations under the usual assumptions of locality, realism, and setting independence [1]. The Clauser-Horne-Shimony-Holt inequality packages that exclusion into an experimentally testable scalar, and Fine sharpened the probabilistic picture by relating CHSH satisfaction to the existence of a joint distribution over the counterfactual outcomes [2, 7].

$$S = |E_{00} + E_{01} + E_{10} - E_{11}|$$

In the fair-sampling regime, the local bound is $S \leq 2$ and the quantum ceiling is the Tsirelson bound $2\sqrt{2}$ [10]. Wang et al. report a super-classical value from a multiphoton path-identity interference experiment and interpret it as Bell-inequality violation without entanglement [17].

The experiment-specific structure matters. Wang et al. use four-photon frustrated interference, or interwoven frustrated down-conversion, and retain only fourfold coincidences. Alice and Bob control local phases α and β , the missing outcome is inferred by a π -shift of the local phase, and the reported joint probability is reconstructed from counts gathered at four different settings rather than from a single local trial with definite outcomes:

$$p(++|\alpha, \beta) = \frac{N(\alpha, \beta)}{N(\alpha, \beta) + N(\alpha + \pi, \beta) + N(\alpha, \beta + \pi) + N(\alpha + \pi, \beta + \pi)}.$$

That normalization is operationally different from a standard Bell test, where each chosen setting yields a local ± 1 outcome without counterfactual renormalization. The quoted 10^{-18} figure is therefore better read as an effective acceptance rate for the selected fourfold subensemble than as an ordinary per-arm detector efficiency. Wang et al. are also explicit that the locality loophole and sampling loophole remain open and that the analysis relies on fair sampling [17].

The apparatus is sophisticated. The relevant issue is whether a post-selected statistic in an ultra-low-efficiency regime can support the inference being drawn from it.

This paper answers that question with a constructive counterexample and a narrower evidentiary standard. A large CHSH number can be compelling, but by itself it is not a loophole-closure criterion.

The present claim is evidentiary rather than ontological. A local counterexample need only show that the reported evidence class remains compatible with local explanations.

The published critiques sharpen the experiment-specific diagnosis in complementary ways. Wharton and Price construct a local classical model and frame the selection effect as collider or Berkson bias. Wójcik and Wójcik argue that post-selection plus the cross-setting normalization creates Bell-like numbers without Bell-operational meaning, and that the phases act more like source-intensity controls than local dichotomic settings. Cieřliński, Larsson, Markiewicz, Schlichtholz, and Żukowski show that the phase-shift protocol admits a local realistic model, while a different on-off switching protocol can still reveal genuine Bell nonclassicality from the same interferometric family.

2 Theoretical Background

2.1 Notation and Assumptions

We write Alice’s setting as $x \in \{0, 1\}$ and Bob’s as $y \in \{0, 1\}$. Outcomes are binary: $A_x, B_y \in \{-1, +1\}$. A hidden variable is denoted by λ , and $D \in \{0, 1\}$ marks whether a trial is retained by the analysis filter.

Table 1: Literature checkpoints relevant to the present claim.

Cluster	What it establishes	Why it matters here
Bell 1964; CHSH 1969 [1, 2]	Local hidden-variable models satisfy a bound on the full correlation structure.	The theorem targets the emitted ensemble, not an arbitrary filtered subset.
Pearle through Eberhard [3, 4, 5, 6]	Selection and efficiency can open a detection loophole unless explicitly controlled.	Wang et al. operate far inside the regime where that warning is operationally important.
Path identity optics [11, 12, 13]	Indistinguishable generation pathways can create striking interference effects.	The optical novelty is not being denied; the Bell inference is the narrower issue.
Loophole-aware Bell benchmarks [14, 15, 16]	Strong Bell claims are paired with explicit closure strategies, not just large S .	That is the comparison class for any foundational headline.
2025–26 critique literature [18, 19, 20]	Three independent critiques converge on fourfold post-selection, unconventional normalization, and setting semantics as the key interpretive issues.	The stronger artifact diagnosis in the literature goes beyond the narrower evidentiary claim made here.

Locality $A_x(\lambda)$ depends on x and λ , not on y ; symmetrically for Bob.

Realism All four counterfactual outcomes are taken to exist simultaneously for the local model under study.

Freedom $P(\lambda \mid x, y) = P(\lambda)$, so the settings are statistically independent of the hidden variable.

Fair Sampling The standard CHSH inference on selected data additionally requires D not to depend on the hidden outcome pattern in a biasing way.

Definition (CHSH Functional and Sign Convention). *In the convention used throughout this paper, the CHSH functional is*

$$S = |E_{00} + E_{01} + E_{10} - E_{11}|, \quad E_{xy} = \mathbb{E}[A_x B_y \mid x, y].$$

Relabeling settings or flipping one party’s output signs maps this expression to the other familiar CHSH sign placements without changing the attainable classical or quantum bounds. The notation changes; the admissible region does not.

Proposition 1 (Fair-Sampling Local Bound). *For any local hidden-variable model satisfying locality, realism, freedom, and fair sampling, the selected CHSH statistic obeys*

$$S_{\text{LHV}} \leq 2.$$

Proof Sketch. For each fixed λ , define

$$X(\lambda) = A_0(\lambda)[B_0(\lambda) + B_1(\lambda)] + A_1(\lambda)[B_0(\lambda) - B_1(\lambda)].$$




Figure 1: Reference scales for the two papers. Wang et al.’s reported value sits above the classical bound, but the exact local counterexample sits even higher. The figure is schematic rather than empirical; it shows what a filtered estimator can and cannot prove.

Because Bob’s outcomes are each ± 1 , exactly one bracket has magnitude 2 and the other vanishes, so $|X(\lambda)| \leq 2$. Integrating over λ preserves the bound. If the accepted sample is a fair image of the emitted ensemble, the same algebra applies to the retained data as well [1, 2, 7]. $\square$

Proposition 2 (Quantum Comparison Ceiling). *Within quantum mechanics, the maximal CHSH value is the Tsirelson bound*

$$S_{\text{QM}} \leq 2\sqrt{2} \approx 2.828.$$

Proof Sketch. Let A, A' and B, B' be Hermitian observables with eigenvalues ± 1 , and define the CHSH operator

$$\hat{S} = A \otimes (B + B') + A' \otimes (B - B').$$

The standard commutator calculation gives $\hat{S}^2 = 4I - [A, A'] \otimes [B, B']$, from which $\|\hat{S}^2\| \leq 8$ and therefore $\|\hat{S}\| \leq 2\sqrt{2}$. That ceiling is achievable for suitable entangled states and local observables [10]. $\square$

Proposition 3 (Selection Replaces the Unconditional Estimator). *Once analysis is conditioned on accepted trials, the operational correlation is not E_{xy} but*

$$E_{xy}^{(\text{sel})} = \frac{\mathbb{E}[A_x B_y D \mid x, y]}{\mathbb{E}[D \mid x, y]} = \mathbb{E}[A_x B_y \mid x, y, D = 1].$$

Proof Sketch. The indicator D selects a conditional ensemble. If D is independent of the hidden outcome pattern given the settings, then the selected estimator agrees with the raw one and the usual CHSH reasoning transfers. If D depends on outcomes or on a setting-outcome interaction,




Figure 2: Effective-rate context. Wang et al. report an effective retained-event rate near 10^{-18} [17], far below the threshold regime usually discussed when one wants detector-loophole-resistant Bell claims [5, 6]. Here η is being used only as a CHSH-style benchmark proxy for the retained sample. Once that benchmark falls low enough, the local post-selection ceiling climbs into super-classical territory without physical embarrassment.

the estimator has changed, and the unqualified inference from $S > 2$ to nonlocality has changed with it [3, 4, 9]. $\square$

That conditionalization issue is the detection loophole in statistical clothing. Pearle, Clauser-Horne, Garg-Mermin, Eberhard, and later reviews all stress that low-efficiency Bell tests need more than a selected scalar summary [3, 6, 4, 5, 9]. A useful CHSH benchmark for post-selected local models is

$$S_{\text{LHV,bench}}(\eta) = \frac{4}{\eta} - 2.$$

We use that expression throughout as the working CHSH benchmark. Appendix A states the scope caveat explicitly: it is informative for CHSH-style post-selected analysis, not a universal replacement for every loophole discussion ever written.

Proposition 4 (Constructive Inflation by Outcome-Coupled Acceptance). *Let $d_{00} = d_{01} = d_{10} = +1$ and $d_{11} = -1$. If a trial with setting pair (x, y) is kept with probability p_{hi} when $A_x B_y = d_{xy}$ and with probability p_{lo} otherwise, then*

$$E_{xy}^{(\text{sel})} = \frac{(p_{\text{hi}} + p_{\text{lo}})E_{xy} + d_{xy}(p_{\text{hi}} - p_{\text{lo}})}{(p_{\text{hi}} + p_{\text{lo}}) + d_{xy}E_{xy}(p_{\text{hi}} - p_{\text{lo}})}.$$

For the deterministic model used here, the raw correlations are $E_{00} = E_{01} = E_{10} = 1/2$ and $E_{11} = -1/2$. Setting $p_{\text{hi}} = 0.9$ and $p_{\text{lo}} = 0.1$ yields the exact toy-model expectations $E_{xy}^{(\text{sel})} = d_{xy} \cdot 13/14$ and therefore $S_{\text{sel}} = 26/7 \approx 3.714$.

Proof Sketch. Write $q_{xy} = P(A_x B_y = d_{xy} \mid x, y)$. Then $q_{xy} = (1 + d_{xy}E_{xy})/2$, the acceptance rate is $\eta_{xy} = p_{\text{hi}}q_{xy} + p_{\text{lo}}(1 - q_{xy})$, and the selected correlation is

$$d_{xy} [p_{\text{hi}}q_{xy} - p_{\text{lo}}(1 - q_{xy})] / \eta_{xy}.$$

Appendix B expands the algebra. The important structural fact is plain: once the filter rewards the sign pattern that maximizes the CHSH combination, the statistic follows the filter. $\square$

The toy model is intentionally simple. That simplicity is useful: if a minimal outcome-coupled filter already produces a super-classical selected statistic, stronger diagnostics are required before local models can be excluded.

The broader review literature on Bell nonlocality places the present argument in context. Bell inequalities are exact consequences of structural assumptions [8]. Experimental Bell claims therefore require not only a large statistic but also an explicit analysis of the assumptions under which that statistic is meaningful.

3 Methods

3.1 Deterministic Local Source

The paper’s Monte Carlo benchmark uses the reference-model workflow from the companion simulator repository. A hidden variable λ is drawn uniformly on $[0, 2\pi)$, and the outcomes are deterministic thresholded cosines:

$$A_x(\lambda) = \text{sign}[\cos(a_x - \lambda)], \quad B_y(\lambda) = \text{sign}[\cos(b_y - \lambda)].$$

The chosen CHSH settings are $(a_0, a_1, b_0, b_1) = (0, \pi/2, \pi/4, -\pi/4)$. For this deterministic sign model, those angles give the exact raw expectations

$$E_{00} = E_{01} = E_{10} = \frac{1}{2}, \quad E_{11} = -\frac{1}{2}, \quad S_{\text{raw}} = 2.$$

The Monte Carlo therefore verifies a simple local baseline rather than searching for one. This is appropriate because the argument requires an explicit counterexample, not an optimized model.

3.2 Outcome-Coupled Acceptance Rule

The filter uses the sign pattern that maximizes the chosen CHSH combination: $d_{00} = d_{01} = d_{10} = +1$ and $d_{11} = -1$. For each trial, define an acceptance indicator D by

$$P(D = 1 \mid A_x, B_y, x, y) = \begin{cases} p_{\text{hi}}, & A_x B_y = d_{xy}, \\ p_{\text{lo}}, & A_x B_y \neq d_{xy}. \end{cases}$$

with $p_{\text{hi}} = 0.9$ and $p_{\text{lo}} = 0.1$. The filter is local in the sense that it is applied after outcomes have been generated from a local source, but it is not fair: it depends directly on the outcome pattern and indirectly on the setting pair through d_{xy} .

3.3 Reproducibility Protocol

1. Sample a setting pair (x, y) uniformly from the four CHSH combinations.
2. Sample λ uniformly on $[0, 2\pi)$ and compute $A_x(\lambda)$ and $B_y(\lambda)$.
3. Record the raw product $A_x B_y$ and the target sign d_{xy} .
4. Accept the trial with probability p_{hi} when $A_x B_y = d_{xy}$, else with probability p_{lo} .
5. Estimate both raw and selected correlations by setting pair, then assemble the CHSH statistic from those four entries.
6. Repeat for 20 runs and report the mean, run-level spread, and 95% t -interval.

The protocol is intentionally compact so that the origin of the inflation remains transparent.

Table 2: Table 1. Parameters for the Monte Carlo rebuttal workflow.

Parameter	Value	Role
Hidden variable	$\lambda \sim \text{Unif}[0, 2\pi)$	Provides an isotropic local source with deterministic outcomes.
CHSH settings	$(0, \pi/2, \pi/4, -\pi/4)$	Produces the exact raw sign pattern $(+1/2, +1/2, +1/2, -1/2)$.
Runs	20 independent runs	Supports a run-level confidence interval for the aggregate Monte Carlo result.
Trials per run	50,000	Samples settings uniformly across the four CHSH pairs.
Selection parameters	$p_{\text{hi}} = 0.9, p_{\text{lo}} = 0.1$	Rewards the CHSH-favored sign pattern and penalizes its complement.
Interval estimate	95% t -interval over runs	Summarizes Monte Carlo variability of the aggregate result row.

Table 3: Table 2. Comparison of the reported experiment with the rebuttal model.

Dataset	CHSH S	Acceptance Benchmark Rate	/ Interpretation
Classical model, no selection	2.001 ± 0.009	100%	Respects the local CHSH bound, as a fair local sample should.
Classical model, post-selected	3.716 ± 0.004	70.0%	Selected statistic exceeds both the classical and Tsirelson bounds.
Wang et al. (2025)	2.275 ± 0.057	about 10^{-18} effective acceptance	Lives deep inside a regime where local post-selection remains available.

4 Results

4.1 Aggregate Monte Carlo Result

The Monte Carlo rebuttal retains the project’s published aggregate result: without post-selection, the local model remains at $S = 2.001$ (95% CI $[1.998, 2.004]$) with 100% acceptance, while the same source under outcome-coupled acceptance yields $S = 3.716$ (95% CI $[3.714, 3.717]$) at 70.0% acceptance. The comparison matters because a local filter already reaches the evidentiary class being interpreted as nonlocal.

4.2 Setting-Pair Decomposition of the Toy Filter

The analytic toy model makes the inflation mechanism numerically transparent. Starting from the exact raw correlations $(+1/2, +1/2, +1/2, -1/2)$, the acceptance rule with $p_{\text{hi}} = 0.9$ and $p_{\text{lo}} = 0.1$ yields the exact selected correlations $(\pm 13/14)$ and a uniform pairwise acceptance rate of 0.7.

Summing the exact selected entries gives

$$S_{\text{sel}} = 13/14 + 13/14 + 13/14 - (-13/14) = 26/7 \approx 3.714,$$

Table 4: Table 3. Exact toy-model expectations by CHSH setting pair.

Setting pair	Favored d_{xy}	sign	Raw E_{xy}	Selected $E_{xy}^{(\text{sel})}$	Acceptance η_{xy}
(a, b)	+1		$1/2 = 0.500$	$13/14 = 0.929$	$7/10 = 0.700$
(a, b')	+1		$1/2 = 0.500$	$13/14 = 0.929$	$7/10 = 0.700$
(a', b)	+1		$1/2 = 0.500$	$13/14 = 0.929$	$7/10 = 0.700$
(a', b')	-1		$-1/2 = -0.500$	$-13/14 = -0.929$	$7/10 = 0.700$

which is fully consistent with the Monte Carlo result row. The analytic toy filter and the numerical rebuttal reach the same conclusion through different formulations.

4.3 Non-Diagnostic Status of the Wang et al. Statistic

Wang et al. report a smaller value than the benchmark above, but that comparison is incidental. The decisive fact is that once the accepted sample is permitted to drift away from the emitted ensemble, local models can already populate the reported regime. At an effective retained-event rate of 10^{-18} —used here only as a CHSH-style benchmark proxy η —the working bound $S_{\text{LHV,bench}}(\eta) = 4/\eta - 2$ is astronomically above 2.275. In that setting, a super-classical selected value is not yet a proof of nonlocality; it is a prompt to inspect the conditioning.

The purpose of the rebuttal is limited. It is enough to show that the reported statistic is not yet exclusive of local explanations.

5 Discussion

5.1 What the Model Proves

A deterministic local source paired with an outcome-coupled acceptance rule can generate a selected CHSH value above both 2 and the Tsirelson bound. A selected scalar alone therefore does not certify nonlocality in a regime where such filters remain experimentally admissible. The conclusion is conditional, but the relevant condition is the one under dispute in the Wang et al. interpretation.

5.2 What the Model Does Not Prove

It does not prove that Wang et al.’s apparatus is classical. It does not deny that a quantum description of the apparatus may be appropriate—the path-identity interference mechanism they exploit is real and interesting in its own right [11, 12, 13]. It does not claim that $4/\eta - 2$ is the last word on every Bell inequality or every experimental design. It shows only that the reported evidence does not yet force the nonlocal interpretation, a conclusion that is also supported by the immediate postpublication response literature [18, 19, 20].

5.3 What the Published Critiques Add

The critique literature goes further than the present formal thesis. Wharton and Price reproduce the geometry with a fully local classical model and interpret the acceptance rule as a collider-bias mechanism. Wójcik and Wójcik trace the apparent nonlocality to post-selection plus the unconventional normalization, arguing that the phases do not play the operational role of standard Bell settings. Cieřliński and coauthors prove that when the local phases are treated as the settings,




Figure 3: The exact local counterexample. At $p_{lo} = 0.10$, the model yields $S = 26/7$ with 70.0% acceptance while the underlying source remains local and classical before selection.

the observed interference admits a local realistic model, even though a different on-off protocol can still reveal genuine Bell nonclassicality in the same optical platform.

This paper adopts that convergence as context while keeping the thesis narrower. A local counterexample already suffices to show that the reported evidence is not yet exclusive. That narrower stance is also consistent with Wang et al.’s own caveats: the locality loophole remains open, the sampling loophole remains open, and the interpretation relies on fair sampling [17].

5.4 What Would Weaken This Rebuttal

The rebuttal would lose force if Wang et al. supplied diagnostics that make the relevant local-selection mechanism implausible. In practical terms, that means setting-pair-resolved acceptance rates, raw pre-selection correlations, no-signaling checks on both raw and selected data, and explicit tests showing that acceptance is not correlated with the hidden outcome structure or with the measurement context. If the selected sample can be shown to be fair, this paper’s central objection recedes accordingly.

The benchmark for persuasive Bell evidence is set by the loophole-free tests of 2015. Hensen et al. used an event-ready architecture, while Giustina et al. and Shalm et al. used entangled-photon tests analyzed with CH/Eberhard-style logic rather than a bare CHSH threshold. All three paired suitable efficiencies with locality protections and explicit diagnostics rather than relying on a single post-selected number [14, 15, 16].




Figure 4: Selection inflates the correlators, not the underlying locality class. Raw correlators of magnitude $1/2$ become retained correlators of magnitude $13/14$ at the paper point. The apparent Bell excess is generated in the conditioning step.

5.5 Inference Discipline

A Bell test is an inference pipeline, not a single scalar. When efficiency is so low that the accepted sample may comprise nearly all observable events, the burden shifts to showing that the observed boundary crossing is attributable to physics rather than to conditioning. Recent work on the physical significance of Bell non-locality underscores that broader debate [21], but that debate begins only after the estimator itself has been disciplined.

A quantum description of the apparatus may still be appropriate. The present point is narrower: exclusivity has not yet been established by the reported evidence.

6 Conclusion

A local hidden-variable source with outcome-coupled acceptance reproduces the evidentiary regime of a post-selected Bell claim and can exceed the Tsirelson bound without entanglement. In that setting, the Wang et al. statistic does not by itself establish quantum nonlocality.

The published critique literature now goes further and largely attributes the reported violation to fourfold post-selection and unconventional normalization. This paper treats that stronger diagnosis as context while retaining the narrower evidentiary thesis.

The remedy is methodological: report the raw correlations, the setting-resolved acceptance structure, and the diagnostics that constrain fair sampling. Until then, the appropriate interpretation of a low-efficiency post-selected violation is evidentiary caution.

Table 5: Table 4. Minimum reporting package for a persuasive low-efficiency Bell claim.

Diagnostic	Why it matters
Raw correlations before selection	Shows whether the effect exists prior to the filtering step.
Acceptance rate by setting pair	Tests whether retention is coupled to measurement context.
No-signaling checks on raw and selected data	Detects filter-induced distortions in the observed marginals.
Independence tests for acceptance vs. outcomes/settings	Directly probes the fair-sampling assumption instead of merely hoping for it.
Sensitivity analysis for plausible filter bias	Quantifies how much apparent violation could be manufactured by asymmetric retention.

7 Appendix

7.1 Appendix A. Detection Benchmark and Threshold

The working benchmark used in this paper is

$$S_{\text{LHV,bench}}(\eta) = \frac{4}{\eta} - 2.$$

Setting that benchmark equal to the quantum ceiling yields the familiar CHSH threshold:

$$\frac{4}{\eta_c} - 2 = 2\sqrt{2} \implies \eta_c = \frac{2}{1 + \sqrt{2}} \approx 0.828.$$

This threshold is the standard reference point for CHSH-style fair-sampling discussions, consistent with the Pearle, Clauser-Horne, Garg-Mermin, and Eberhard literature [3, 6, 4, 5]. It is not universal across every Bell inequality variant: Clauser-Horne and Eberhard-type analyses can shift the efficiency discussion, especially for non-maximally entangled states. In this paper, η is being used as a compact CHSH benchmark parameter for the retained-sample rate, because Wang et al. frame their claim as a CHSH-style Bell violation even though their experimental rate is better described as effective acceptance than as ordinary detector efficiency.

7.2 Appendix B. Derivation of the Toy Post-Selection Estimator

Fix a setting pair (x, y) and favored sign $d_{xy} \in \{-1, +1\}$. Let $q_{xy} = P(A_x B_y = d_{xy} \mid x, y)$. Then

$$q_{xy} = \frac{1 + d_{xy} E_{xy}}{2}$$

because $E_{xy} = P(AB = +1 \mid x, y) - P(AB = -1 \mid x, y)$. Under the acceptance rule p_{hi} for favored outcomes and p_{lo} otherwise, the pairwise acceptance rate is

$$\eta_{xy} = p_{\text{hi}} q_{xy} + p_{\text{lo}} (1 - q_{xy}) = \frac{(p_{\text{hi}} + p_{\text{lo}}) + d_{xy} E_{xy} (p_{\text{hi}} - p_{\text{lo}})}{2}$$

and the selected correlation is

$$E_{xy}^{(\text{sel})} = d_{xy} \frac{p_{\text{hi}} q_{xy} - p_{\text{lo}} (1 - q_{xy})}{p_{\text{hi}} q_{xy} + p_{\text{lo}} (1 - q_{xy})} = \frac{(p_{\text{hi}} + p_{\text{lo}}) E_{xy} + d_{xy} (p_{\text{hi}} - p_{\text{lo}})}{(p_{\text{hi}} + p_{\text{lo}}) + d_{xy} E_{xy} (p_{\text{hi}} - p_{\text{lo}})}.$$

Table 6: Table 5. Symbols and conventions used in the paper.

Symbol	Meaning
x, y	Alice/Bob setting labels in $\{0, 1\}$.
A_x, B_y	Binary outcomes taking values ± 1 .
λ	Hidden variable sampled from a uniform local source.
D	Acceptance indicator for whether a trial is retained.
E_{xy}	Raw correlation $\mathbb{E}[A_x B_y \mid x, y]$.
$E_{xy}^{(\text{sel})}$	Selected correlation conditioned on $D = 1$.
d_{xy}	Favored product sign for the CHSH-maximizing filter.
$S = E_{00} + E_{01} + E_{10} - E_{11} $	The CHSH sign convention used throughout; other conventions are equivalent up to relabeling.

Substituting the exact raw values $E_{00} = E_{01} = E_{10} = 1/2$, $E_{11} = -1/2$, and $(p_{\text{hi}}, p_{\text{lo}}) = (0.9, 0.1)$ gives

$$\eta_{xy} = \frac{1 + 0.8 \cdot \frac{1}{2}}{2} = 0.7, \quad E_{xy}^{(\text{sel})} = d_{xy} \frac{1.0 \cdot \frac{1}{2} + 0.8}{1.0 + 0.8 \cdot \frac{1}{2}} = d_{xy} \frac{13}{14}.$$

Hence $S_{\text{sel}} = 26/7 \approx 3.714$. More generally, whenever the filter rewards the sign pattern aligned with the chosen CHSH combination, the statistic inherits the filter’s preferences.

7.3 Appendix C. Sign Conventions and Notation Glossary

8 References

References

- [1] J. S. Bell, “On the Einstein Podolsky Rosen paradox,” *Physics Physique Fizika* 1, 195–200 (1964). <https://doi.org/10.1103/PhysicsPhysiqueFizika.1.195>
- [2] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, “Proposed experiment to test local hidden-variable theories,” *Physical Review Letters* 23, 880–884 (1969). <https://doi.org/10.1103/PhysRevLett.23.880>
- [3] P. M. Pearle, “Hidden-variable example based upon data rejection,” *Physical Review D* 2, 1418–1425 (1970). <https://doi.org/10.1103/PhysRevD.2.1418>
- [4] J. F. Clauser and M. A. Horne, “Experimental consequences of objective local theories,” *Physical Review D* 10, 526–535 (1974). <https://doi.org/10.1103/PhysRevD.10.526>
- [5] A. Garg and N. D. Mermin, “Detector inefficiencies in the Einstein-Podolsky-Rosen experiment,” *Physical Review D* 35, 3831–3835 (1987). <https://doi.org/10.1103/PhysRevD.35.3831>
- [6] P. H. Eberhard, “Background level and counter efficiencies required for a loophole-free Einstein-Podolsky-Rosen experiment,” *Physical Review A* 47, R747–R750 (1993). <https://doi.org/10.1103/PhysRevA.47.R747>
- [7] A. Fine, “Hidden Variables, Joint Probability, and the Bell Inequalities,” *Physical Review Letters* 48, 291–295 (1982). <https://doi.org/10.1103/PhysRevLett.48.291>

- [8] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, “Bell nonlocality,” *Reviews of Modern Physics* 86, 419–478 (2014). <https://doi.org/10.1103/RevModPhys.86.419>
- [9] J.-Å. Larsson, “Loopholes in Bell inequality tests of local realism,” *Journal of Physics A* 47, 424003 (2014). <https://doi.org/10.1088/1751-8113/47/42/424003>
- [10] B. S. Tsirelson, “Quantum generalizations of Bell’s inequality,” *Letters in Mathematical Physics* 4, 93–98 (1980). <https://doi.org/10.1007/BF00417500>
- [11] X. Y. Zou, L. J. Wang, and L. Mandel, “Induced coherence and indistinguishability in optical interference,” *Physical Review Letters* 67, 318–321 (1991). <https://doi.org/10.1103/PhysRevLett.67.318>
- [12] M. Krenn, M. Malik, T. Scheidl, R. Ursin, and A. Zeilinger, “Entanglement by path identity,” *Physical Review Letters* 118, 080401 (2017). <https://doi.org/10.1103/PhysRevLett.118.080401>
- [13] A. Hochrainer, M. Lahiri, and A. Zeilinger, “Quantum indistinguishability by path identity and with undetected photons,” *Reviews of Modern Physics* 94, 025007 (2022). <https://doi.org/10.1103/RevModPhys.94.025007>
- [14] B. Hensen et al., “Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres,” *Nature* 526, 682–686 (2015). <https://doi.org/10.1038/nature15759>
- [15] M. Giustina et al., “Significant-loophole-free test of Bell’s theorem with entangled photons,” *Physical Review Letters* 115, 250401 (2015). <https://doi.org/10.1103/PhysRevLett.115.250401>
- [16] L. K. Shalm et al., “Strong loophole-free test of local realism,” *Physical Review Letters* 115, 250402 (2015). <https://doi.org/10.1103/PhysRevLett.115.250402>
- [17] K. Wang et al., “Violation of Bell inequality with unentangled photons,” *Science Advances* 11(31), eadr1794 (2025). <https://doi.org/10.1126/sciadv.adr1794>
- [18] K. B. Wharton and H. Price, “Bell Inequality Violations Without Entanglement? It’s Just Postselection,” arXiv:2508.13431 (2025). <https://arxiv.org/abs/2508.13431>
- [19] A. Wójcik and J. Wójcik, “Simple explanation of apparent Bell nonlocality of unentangled photons,” arXiv:2509.03127 (2025). <https://arxiv.org/abs/2509.03127>
- [20] P. Cieřliński, J.-Å. Larsson, M. Markiewicz, K. Schlichtholz, and M. Żukowski, “Unquestionable Bell theorem for interwoven frustrated down conversion processes,” *Physical Review Letters* 136, 090206 (2026); arXiv:2508.19207. <https://arxiv.org/abs/2508.19207>
- [21] C. Vieira, R. Ramanathan, and A. Cabello, “Test of the physical significance of Bell non-locality,” *Nature Communications* 16, 4390 (2025). <https://www.nature.com/articles/s41467-025-59247-7>