

Bell Violations Without Entanglement?

A Reproducible Rebuttal of Wang et al. (2025)

Meshal Alawein
UC Berkeley
contact@meshal.ai

Version 1.3
March 30, 2026

Abstract

A recent *Science Advances* paper reports $S = 2.275 \pm 0.057$ at an effective detection efficiency of roughly 10^{-18} and interprets that statistic as Bell-inequality violation without entanglement [17]. The present paper makes a narrower claim. In such a severely filtered regime, the scalar fact $S > 2$ is not exclusionary evidence against local models unless the selection mechanism is independently constrained. We formalize this point in three steps. First, we write the selected estimator as a conditional expectation on accepted events and identify fair sampling as the assumption that transfers the classical CHSH bound from the emitted ensemble to the retained one. Second, we construct a deterministic local hidden-variable model with an outcome-coupled acceptance rule and derive the exact toy-model expectations $E_{\text{sel}} = \pm 13/14$ and $S_{\text{sel}} = 26/7 \approx 3.714$. Third, we compare those exact expectations with the site’s paper-grade Monte Carlo artifact, which yields $S = 3.716$ (95% CI [3.714, 3.717]) at **70.0%** acceptance, and with the Wang et al. report. The evidentiary consequence is limited but decisive: in an ultra-low-efficiency Bell test, a post-selected CHSH value above 2 is suggestive at most until raw correlations, setting-resolved acceptance rates, and fair-sampling diagnostics are reported.

Claim. In the ultra-low-efficiency regime, a post-selected CHSH value above 2 is non-exclusionary evidence until the selection rule itself is constrained.

Published findings.

- The paper-grade Monte Carlo artifact remains $S = 3.716$ (95% CI [3.714, 3.717]) at **70.0%** acceptance for a purely local source.
- The selected estimator is a conditional expectation on accepted events, so the naive $S \leq 2$ inference survives only under fair-sampling assumptions.
- At Wang et al.’s reported efficiency of about 10^{-18} , diagnostic structure matters more than rhetorical velocity.

1 Introduction

Bell’s theorem excludes local hidden-variable explanations for the full set of quantum correlations under the usual assumptions of locality, realism, and setting independence [1]. The Clauser-Horne-Shimony-Holt inequality packages that exclusion into an experimentally testable scalar, and

Table 1: Literature checkpoints relevant to the present claim.

Cluster	What it establishes	Why it matters here
Bell 1964; CHSH 1969 [1, 2]	Local hidden-variable models satisfy a bound on the full correlation structure.	The theorem targets the emitted ensemble, not an arbitrary filtered subset.
Pearle through Eberhard [3, 4, 5, 6]	Selection and efficiency can open a detection loophole unless explicitly controlled.	Wang et al. operate far inside the regime where that warning is operationally important.
Path identity optics [11, 12, 13]	Indistinguishable generation pathways can create striking interference effects.	The optical novelty is not being denied; the Bell inference is the narrower issue.
Loophole-aware Bell benchmarks [14, 15, 16]	Strong Bell claims are paired with explicit closure strategies, not just large S .	That is the comparison class for any foundational headline.
Immediate 2025 responses [18, 19, 20]	The postpublication debate quickly focused on postselection and interpretation.	The present paper joins that narrower, more technical conversation.

Fine sharpened the probabilistic picture by relating CHSH satisfaction to the existence of a joint distribution over the counterfactual outcomes [2, 7].

$$S = |E_{00} + E_{01} + E_{10} - E_{11}|$$

In the fair-sampling regime, the local bound is $S \leq 2$ and the quantum ceiling is the Tsirelson bound $2\sqrt{2}$ [10]. Wang et al. report a super-classical value from a multiphoton path-identity interference experiment and interpret it as Bell-inequality violation without entanglement [17].

The question here is therefore not whether the apparatus is sophisticated. It plainly is. The question is whether a post-selected statistic in an ultra-low-efficiency regime can carry the inferential weight being assigned to it.

This paper answers that question with a constructive counterexample and a narrower standard of evidence. A large CHSH number can be impressive; impressiveness is not, by itself, a loophole-closure criterion.

The claim is about inference, not ontology. A local counterexample need only show that the reported evidence class remains available to local explanations.

2 Theoretical Background

2.1 Notation and Assumptions

We write Alice’s setting as $x \in \{0, 1\}$ and Bob’s as $y \in \{0, 1\}$. Outcomes are binary: $A_x, B_y \in \{-1, +1\}$. A hidden variable is denoted by λ , and $D \in \{0, 1\}$ marks whether a trial is retained by the analysis filter.

Locality $A_x(\lambda)$ depends on x and λ , not on y ; symmetrically for Bob.




Figure 1: Reference scales for the two papers. Wang et al.’s reported value sits above the classical bound, but the exact local counterexample sits even higher. The figure is not an empirical fit; it is a warning about what a filtered estimator can and cannot prove.

Realism All four counterfactual outcomes are taken to exist simultaneously for the local model under study.

Freedom $P(\lambda | x, y) = P(\lambda)$, so the settings are statistically independent of the hidden variable.

Fair Sampling The standard CHSH inference on selected data additionally requires D not to depend on the hidden outcome pattern in a biasing way.

Definition (CHSH Functional and Sign Convention). *In the convention used throughout this paper, the CHSH functional is*

$$S = |E_{00} + E_{01} + E_{10} - E_{11}|, \quad E_{xy} = \mathbb{E}[A_x B_y | x, y].$$

Relabeling settings or flipping one party’s output signs maps this expression to the other familiar CHSH sign placements without changing the attainable classical or quantum bounds. The notation changes; the admissible region does not.

Proposition 1 (Fair-Sampling Local Bound). *For any local hidden-variable model satisfying locality, realism, freedom, and fair sampling, the selected CHSH statistic obeys*

$$S_{\text{LHV}} \leq 2.$$

Proof Sketch. For each fixed λ , define

$$X(\lambda) = A_0(\lambda)[B_0(\lambda) + B_1(\lambda)] + A_1(\lambda)[B_0(\lambda) - B_1(\lambda)].$$

Because Bob’s outcomes are each ± 1 , exactly one bracket has magnitude 2 and the other vanishes, so $|X(\lambda)| \leq 2$. Integrating over λ preserves the bound. If the accepted sample is a fair image of the emitted ensemble, the same algebra applies to the retained data as well [1, 2, 7]. $\square$

Proposition 2 (Quantum Comparison Ceiling). *Within quantum mechanics, the maximal CHSH value is the Tsirelson bound*

$$S_{\text{QM}} \leq 2\sqrt{2} \approx 2.828.$$

Proof Sketch. Let A, A' and B, B' be Hermitian observables with eigenvalues ± 1 , and define the CHSH operator

$$\hat{S} = A \otimes (B + B') + A' \otimes (B - B').$$

The standard commutator calculation gives $\hat{S}^2 = 4I - [A, A'] \otimes [B, B']$, from which $\|\hat{S}^2\| \leq 8$ and therefore $\|\hat{S}\| \leq 2\sqrt{2}$. That ceiling is achievable for suitable entangled states and local observables [10]. $\square$

Proposition 3 (Selection Replaces the Unconditional Estimator). *Once analysis is conditioned on accepted trials, the operational correlation is not E_{xy} but*

$$E_{xy}^{(\text{sel})} = \frac{\mathbb{E}[A_x B_y D \mid x, y]}{\mathbb{E}[D \mid x, y]} = \mathbb{E}[A_x B_y \mid x, y, D = 1].$$

Proof Sketch. The indicator D selects a conditional ensemble. If D is independent of the hidden outcome pattern given the settings, then the selected estimator agrees with the raw one and the usual CHSH reasoning transfers. If D depends on outcomes or on a setting-outcome interaction, the estimator has changed, and the unqualified inference from $S > 2$ to nonlocality has changed with it [3, 4, 9]. $\square$

That conditionalization issue is the detection loophole in statistical clothing. Pearle, Clauser-Horne, Garg-Mermin, Eberhard, and later reviews all stress that low-efficiency Bell tests need more than a selected scalar summary [3, 6, 4, 5, 9]. A useful CHSH benchmark for post-selected local models is

$$S_{\text{LHV,bench}}(\eta) = \frac{4}{\eta} - 2.$$

We use that expression throughout as the working efficiency benchmark. Appendix A states the scope caveat explicitly: it is informative for CHSH-style post-selected analysis, not a universal replacement for every loophole discussion ever written.

Proposition 4 (Constructive Inflation by Outcome-Coupled Acceptance). *Let $d_{00} = d_{01} = d_{10} = +1$ and $d_{11} = -1$. If a trial with setting pair (x, y) is kept with probability p_{hi} when $A_x B_y = d_{xy}$ and with probability p_{lo} otherwise, then*

$$E_{xy}^{(\text{sel})} = \frac{(p_{\text{hi}} + p_{\text{lo}})E_{xy} + d_{xy}(p_{\text{hi}} - p_{\text{lo}})}{(p_{\text{hi}} + p_{\text{lo}}) + d_{xy}E_{xy}(p_{\text{hi}} - p_{\text{lo}})}.$$

For the deterministic model used here, the raw correlations are $E_{00} = E_{01} = E_{10} = 1/2$ and $E_{11} = -1/2$. Setting $p_{\text{hi}} = 0.9$ and $p_{\text{lo}} = 0.1$ yields the exact toy-model expectations $E_{xy}^{(\text{sel})} = d_{xy} \cdot 13/14$ and therefore $S_{\text{sel}} = 26/7 \approx 3.714$.

Proof Sketch. Write $q_{xy} = P(A_x B_y = d_{xy} \mid x, y)$. Then $q_{xy} = (1 + d_{xy}E_{xy})/2$, the acceptance rate is $\eta_{xy} = p_{\text{hi}}q_{xy} + p_{\text{lo}}(1 - q_{xy})$, and the selected correlation is

$$d_{xy} [p_{\text{hi}}q_{xy} - p_{\text{lo}}(1 - q_{xy})] / \eta_{xy}.$$

Appendix B expands the algebra. The important structural fact is plain: once the filter rewards the sign pattern that maximizes the CHSH combination, the statistic follows the filter. $\square$




Figure 2: Efficiency context. Wang et al. report an effective efficiency near 10^{-18} [17], far below the threshold regime usually discussed when one wants detector-loophole-resistant Bell claims [5, 6]. Once efficiency falls below the CHSH benchmark threshold, the local post-selection ceiling climbs into super-classical territory without physical embarrassment.

The toy model is intentionally plain. If a plain filter can do the job, one should not demand baroque metaphysics before demanding better diagnostics.

The broader review literature on Bell nonlocality is useful here because it clarifies both sides at once. On one side, Bell inequalities are not vague metaphors; they are exact consequences of structural assumptions [8]. On the other side, experimental Bell claims are not established by enthusiasm alone. They are established by pairing a large statistic with a careful analysis of the assumptions under which the statistic carries that meaning.

3 Methods

3.1 Deterministic Local Source

The paper-grade artifact uses the reference-model workflow from the companion simulator repository. A hidden variable λ is drawn uniformly on $[0, 2\pi)$, and the outcomes are deterministic thresholded cosines:

$$A_x(\lambda) = \text{sign}[\cos(a_x - \lambda)], \quad B_y(\lambda) = \text{sign}[\cos(b_y - \lambda)].$$

The chosen CHSH settings are $(a_0, a_1, b_0, b_1) = (0, \pi/2, \pi/4, -\pi/4)$. For this deterministic sign model, those angles give the exact raw expectations

$$E_{00} = E_{01} = E_{10} = \frac{1}{2}, \quad E_{11} = -\frac{1}{2}, \quad S_{\text{raw}} = 2.$$

The Monte Carlo therefore verifies a simple local baseline rather than searching for one. That is a feature, not a concession.

Table 2: Table 1. Parameters for the paper-grade rebuttal workflow.

Parameter	Value	Role
Hidden variable	$\lambda \sim \text{Unif}[0, 2\pi)$	Provides an isotropic local source with deterministic outcomes.
CHSH settings	$(0, \pi/2, \pi/4, -\pi/4)$	Produces the exact raw sign pattern $(+1/2, +1/2, +1/2, -1/2)$.
Runs	20 independent runs	Supports a run-level confidence interval for the aggregate Monte Carlo result.
Trials per run	50,000	Samples settings uniformly across the four CHSH pairs.
Selection parameters	$p_{\text{hi}} = 0.9, p_{\text{lo}} = 0.1$	Rewards the CHSH-favored sign pattern and penalizes its complement.
Interval estimate	95% t -interval over runs	Summarizes Monte Carlo variability of the aggregate artifact row.

3.2 Outcome-Coupled Acceptance Rule

The filter uses the sign pattern that maximizes the chosen CHSH combination: $d_{00} = d_{01} = d_{10} = +1$ and $d_{11} = -1$. For each trial, define an acceptance indicator D by

$$P(D = 1 \mid A_x, B_y, x, y) = \begin{cases} p_{\text{hi}}, & A_x B_y = d_{xy}, \\ p_{\text{lo}}, & A_x B_y \neq d_{xy}. \end{cases}$$

with $p_{\text{hi}} = 0.9$ and $p_{\text{lo}} = 0.1$. The filter is local in the sense that it is applied after outcomes have been generated from a local source, but it is not fair: it depends directly on the outcome pattern and indirectly on the setting pair through d_{xy} .

3.3 Reproducibility Protocol

1. Sample a setting pair (x, y) uniformly from the four CHSH combinations.
2. Sample λ uniformly on $[0, 2\pi)$ and compute $A_x(\lambda)$ and $B_y(\lambda)$.
3. Record the raw product $A_x B_y$ and the target sign d_{xy} .
4. Accept the trial with probability p_{hi} when $A_x B_y = d_{xy}$, else with probability p_{lo} .
5. Estimate both raw and selected correlations by setting pair, then assemble the CHSH statistic from those four entries.
6. Repeat for 20 runs and report the mean, run-level spread, and 95% t -interval.

The algorithm is short because the mechanism is short. Selection bias rarely requires a symphony.

4 Results

4.1 Aggregate Monte Carlo Result

The paper-grade rebuttal keeps the project’s published aggregate row: without post-selection the local model remains at $S = 2.001$ (95% CI $[1.998, 2.004]$) with 100% acceptance, while the

Table 3: Table 2. Comparison of the reported experiment with the rebuttal model.

Dataset	CHSH S	Acceptance / Efficiency	Interpretation
Classical model, no selection	2.001 ± 0.009	100%	Respects the local CHSH bound, as a fair local sample should.
Classical model, post-selected	3.716 ± 0.004	70.0%	Selection artifact exceeds both the classical and Tsirelson bounds.
Wang et al. (2025)	2.275 ± 0.057	about 10^{-18}	Lives deep inside a regime where local post-selection remains available.

Table 4: Table 3. Exact toy-model expectations by CHSH setting pair.

Setting pair	Favored d_{xy}	sign	Raw E_{xy}	Selected $E_{xy}^{(\text{sel})}$	Acceptance η_{xy}
(a, b)	+1		$1/2 = 0.500$	$13/14 = 0.929$	$7/10 = 0.700$
(a, b')	+1		$1/2 = 0.500$	$13/14 = 0.929$	$7/10 = 0.700$
(a', b)	+1		$1/2 = 0.500$	$13/14 = 0.929$	$7/10 = 0.700$
(a', b')	-1		$-1/2 = -0.500$	$-13/14 = -0.929$	$7/10 = 0.700$

same source under outcome-coupled acceptance yields $S = 3.716$ (95% CI $[3.714, 3.717]$) at 70.0% acceptance. The evidentiary point is not that the artifact is larger than Wang et al.’s value. It is that a local filter reaches the same class of evidence being interpreted as nonlocal.

4.2 Setting-Pair Decomposition of the Toy Filter

The analytic toy model makes the inflation mechanism numerically transparent. Starting from the exact raw correlations $(+1/2, +1/2, +1/2, -1/2)$, the acceptance rule with $p_{\text{hi}} = 0.9$ and $p_{\text{lo}} = 0.1$ yields the exact selected correlations $(\pm 13/14)$ and a uniform pairwise acceptance rate of 0.7.

Summing the exact selected entries gives

$$S_{\text{sel}} = 13/14 + 13/14 + 13/14 - (-13/14) = 26/7 \approx 3.714,$$

which is fully consistent with the Monte Carlo artifact row. The analytic toy filter and the numerical rebuttal are telling the same story in different dialects.

4.3 Why Wang et al. Remain Non-Diagnostic Here

Wang et al. report a smaller value than the artifact above, but that comparison is incidental. The decisive fact is that once the accepted sample is permitted to drift away from the emitted ensemble, local models can already populate the reported regime. At $\eta \asymp 10^{-18}$, the working CHSH benchmark $S_{\text{LHV,bench}}(\eta) = 4/\eta - 2$ is astronomically above 2.275. In that setting, a super-classical selected value is not yet a proof of nonlocality; it is a prompt to inspect the conditioning.

The rebuttal need not outrun the reported statistic. It only needs to show that the reported statistic is not yet exclusive.




Figure 3: The exact local counterexample. At $p_{lo} = 0.10$, the model yields $S = 26/7$ with 70.0% acceptance while the underlying source remains local and classical before selection.

5 Discussion

5.1 What the Model Proves

The constructive result is limited but sharp. A deterministic local source, paired with an outcome-coupled acceptance rule, can generate a selected CHSH value above 2 and above the Tsirelson bound. Therefore a selected scalar by itself does not certify nonlocality in a regime where such filters remain experimentally admissible. The logic is conditional, but the condition is exactly the one that matters here.

5.2 What the Model Does Not Prove

It does not prove that Wang et al.’s apparatus is classical. It does not deny that a quantum description of the apparatus may be appropriate—the path-identity interference mechanism they exploit is real and interesting in its own right [11, 12, 13]. It does not claim that $4/\eta - 2$ is the last word on every Bell inequality or every experimental design. What it does say is simpler: the reported evidence does not yet force the nonlocal interpretation, and the immediate postpublication response literature concurs [18, 19, 20].

5.3 What Would Weaken This Rebuttal

The rebuttal would lose force if Wang et al. supplied diagnostics that make the relevant local-selection mechanism implausible. In practical terms, that means setting-pair-resolved acceptance rates, raw pre-selection correlations, no-signaling checks on both raw and selected data, and explicit tests showing that acceptance is not correlated with the hidden outcome structure or with the




Figure 4: Selection inflates the correlators, not the underlying locality class. Raw correlators of magnitude $1/2$ become retained correlators of magnitude $13/14$ at the paper point. The apparent Bell excess is generated in the conditioning step.

measurement context. If the selected sample can be shown to be fair, this paper’s central objection recedes accordingly.

The benchmark for persuasive Bell evidence is set by the loophole-free tests of 2015, which paired suitable efficiencies with locality protections and aggressive diagnostics rather than relying on a single post-selected number [14, 15, 16].

5.4 Inference Discipline

A Bell test is an inference pipeline, not a single scalar. When efficiency is so low that the accepted sample may be almost all one has, the burden shifts from celebrating a boundary crossing to demonstrating that the boundary was crossed by physics rather than by conditioning. The broader foundational discussion has not ended in 2015, as recent work on the physical significance of Bell non-locality makes clear [21]. But that larger debate begins only after the estimator has been disciplined. Nature may be subtle; statistics, when conditioned hard enough, are merely obedient.

*The strongest compatible objection is still: “perhaps the apparatus is genuinely quantum.”
Exactly so. Compatibility is the point. Exclusivity is what has not yet been shown.*

6 Conclusion

The conclusion is narrower than a manifesto and stronger than a complaint. A local hidden-variable source with outcome-coupled acceptance reproduces the evidentiary regime of a post-selected Bell claim and can exceed the Tsirelson bound without entanglement. The Wang et al. statistic,

Table 5: Table 4. Minimum reporting package for a persuasive low-efficiency Bell claim.

Diagnostic	Why it matters
Raw correlations before selection	Shows whether the effect exists prior to the filtering step.
Acceptance rate by setting pair	Tests whether retention is coupled to measurement context.
No-signaling checks on raw and selected data	Detects filter-induced distortions in the observed marginals.
Independence tests for acceptance vs. outcomes/settings	Directly probes the fair-sampling assumption instead of merely hoping for it.
Sensitivity analysis for plausible filter bias	Quantifies how much apparent violation could be manufactured by asymmetric retention.

considered as a selected CHSH number at ultra-low effective efficiency, therefore does not by itself establish quantum nonlocality.

The remedy is methodological rather than theatrical: report the raw correlations, the setting-resolved acceptance structure, and the diagnostics that constrain fair sampling. Until then, the correct interpretation of a low-efficiency post-selected violation is caution, not coronation.

7 Appendix

7.1 Appendix A. Detection Benchmark and Threshold

The working benchmark used in this paper is

$$S_{\text{LHV,bench}}(\eta) = \frac{4}{\eta} - 2.$$

Setting that benchmark equal to the quantum ceiling yields the familiar CHSH threshold:

$$\frac{4}{\eta_c} - 2 = 2\sqrt{2} \implies \eta_c = \frac{2}{1 + \sqrt{2}} \approx 0.828.$$

This threshold is the standard reference point for CHSH-style fair-sampling discussions, consistent with the Pearle, Clauser-Horne, Garg-Mermin, and Eberhard literature [3, 6, 4, 5]. It is not universal across every Bell inequality variant: Clauser-Horne and Eberhard-type analyses can shift the efficiency discussion, especially for non-maximally entangled states. The benchmark is used here because Wang et al. frame their claim as a CHSH-style Bell violation.

7.2 Appendix B. Derivation of the Toy Post-Selection Estimator

Fix a setting pair (x, y) and favored sign $d_{xy} \in \{-1, +1\}$. Let $q_{xy} = P(A_x B_y = d_{xy} \mid x, y)$. Then

$$q_{xy} = \frac{1 + d_{xy} E_{xy}}{2}$$

because $E_{xy} = P(AB = +1 \mid x, y) - P(AB = -1 \mid x, y)$. Under the acceptance rule p_{hi} for favored outcomes and p_{lo} otherwise, the pairwise acceptance rate is

$$\eta_{xy} = p_{\text{hi}} q_{xy} + p_{\text{lo}} (1 - q_{xy}) = \frac{(p_{\text{hi}} + p_{\text{lo}}) + d_{xy} E_{xy} (p_{\text{hi}} - p_{\text{lo}})}{2}$$

Table 6: Table 5. Symbols and conventions used in the paper.

Symbol	Meaning
x, y	Alice/Bob setting labels in $\{0, 1\}$.
A_x, B_y	Binary outcomes taking values ± 1 .
λ	Hidden variable sampled from a uniform local source.
D	Acceptance indicator for whether a trial is retained.
E_{xy}	Raw correlation $\mathbb{E}[A_x B_y \mid x, y]$.
$E_{xy}^{(\text{sel})}$	Selected correlation conditioned on $D = 1$.
d_{xy}	Favored product sign for the CHSH-maximizing filter.
$S = E_{00} + E_{01} + E_{10} - E_{11} $	The CHSH sign convention used throughout; other conventions are equivalent up to relabeling.

and the selected correlation is

$$E_{xy}^{(\text{sel})} = d_{xy} \frac{p_{\text{hi}} q_{xy} - p_{\text{lo}}(1 - q_{xy})}{p_{\text{hi}} q_{xy} + p_{\text{lo}}(1 - q_{xy})} = \frac{(p_{\text{hi}} + p_{\text{lo}})E_{xy} + d_{xy}(p_{\text{hi}} - p_{\text{lo}})}{(p_{\text{hi}} + p_{\text{lo}}) + d_{xy}E_{xy}(p_{\text{hi}} - p_{\text{lo}})}.$$

Substituting the exact raw values $E_{00} = E_{01} = E_{10} = 1/2$, $E_{11} = -1/2$, and $(p_{\text{hi}}, p_{\text{lo}}) = (0.9, 0.1)$ gives

$$\eta_{xy} = \frac{1 + 0.8 \cdot \frac{1}{2}}{2} = 0.7, \quad E_{xy}^{(\text{sel})} = d_{xy} \frac{1.0 \cdot \frac{1}{2} + 0.8}{1.0 + 0.8 \cdot \frac{1}{2}} = d_{xy} \frac{13}{14}.$$

Hence $S_{\text{sel}} = 26/7 \approx 3.714$. The stronger moral is structural: whenever the filter rewards the sign pattern aligned with the chosen CHSH combination, the statistic inherits the filter’s preferences.

7.3 Appendix C. Sign Conventions and Notation Glossary

8 References

References

- [1] J. S. Bell, “On the Einstein Podolsky Rosen paradox,” *Physics Physique Fizika* 1, 195–200 (1964). <https://doi.org/10.1103/PhysicsPhysiqueFizika.1.195>
- [2] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, “Proposed experiment to test local hidden-variable theories,” *Physical Review Letters* 23, 880–884 (1969). <https://doi.org/10.1103/PhysRevLett.23.880>
- [3] P. M. Pearle, “Hidden-variable example based upon data rejection,” *Physical Review D* 2, 1418–1425 (1970). <https://doi.org/10.1103/PhysRevD.2.1418>
- [4] J. F. Clauser and M. A. Horne, “Experimental consequences of objective local theories,” *Physical Review D* 10, 526–535 (1974). <https://doi.org/10.1103/PhysRevD.10.526>
- [5] A. Garg and N. D. Mermin, “Detector inefficiencies in the Einstein-Podolsky-Rosen experiment,” *Physical Review D* 35, 3831–3835 (1987). <https://doi.org/10.1103/PhysRevD.35.3831>
- [6] P. H. Eberhard, “Background level and counter efficiencies required for a loophole-free Einstein-Podolsky-Rosen experiment,” *Physical Review A* 47, R747–R750 (1993). <https://doi.org/10.1103/PhysRevA.47.R747>

- [7] A. Fine, “Hidden Variables, Joint Probability, and the Bell Inequalities,” *Physical Review Letters* 48, 291–295 (1982). <https://doi.org/10.1103/PhysRevLett.48.291>
- [8] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, “Bell nonlocality,” *Reviews of Modern Physics* 86, 419–478 (2014). <https://doi.org/10.1103/RevModPhys.86.419>
- [9] J.-Å. Larsson, “Loopholes in Bell inequality tests of local realism,” *Journal of Physics A* 47, 424003 (2014). <https://doi.org/10.1088/1751-8113/47/42/424003>
- [10] B. S. Tsirelson, “Quantum generalizations of Bell’s inequality,” *Letters in Mathematical Physics* 4, 93–98 (1980). <https://doi.org/10.1007/BF00417500>
- [11] X. Y. Zou, L. J. Wang, and L. Mandel, “Induced coherence and indistinguishability in optical interference,” *Physical Review Letters* 67, 318–321 (1991). <https://doi.org/10.1103/PhysRevLett.67.318>
- [12] M. Krenn, M. Malik, T. Scheidl, R. Ursin, and A. Zeilinger, “Entanglement by path identity,” *Physical Review Letters* 118, 080401 (2017). <https://doi.org/10.1103/PhysRevLett.118.080401>
- [13] A. Hochrainer, M. Lahiri, and A. Zeilinger, “Quantum indistinguishability by path identity and with undetected photons,” *Reviews of Modern Physics* 94, 025007 (2022). <https://doi.org/10.1103/RevModPhys.94.025007>
- [14] B. Hensen et al., “Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres,” *Nature* 526, 682–686 (2015). <https://doi.org/10.1038/nature15759>
- [15] M. Giustina et al., “Significant-loophole-free test of Bell’s theorem with entangled photons,” *Physical Review Letters* 115, 250401 (2015). <https://doi.org/10.1103/PhysRevLett.115.250401>
- [16] L. K. Shalm et al., “Strong loophole-free test of local realism,” *Physical Review Letters* 115, 250402 (2015). <https://doi.org/10.1103/PhysRevLett.115.250402>
- [17] K. Wang et al., “Violation of Bell inequality with unentangled photons,” *Science Advances* 11(31), eadr1794 (2025). <https://doi.org/10.1126/sciadv.adr1794>
- [18] K. B. Wharton and H. Price, “Bell Inequality Violations Without Entanglement? It’s Just Postselection,” arXiv:2508.13431 (2025). <https://arxiv.org/abs/2508.13431>
- [19] A. Wójcik and D. K. Wójcik, “Simple explanation of apparent Bell nonlocality of unentangled photons,” arXiv:2509.03127 (2025). <https://arxiv.org/abs/2509.03127>
- [20] J. Cieslinski et al., “Unquestionable Bell theorem for interwoven frustrated down conversion processes,” arXiv:2508.19207 (2025). <https://arxiv.org/abs/2508.19207>
- [21] C. Vieira, R. Ramanathan, and A. Cabello, “Test of the physical significance of Bell non-locality,” *Nature Communications* 16, 4390 (2025). <https://www.nature.com/articles/s41467-025-59247-7>