

bem: modeling for neutron Bragg-edge imaging

Jiao Y. Y. Lin*1 and Gian Song1

1 Oak Ridge National Laboratory

DOI: 10.21105/joss.00973

Software

■ Review ♂

Repository ♂Archive ♂

Submitted: 28 August 2018 Published: 18 October 2018

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC-BY).

Summary

Due to its zero net charge, neutron is a unique probe of materials. Low neutron absorption and scattering cross sections by most nuclei make it suitable for studying bulk samples. Unlike X-ray scattering, neutron form factors are not monotonically dependent on atomic numbers; the fact that the neutron scattering cross section of hydrogen is large makes neutron a useful tool in biology. In the past half century, neutron imaging has seen growing applications in various scientific fields including physics, engineering sciences, biology, and archaeology (Strobl et al., 2009).

With energy-resolved neutron imaging techniques, neutron Bragg-edge imaging has recently found applications for materials science in phase mapping, stress/strain mapping, and texture analysis (Josic, Steuwer, & Lehmann, 2010, Sato (2017)). To model Bragg-edge neutron imaging data, it is necessary to calculate the total neutron cross section of a sample. This open-source python package provides easy-to-use functions to calculate coherent elastic (diffraction), incoherent elastic, coherent inelastic, and incoherent inelastic scattering cross sections, as well as absorption cross sections based on approximations and formulas in (Vogel, 2000). Also implemented are algorithms that take into account the March-Dollase texture model, and the Jorgensen peak profile (Vogel, 2000).

Notice of Copyright

This manuscript has been authored by UT-Battelle, LLC under Contract No. DE-AC05-00OR22725 with the U.S. Department of Energy. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes. The Department of Energy will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan (http://energy.gov/downloads/doe-public-access-plan).

Acknowledgements

This work is sponsored by the Laboratory Directed Research and Development Program of Oak Ridge National Laboratory, managed by UT-Battelle LLC, for DOE. Part of this research is supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, User Facilities under contract number DE-AC05-00OR22725.

^{*}Corresponding author: linjiao@ornl.gov



References

Josic, L., Steuwer, A., & Lehmann, E. (2010). Energy selective neutron radiography in material research. Applied Physics A, 99(3), 515–522. doi:10.1007/s00339-010-5602-7

Sato, H. (2017). Deriving quantitative crystallographic information from the wavelength-resolved neutron transmission analysis performed in imaging mode. *Journal of Imaging*, 4(1), 7. doi:10.3390/jimaging4010007

Strobl, M., Manke, I., Kardjilov, N., Hilger, A., Dawson, M., & Banhart, J. (2009). Advances in neutron radiography and tomography. *Journal of Physics D: Applied Physics*, 42(24), 243001. doi:10.1088/0022-3727/42/24/243001

Vogel, S. (2000). A riet veld-approach for the analysis of neutron time-of-flight transmission data (PhD thesis). The University of Kiel. Retrieved from https://core.ac.uk/download/pdf/56723242.pdf