

SeisModels.jl: A Julia package for models of the Earth's interior

Andy Nowacki¹

¹ School of Earth and Environment, University of Leeds

DOI: [10.21105/joss.02043](https://doi.org/10.21105/joss.02043)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Katy Barnhart](#) ↗

Reviewers:

- [@daniellivingston](#)
- [@fhorrobin](#)

Submitted: 13 January 2020

Published: 27 January 2020

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

Summary

The interior structure of the Earth and other quasi-spherical planets reflects their history, development, present-day state and dynamics. Hence inferring their structure is of great interest, and requires the ability to rapidly compute their bulk properties such as total mass, moment of inertia, gravitational acceleration, interior pressure, and so on, in order to compare model predictions with data. `SeisModels.jl` is a Julia package designed to allow such computation.

`SeisModels.jl` can be used to represent arbitrary models of quasi-spherical bodies. In its current release, radially-symmetric bodies can be represented easily using spherical shells whose properties are linearly varying with radius, constant with radius, or parameterised by a set of polynomial coefficients. Density, elastic isotropic velocity (P-wave and S-wave), elastic radially-anisotropic velocity and attenuation are supported as basic model parameters, permitting the computation at arbitrary radius of gravity, pressure, mass, and so on. Inbuilt models include PREM (Dziewoński & Anderson, 1981), AK135 (Kennett, Engdahl, & Buland, 1995) and iasp91 (Kennett & Engdahl, 1991) for the Earth, and the Moon model of Weber, Lin, Garnero, Williams, & Lognonné (2011).

Due to Julia's multiple dispatch programming paradigm, extending `SeisModels.jl` to represent bodies of different types or ones parameterised differently to the inbuilt types is simple and incurs no performance penalty. The software has been benchmarked against independent implementations and computes properties quickly enough to test millions of models against data, enabling robust parameter searches or Monte Carlo sampling. For the example of working with PREM on a desktop computer with an Intel E5-1650 CPU, it takes only 12 ms to calculate the pressure at the centre of the Earth, whilst surface gravity is returned within 5 μ s.

`SeisModels.jl` has been used for research projects and teaching, and provides the ability to read and write files in appropriate formats to calculate normal mode eigenfrequencies and -functions using the [Mineos](#) software, and ray-theoretical travel times using the Java TauP (Crotwell, Owens, & Ritsema, 1999) and Python Obspy (Beyreuther et al., 2010) packages. `SeisModels.jl` is used by the [Mineos.jl](#) Julia module to directly compute normal mode properties for planetary models.

Documentation is available at a [dedicated website](#). The package comes with a test set which can be easily run by users—after adding the package, simply run `import Pkg; Pkg.test("SeisModels")` in Julia.

Acknowledgements

The author was supported by a NERC standard grant (NE/R001154/1). The Deep Earth Research Group at the School of Earth and Environment, University of Leeds is acknowledged

for inspiring the creation of this package, and Andrew Walker for the publication of it.

References

- Beyreuther, M., Barsch, R., Krischer, L., Megies, T., Behr, Y., & Wassermann, J. (2010). ObsPy: A Python toolbox for seismology. *Seismological Research Letters*, 81(3), 530–533. doi:[10.1785/gssrl.81.3.530](https://doi.org/10.1785/gssrl.81.3.530)
- Crotwell, H. P., Owens, T. J., & Ritsema, J. (1999). The taup Toolkit: Flexible seismic travel-time and ray-path utilities. *Seismological Research Letters*, 70(2), 154–160. doi:[10.1785/gssrl.70.2.154](https://doi.org/10.1785/gssrl.70.2.154)
- Dziewoński, A., & Anderson, D. (1981). Preliminary reference Earth model. *Physics of The Earth and Planetary Interiors*, 25(4), 297–356. doi:[10.1016/0031-9201\(81\)90046-7](https://doi.org/10.1016/0031-9201(81)90046-7)
- Kennett, B. L. N., & Engdahl, E. (1991). Traveltimes for global earthquake location and phase identification. *Geophysical Journal International*, 105(2), 429–465. doi:[10.1111/j.1365-246X.1991.tb06724.x](https://doi.org/10.1111/j.1365-246X.1991.tb06724.x)
- Kennett, B. L. N., Engdahl, E., & Buland, R. (1995). Constraints on seismic velocities in the Earth from travel-times. *Geophysical Journal International*, 122(1), 108–124. doi:[10.1111/j.1365-246X.1995.tb03540.x](https://doi.org/10.1111/j.1365-246X.1995.tb03540.x)
- Weber, R. C., Lin, P.-Y., Garnero, E. J., Williams, Q., & Lognonné, P. (2011). Seismic detection of the lunar core. *Science*, 331(6015), 309–312. doi:[10.1126/science.1199375](https://doi.org/10.1126/science.1199375)