

Gromov–Hausdorff distance for directed spaces

Lydia Mezrag

Université de Montréal - MILA Quebec AI institute

joint work with Lisbeth Fajstrup, Brittany Terese Fasy, Wenwen Li, Tatum Rask, Francesca Tombari, and Ziva Urbancic

ComPer 2025: Workshop on Computational Topology

- Introduction
- Preliminaries
- Directed GH-distance: version 1
- Directed GH-distance: version 2
- Directed GH-distance: version 3
- References

- **[Gromov]** - Metric Structures for Riemannian and Non-Riemannian Spaces
- **Kalton & Ostrovskii** - Distances between Banach spaces, ...
- **Length structures:** $len : \mathcal{C}(I) \rightarrow [0, \infty)$ satisfying some axioms (positivity, monotonicity, additivity, invariance under reparametrization and 'continuity').
- **Length spaces.**

(X, d) metric space \longrightarrow Metric Length structure len



new metric d_{len} associated with len

Let (X, d) be a metric space. The **Length** of a path γ is given by

$$l(\gamma) = \sup_{[N]} \sum_{i=1}^N d(\gamma(t_{i-1}), \gamma(t_i)), \quad (1)$$

$[N]$ = partition of I of length N . $l(\gamma) < \infty$ = **rectifiable path**.

Definition 1 (Directed spaces).

A pair $(X, \vec{P}(X))$, is called **d-space** if X is a topological space and $\vec{P}(X)$ is a collection of paths such that:

- ① every constant path belongs to $\vec{P}(X)$,
- ② $\vec{P}(X)$ is closed under non-decreasing reparameterization and subpath.
- ③ if $\gamma, \gamma' \in \vec{P}(X)$ such that $\gamma(1) = \gamma'(0)$, then the concatenation $\gamma\gamma' \in \vec{P}(X)$.

Let \vec{X} and \vec{Y} be d-spaces. A **d-map** $\vec{F}: \vec{X} \rightarrow \vec{Y}$ is a continuous map $F: X \rightarrow Y \mid \forall \gamma \in \vec{P}(X), \text{ the composition } F \circ \gamma \in \vec{P}(Y)$. This defines a category **dTop**. Note that a bijective d-map $\vec{F}: \vec{X} \rightarrow \vec{Y}$ whose inverse is a d-map is called **d-invertible**.

Example 2.

- **Discrete d-space** $(X, \vec{P}(X)) : X \in \mathbf{Top}$ and $\vec{P}(X) = \text{c}^{st}\text{-paths}$.
- **Trivial d-space** $(X, \vec{P}(X)) : X \in \mathbf{Top}$ and $\vec{P}(X) = \mathcal{C}(I, X)$.

Example 3.

Partially ordered set (X, \leq) and $\vec{P}(X) \subseteq \{\gamma \in \mathcal{C} : \gamma(s) \leq \gamma(t) \ \forall s \leq t\}$

A **rectifiable d-space** is a d-space $(X, \vec{P}(X))$, where X is a metric space and every d-path in $\vec{P}(X)$ is **rectifiable**.

- (M, d) metric space, then $\forall A, B \subseteq M$, their **Hausdorff distance** is defined by:

$$d_H(A, B) = \max \left\{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(A, b) \right\} \quad (2)$$

- Given (X, d^X) , (Y, d^Y) , the **Gromov–Hausdorff distance** is defined by:

$$d_{GH}(X, Y) = \inf_{f, g} d_H(f(X), g(Y))$$

where $f: X \hookrightarrow Z$ and $g: Y \hookrightarrow Z$ are isometric embeddings.

Alternate (equivalent definitions)

- $d_{GH}(X, Y) = \inf_{d \in \mathcal{D}(d^X, d^Y)} d_H^{(X \sqcup Y, d)}(X, Y)$
- $d_{GH}(X, Y) = \frac{1}{2} \inf_{\mathcal{R}} \text{dis}(\mathcal{R})$,
- $d_{GH}(X, Y) = \frac{1}{2} \inf_{f, g} \max\{\text{dis}(f), \text{dis}(g), \text{codis}(f, g)\}$

Here $\mathcal{R} \subseteq X \times Y$ is a **correspondence** and

$$\text{dis}(\mathcal{R}) = \sup_{(x, y), (x', y') \in \mathcal{R}} |d^X(x, x') - d^Y(y, y')|.$$

A **zigzag path** btw x and x' is a sequence of d -paths
 $(\gamma_i)_{i=1}^m \mid \gamma_i \in \vec{P}(p_{i-1}, p_i) \cup \vec{P}(p_i, p_{i-1}), x = p_0 \text{ and } x' = p_m.$

Definition 4 (Zigzag distance).

(X, d) metric space and \vec{X} d -space. The **zigzag distance** induced by d is defined by

$$d_{ZZ}(x, x') = \inf_{\gamma \in \vec{P}_{ZZ}(x, x')} l_{ZZ}(\gamma).$$

Proposition 1.

- ① Let \vec{X} be a d -space and d_{ZZ} the zigzag distance induced by d . Then

$$d_{ZZ}(x, x') \geq d(x, x').$$

- ② The zigzag distance, d_{ZZ} , on \vec{X} is an extended metric.
③ Every path γ is C^0 wrt d_{ZZ} .

Given a directed metric space (\vec{X}, d_{ZZ}) , we can construct a new directed metric space $(\vec{X}, (d_{ZZ})_{ZZ})$ where the d-paths are continuous with respect to d_{ZZ} . This construction can be iterated.

Proposition 2 (Idempotence).

Let (\vec{X}, d_{ZZ}) be a directed metric space. Then, $(d_{ZZ})_{ZZ} = d_{ZZ}$.

Change of convention! Topology now is wrt d_{ZZ} .

Definition 5.

Consider two directed metric spaces (\vec{X}, d_{ZZ}^X) and (\vec{Y}, d_{ZZ}^Y) . A d-map $\vec{F}: \vec{X} \rightarrow \vec{Y}$ is called a **d-isometry** if for any x, x' in X ,

$$d_{ZZ}^X(x, x') = d_{ZZ}^Y(\vec{F}(x), \vec{F}(x')).$$

- Two d-spaces \vec{X} and \vec{Y} are **d-isometric** if there is a bijective d-isometry $\vec{F}: \vec{X} \rightarrow \vec{Y}$ such that its inverse ($F^{-1}: Y \rightarrow X$) is a d-map.
- The **directed Hausdorff distance** of \vec{X} and \vec{Y} is defined by

$$\vec{d}_H(\vec{X}, \vec{Y}) = d_H((X, d_{ZZ}), (Y, d_{ZZ})). \quad (3)$$

Definition 6 (GH-distance version 1).

The **directed Gromov–Hausdorff distance** between two directed metric spaces (\vec{X}, d_{ZZ}^X) and (\vec{Y}, d_{ZZ}^Y) is given by:

$$\vec{d}_{GH}(\vec{X}, \vec{Y}) = \inf_{\vec{F}, \vec{G}} \vec{d}_H(f(\vec{X}), g(\vec{Y})), \quad (4)$$

where $\vec{F}: \vec{X} \rightarrow \vec{Z}$ and $\vec{G}: \vec{Y} \rightarrow \vec{Z}$ are directed isometries into some directed metric space (\vec{Z}, d_{ZZ}^Z) .

Theorem 7.

Let (\vec{X}, d_{ZZ}^X) and (\vec{Y}, d_{ZZ}^Y) be directed metric spaces. Then

$$d_{GH}((X, d_{ZZ}^X), (Y, d_{ZZ}^Y)) = \vec{d}_{GH}((\vec{X}, d_{ZZ}^X), (\vec{Y}, d_{ZZ}^Y))$$

Corollary 1.

The directed Gromov–Hausdorff distance is a metric on the space of isometry classes of compact directed metric spaces.

Note Corollary 1 requires compactness of the directed metric space (\vec{X}, d_{ZZ}) .

- For any zigzag connected d-space, $\vec{d}_{GH}(\vec{X}, \vec{X}) = 0$.
- Let (\vec{X}, d_{ZZ}^X) and (\vec{Y}, d_{ZZ}^Y) be a compact directed metric space. Then
 - 1 $\vec{d}_{GH}(\vec{X}, \vec{Y}) \leq \frac{1}{2} \max\{\text{Diam}(\vec{X}), \text{Diam}(\vec{Y})\}$.
 - 2 $\vec{d}_{GH}(\vec{X}, \vec{Y}) = \frac{1}{2} \text{Diam}(\vec{Y})$, if $X = \{x_0\}$.

Conjecture 1.

Let (\vec{X}, d_{ZZ}^X) and (\vec{Y}, d_{ZZ}^Y) be directed compact metric spaces, then

$$d_{GH}(X, Y) \leq \vec{d}_{GH}(\vec{X}, \vec{Y}).$$

If $\vec{F}: (\vec{X}, d_{ZZ}^X) \rightarrow (\vec{Y}, d_{ZZ}^Y)$ is a d-map, we define its distortion by

$$\text{dis}(\vec{F}) = \sup_{x, x' \in \vec{X}} |d_{ZZ}^X(x, x') - d_{ZZ}^Y(\vec{F}(x), \vec{F}(x'))|. \quad (5)$$

Codistorsion is defined analogously.

Definition 8 (GH-distance version 2).

The **distortion distance** between directed metric spaces (\vec{X}, d_{ZZ}^X) and (\vec{Y}, d_{ZZ}^Y) is defined as

$$\vec{d}_{\text{dis}}(\vec{X}, \vec{Y}) = \frac{1}{2} \inf_{\vec{F}, \vec{G}} \max\{\text{dis}(\vec{F}), \text{dis}(\vec{G}), \text{codis}(\vec{F}, \vec{G})\}, \quad (6)$$

- **Triangle Inequality:** \vec{X} , \vec{Y} , and \vec{Z} directed metric spaces. Then,

$$\vec{d}_{\text{dis}}(\vec{X}, \vec{Z}) \leq \vec{d}_{\text{dis}}(\vec{X}, \vec{Y}) + \vec{d}_{\text{dis}}(\vec{Y}, \vec{Z}).$$

- (\vec{X}, d_{ZZ}^X) and (\vec{Y}, d_{ZZ}^Y) compact directed metric spaces. Then,

$$\vec{d}_{\text{dis}}(\vec{X}, \vec{Y}) \leq \frac{1}{2} \max\{\text{diam}(\vec{X}), \text{diam}(\vec{Y})\},$$

where $\text{diam}(\vec{X}) = \text{diam}(X, d_{ZZ}^X)$ and $\text{diam}(\vec{Y}) = \text{diam}(Y, d_{ZZ}^Y)$

Theorem 9.

D-maps $\vec{F} : \vec{X} \rightarrow \vec{Y}$ and $\vec{G} : \vec{Y} \rightarrow \vec{X}$ for which $\text{dis}(\vec{F}) = \text{dis}(\vec{G}) = \text{codis}(\vec{F}, \vec{G}) = 0$ exist if and only if spaces \vec{X} and \vec{Y} are d-isometric.

Proposition 3.

For any bounded directed metric spaces \vec{X} and \vec{Y} ,

$$\vec{d}_{GH}(\vec{X}, \vec{Y}) \leq \vec{d}_{\text{dis}}(\vec{X}, \vec{Y}).$$

- \vec{d}_{dis} is an **extended pseudo-metric** on the space of d-isometry classes of directed metric spaces.

A **d-relation** \mathcal{R} between \vec{X} and \vec{Y} is a relation between X and Y |
 $\forall (x, y) \text{ and } (x', y') \in \mathcal{R}$

- $\forall \gamma_1 \in \vec{P}(x, x') \exists \gamma_2 \in \vec{P}(y, y')$,
- similarly, $\forall \gamma_2 \in \vec{P}(y, y') \exists \gamma_1 \in \vec{P}(x, x')$.

d-correspondence = d-relation + correspondence.

Definition 10 (GH-distance version 3).

The **d-correspondence distortion distance** between directed metric spaces (\vec{X}, d_{ZZ}^X) and (\vec{Y}, d_{ZZ}^Y) is defined as

$$\vec{d}_{\text{c-dis}}(\vec{X}, \vec{Y}) = \frac{1}{2} \inf_{\mathcal{R}} \text{dis}(\mathcal{R}), \quad (7)$$

where \mathcal{R} varies over all the d-correspondences between \vec{X} and \vec{Y} .

Proposition 4.

For every two compact directed metric spaces \vec{X} and \vec{Y} ,

$$\vec{d}_{GH}(\vec{X}, \vec{Y}) \leq \vec{d}_{\text{c-dis}}(\vec{X}, \vec{Y}).$$

References



Fajstrup, Lisbeth and Goubault, Eric and Haucourt, Emmanuel and Mimram, Samuel and Raussen, Martin

Directed Algebraic Topology and Concurrency

Springer, 2016.



Gaucher, Philippe

Homotopy invariants of higher dimensional categories and concurrency in computer science

Mathematical Structures in Computer Science, Cambridge University Press, 2000.



Gaucher, Philippe

Homotopy invariants of higher dimensional categories and concurrency in computer science

Mathematical Structures in Computer Science, Cambridge University Press, 2000.



Kalton, Nigel J and Ostrovskii, Mikhail I

Distances between Banach spaces

Walter de Gruyter, 1999.



Lim, Sunhyuk and Mémoli, Facundo and Smith, Zane
The Gromov-Hausdorff Distance Between Spheres
MSP (Mathematical Sciences Publishers), 2023



Gromov, Mikhael and Katz, Misha and Pansu, Pierre and Semmes, Stephen
Metric structures for Riemannian and non-Riemannian spaces
Springer, 2023.



Mennucci, Andrea CG
On asymmetric distances
Analysis and Geometry in Metric Spaces, Versita, 2013.



Fajstrup, Lisbeth and Costa, João Pita
On the Hierarchy of d-structures
Order, Springer, 2017.



Burago, Dmitri and Burago, Yuri and Ivanov, Sergei
A Course in Metric Geometry
American Mathematical Society, 2001.



Chowdhury, Samir and Mémoli, Facundo
Persistent path homology of directed networks
Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms, 2018.