

Computing Stabilized Decompositions

Jan Jendrysiak
with Håvard Bjerkevik and Fabian Lenzen

TU Graz

October 11, 2025

Structure

1 Decompositions


2 Stabilization of Decompositions

3 Computing the Pruning





Persistence under noise

Goal



Estimate the homology groups of a noisy sample.



Alpha Complex



Persistent Homology under noise




Persistent Homology under noise




Persistent Homology under noise



Persistence barcode



Bifiltrations




Multiparameter Persistence Modules

$$\begin{array}{ccccccc}
 & \mathbb{K} & \longrightarrow & 0 & \longrightarrow & 0 & \\
 & \uparrow & & \uparrow & & \uparrow & \\
 H_1(-; \mathbb{K}) \Rightarrow & \left[\begin{matrix} 1 & 0 \end{matrix} \right] & \longrightarrow & \left[\begin{matrix} 0 & 1 \end{matrix} \right] & \longrightarrow & & \\
 & \uparrow & & \uparrow & & \uparrow & \\
 & \mathbb{K}^2 & \longrightarrow & \mathbb{K} & \longrightarrow & \mathbb{K} & \\
 & \uparrow & & \uparrow & & \uparrow & \\
 & \left[\begin{matrix} 1 \\ 0 \end{matrix} \right] & \longrightarrow & & & & \\
 & \uparrow & & & & & \\
 & \mathbb{K} & \longrightarrow & 0 & \longrightarrow & 0 &
 \end{array}$$

Definition

A *Multiparameter Persistence Module* M is a functor $M: \mathbb{R}^d \rightarrow \text{Vect}_K$.

Hilbert Functions



Definition

Define the *thickness* $\theta(M)$ as $\sup_{\alpha \in \mathbb{R}^2} \dim_{\mathbb{K}} M_\alpha$

Decompositions

Let M be a multiparameter persistence module.

Decompositions

Let M be a multiparameter persistence module.

Goal

Find $\{M_i\}_{i \in I}$ such that $M \xrightarrow{\sim} \bigoplus_{i \in I} M_i$.

with M_i indecomposable.

Decompositions

$$\begin{array}{ccccccc} \mathbb{F} & \longrightarrow & 0 & \longrightarrow & 0 \\ \left[\begin{matrix} 1 & 0 \end{matrix} \right] \uparrow & & \uparrow & & \uparrow \\ \mathbb{F}_2^2 & \xrightarrow{\left[\begin{matrix} 0 & 1 \end{matrix} \right]} & \mathbb{F}_2 & \longrightarrow & \mathbb{F}_2 \\ \left[\begin{matrix} 1 \\ 0 \end{matrix} \right] \uparrow & & \uparrow & & \uparrow \\ \mathbb{F}_2 & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

Decompositions

decomposes as

$$\begin{array}{ccccccc} \mathbb{F}_2 & \longrightarrow & 0 & \longrightarrow & 0 & & \\ \uparrow & & \uparrow & & \uparrow & & \\ \mathbb{F}_2 & \longrightarrow & 0 & \longrightarrow & 0 & \oplus & \mathbb{F}_2 \\ \uparrow & & \uparrow & & \uparrow & & \\ \mathbb{F}_2 & \longrightarrow & 0 & \longrightarrow & 0 & & \\ \uparrow & & \uparrow & & \uparrow & & \\ 0 & \longrightarrow & 0 & \longrightarrow & 0 & & \end{array}$$

0 → 0 → 0

Motivation

Decompositions of Persistence Modules are useful for:

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions
 - Landscapes

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions
 - Landscapes
 - Skyscraper Invariant

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions
 - Landscapes
 - Skyscraper Invariant
 - ...

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions
 - Landscapes
 - Skyscraper Invariant
 - ...
- ② Additive algebraic constructions:

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions
 - Landscapes
 - Skyscraper Invariant
 - ...
- ② Additive algebraic constructions:
 - Homomorphism groups

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions
 - Landscapes
 - Skyscraper Invariant
 - ...
- ② Additive algebraic constructions:
 - Homomorphism groups
 - Submodules

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions
 - Landscapes
 - Skyscraper Invariant
 - ...
- ② Additive algebraic constructions:
 - Homomorphism groups
 - Submodules
 - Resolutions

Motivation

Decompositions of Persistence Modules are useful for:

- ① Computation of additive invariants and distances.
 - Rank Invariant \Rightarrow Signed Barcodes
 - Fibred Barcode \Rightarrow Matching Distance
 - Generalized Rank Invariant
 - Interval-Resolutions
 - Landscapes
 - Skyscraper Invariant
 - ...
- ② Additive algebraic constructions:
 - Homomorphism groups
 - Submodules
 - Resolutions
- ③ As an invariant itself, if the components are small.

Current Situation

What we can do

We can compute decompositions reasonably fast with AIDA.
Between $\sim n^{1.5}$ and $\sim n^3$ for density-scale bifiltrations.

Current Situation

What we can do

We can compute decompositions reasonably fast with AIDA.
Between $\sim n^{1.5}$ and $\sim n^3$ for density-scale bifiltrations.
For interval-decomposable modules in $\mathcal{O}(n^3)$.

Current Situation

What we can do

We can compute decompositions reasonably fast with AIDA.
Between $\sim n^{1.5}$ and $\sim n^3$ for density-scale bifiltrations.
For interval-decomposable modules in $\mathcal{O}(n^3)$.

Obstructions

- Complicated topologies create large indecomposable summands.
- Every Persistence Module is close to being indecomposable.

Instability of Decompositions


Theorem (Bauer, Scoccola '22)

For any $\epsilon > 0$ and finitely presented two-parameter module M , there is an indecomposable module M' that is ϵ -interleaved with M .

Instability of Decompositions

Theorem (Bauer, Scoccola '22)

For any $\epsilon > 0$ and finitely presented two-parameter module M , there is an indecomposable module M' that is ϵ -interleaved with M .



Experimental Evidence




Figure: 246 Generators, 357 Relations,

Experimental Evidence







Figure: 246 Generators, 357 Relations, decomposes into 192 components.







Hilbert Function




Hilbert Function







Hilbert Function





Hilbert Function



3D Sampled Points with Density









4892

generators, 3390 relations, 2742 components





This module is *not* close to being highly decomposable.

Structure

1 Decompositions

2 Stabilization of Decompositions

3 Computing the Pruning

The Pruning Construction

To combat the problem of “almost-indecomposability”, (Bjerkevik ’24) developed the Pruning construction.

The Pruning Construction

To combat the problem of “almost-indecomposability”, (Bjerkevik ’24) developed the Pruning construction.

Theorem

Let $\varepsilon \geq 0$, and let M and N be ε -interleaved pfd modules with $\theta(M) < \infty$. Then $\text{Pru}_\varepsilon(M)$ is a $2\varepsilon\theta(M)$ -refinement of both M and N . In particular $d_I(M, \text{Pru}_\varepsilon(M)) \leq 2\varepsilon\theta(M)$.

Algebraic Preliminaries

Definition

Let M be a persistence module and $\alpha \in \mathbb{R}^2$. We define its *shift* $M[\alpha]$ via

$$M[\alpha]_\beta = M_{\alpha+\beta} \quad M[\alpha]_{\beta \rightarrow \gamma} = M_{(\alpha+\beta) \rightarrow (\alpha+\gamma)}.$$

The structure maps induce natural homomorphisms

$$\text{sh}_M^\alpha: M \rightarrow M[\alpha].$$

Algebraic Preliminaries


Definition

Let M be a persistence module and $\alpha \in \mathbb{R}^2$. We define its *shift* $M[\alpha]$ via

$$M[\alpha]_\beta = M_{\alpha+\beta} \quad M[\alpha]_{\beta \rightarrow \gamma} = M_{(\alpha+\beta) \rightarrow (\alpha+\gamma)}.$$

The structure maps induce natural homomorphisms

$$\text{sh}_M^\alpha: M \rightarrow M[\alpha].$$



Algebraic Preliminaries

Definition

We define functors

$$\text{Ker}_\alpha(M) := \ker \text{sh}_M^\alpha$$

$$\text{Im}_\alpha(M) := \text{im } \text{sh}_{M[-\alpha]}^\alpha$$


Algebraic Preliminaries

Definition

We define functors

$$\text{Ker}_\alpha(M) := \ker \text{sh}_M^\alpha$$

$$\text{Im}_\alpha(M) := \text{im } \text{sh}_{M[-\alpha]}^\alpha$$



Erosion

Definition

Let $\varepsilon \in \mathbb{R}$. Define the *Erosion* of M as


$$\text{Er}_\varepsilon(M) := \text{Im}_{\bar{\varepsilon}}(M) \Big/ (\text{Ker}_{\bar{\varepsilon}}(M) \cap \text{Im}_{\bar{\varepsilon}}(M)) .$$

Erosion

Definition

Let $\varepsilon \in \mathbb{R}$. Define the *Erosion* of M as

$$\text{Er}_\varepsilon(M) := \text{Im}_{\bar{\varepsilon}}(M) \Big/ (\text{Ker}_{\bar{\varepsilon}}(M) \cap \text{Im}_{\bar{\varepsilon}}(M)) .$$





Erosion

Definition

Let $\varepsilon \in \mathbb{R}$. Define the *Erosion* of M as

$$\text{Er}_\varepsilon(M) := \text{Im}_{\bar{\varepsilon}}(M) \Big/ (\text{Ker}_{\bar{\varepsilon}}(M) \cap \text{Im}_{\bar{\varepsilon}}(M)) .$$



The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.

Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.

Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

Define K to be the smallest submodule of I such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \subseteq K$.

The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.

Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

Define K to be the smallest submodule of I such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \subseteq K$.

$$(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \supset (\text{sh}_I^{2\varepsilon})^{-1}(f(0))$$

The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.

Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

Define K to be the smallest submodule of I such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \subseteq K$.

$$(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \supset (\text{sh}_I^{2\varepsilon})^{-1}(f(0)) = \text{Ker}_{2\varepsilon}(M)$$

The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.

Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

Define K to be the smallest submodule of I such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \subseteq K$.

$$(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \supset (\text{sh}_I^{2\varepsilon})^{-1}(f(0)) = \text{Ker}_{2\varepsilon}(M)$$

$$\text{Pru}_\varepsilon(M) := I/K.$$

The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.

Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

Define K to be the smallest submodule of I such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \subseteq K$.

$$(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \supset (\text{sh}_I^{2\varepsilon})^{-1}(f(0)) = \text{Ker}_{2\varepsilon}(M)$$

$$\text{Pru}_\varepsilon(M) := I/K.$$

”Pruning is Eroding, stabilised over $\text{Hom}(M, M[2\varepsilon])$.”

The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.


Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

Define K to be the smallest submodule of I such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \subseteq K$.

$$(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \supset (\text{sh}_I^{2\varepsilon})^{-1}(f(0)) = \text{Ker}_{2\varepsilon}(M)$$

$$\text{Pru}_\varepsilon(M) := I/K.$$

”Pruning is Eroding, stabilised over $\text{Hom}(M, M[2\varepsilon])$.”



The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.

Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

Define K to be the smallest submodule of I such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \subseteq K$.

$$(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \supset (\text{sh}_I^{2\varepsilon})^{-1}(f(0))$$

The Pruning Construction

Definition

Let $M \in \text{A-GrMod}$ and $\varepsilon \in \mathbb{R}_+$.


Define I to be the largest submodule of M such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $f(I) \subseteq \text{sh}_M^{2\varepsilon}(I)$.

Define K to be the smallest submodule of I such that for any $f: M \rightarrow M[2\varepsilon]$ it holds that $(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \subseteq K$.

$$(\text{sh}_I^{2\varepsilon})^{-1}(f(K)) \supset (\text{sh}_I^{2\varepsilon})^{-1}(f(0)) = \text{Ker}_{2\varepsilon}(M)$$

$$\text{Pru}_\varepsilon(M) := I/K.$$

”Pruning is Eroding, stabilised over $\text{Hom}(M, M[2\varepsilon])$.”



Structure

1 Decompositions

2 Stabilization of Decompositions

3 Computing the Pruning

Computing the Pruning

Lemma

Set $I_0 := X$, $K_0 := 0$. We define iteratively

$$I_i := \bigcap_{f \in \text{End}(X)_{2\epsilon}} f^{-1} \circ \text{sh}_X^{2\epsilon}(I_{i-1}) \text{ and} \quad (1)$$

$$K_i := \sum_{f \in \text{End}(X)_{2\epsilon}} (\text{sh}_I^{2\epsilon})^{-1} \circ f(K_{i-1}). \quad (2)$$

Then $I = \bigcap^{\infty} I_i$ and $K = \bigcup^{\infty} K_i$.

Computing the Pruning

Lemma

Set $I_0 := X$, $K_0 := 0$. We define iteratively

$$I_i := \bigcap_{f \in \text{End}(X)_{2\epsilon}} f^{-1} \circ \text{sh}_X^{2\epsilon}(I_{i-1}) \text{ and} \quad (1)$$

$$K_i := \sum_{f \in \text{End}(X)_{2\epsilon}} (\text{sh}_I^{2\epsilon})^{-1} \circ f(K_{i-1}). \quad (2)$$

Then $I = \bigcap^{\infty} I_i$ and $K = \bigcup^{\infty} K_i$.

⇒ This implies an algorithm:

Choose a basis $\{f_i\}$ for $\text{Hom}(M, M[2\epsilon])$ and compute the above.

Computing the Pruning

Lemma

Set $I_0 := X$, $K_0 := 0$. We define iteratively

$$I_i := \bigcap_{f \in \text{End}(X)_{2\epsilon}} f^{-1} \circ \text{sh}_X^{2\epsilon}(I_{i-1}) \text{ and} \quad (1)$$

$$K_i := \sum_{f \in \text{End}(X)_{2\epsilon}} (\text{sh}_I^{2\epsilon})^{-1} \circ f(K_{i-1}). \quad (2)$$

Then $I = \bigcap^{\infty} I_i$ and $K = \bigcup^{\infty} K_i$.

⇒ This implies an algorithm:

Choose a basis $\{f_i\}$ for $\text{Hom}(M, M[2\epsilon])$ and compute the above.

$\dim_{\mathbb{K}} \text{Hom}(M, M[2\epsilon]) \in \mathcal{O}(n^2)$.

Size of $\text{Hom}(M, M[2\varepsilon])$

	Random Points - 78 g			Random Points - 254 g		
ε	0.00	0.01	0.02	0.00	0.01	0.02
$\text{End}(M)_\varepsilon$	713	914	1111	5010	7632	10383

	Two circles - 357 g			Torus - 647 g		
ε	0.00	0.01	0.02	0.00	0.01	0.02
$\text{End}(M)_\varepsilon$	16192	17089	17844	7230	9307	11635

Observation

Let $f = \text{sh}_M^{2\varepsilon} \circ g$ where $g: M \rightarrow M$,
then f can be skipped in the construction of $\text{Pru}_\varepsilon(M)$.

Observation

Let $f = \text{sh}_M^{2\varepsilon} \circ g$ where $g: M \rightarrow M$,
then f can be skipped in the construction of $\text{Pru}_\varepsilon(M)$.

Proposition


Can instead consider a basis for

$$\text{Hom}(M, M[2\varepsilon]) \Big/ (\text{sh}_M^{2\varepsilon})_* \text{Hom}(M, M)$$


Size of $\text{Hom}(M, M[2\varepsilon])$

	Two circles			Torus		
	0.005	0.01	0.02	0.005	0.01	0.02
$\text{End}(M)_0$	16192	16192	16192	7230	7230	7230
$\text{End}(M)_{2\varepsilon}$	16645	17089	17844	8241	9307	11635
$\text{End}(M)_{2\varepsilon/0}$	453	897	1652	1011	2077	4405



Results



Results




Results




192 components

Results




192 components




72 components

Results




Results




Results

Hilbert Function





Hilbert Function




Results



and its largest component




Results




Results



Results




1 component



69 components

Results



Open Problems

- The number of iterations is bounded by $\theta(M)$ - can we do better?


Open Problems

- The number of iterations is bounded by $\theta(M)$ - can we do better?
- Notice that

$$\text{Hom}(M, M[2\varepsilon]) / (\text{sh}_M^{2\varepsilon})_* \text{Hom}(M, M) = \left(\text{End}(M) \otimes F[2\varepsilon]/F \right)_0.$$

Can you compute this faster than $\text{Hom}(M, M[2\varepsilon])$?

Thank you!



Code

<https://github.com/JanJend/Stable-Decomposition>