

# Coexistence of sexually antagonistic alleles

Alba Cervantes-Loreto<sup>1</sup>, Michelle L. Marraffini<sup>1</sup>, Daniel B. Stouffer<sup>1</sup>, and

Sarah P. Flanagan<sup>1</sup>

<sup>1</sup>Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury,

Christchurch 8140, New Zealand

---

**Words in abstract** 203

**Words in manuscript** 5130

**Number of references** 50

**Number of figures** 4

**Number of tables** 2

**Number of text boxes** 0

**Corresponding author** Alba Cervantes-Loreto

**Phone** +64 369 2880

**Email** alba.cervantesloreto@pg.canterbury.ac.nz

---

# 1 Abstract

Sexually antagonistic selection (SAS) occurs when the selection in the traits or loci differs between the sexes. This sexual conflict offers the opportunity for maintaining polymorphism in a population, but it often results in the eventual fixation of the fitter allele. However, the effects of SAS have generally been studied under strong simplifying assumptions, such as constant populations and homogeneous environments, which could considerably change the expected outcomes of SAS. Thus, in this study, we examined how fluctuations in selection and population sizes contributed to the coexistence of sexually antagonistic alleles by adopting an ecological framework that allowed us to examine evolutionary dynamics through the same lens as the coexistence of competing species. We performed simulations of alleles invading a population while allowing selection and populations sizes to fluctuate over time. Then, we quantified coexistence outcomes and the relative contribution of each type of fluctuation to each alleles' invasion growth rate. Our results showed that environmental fluctuations can dramatically increase the expected genetic variation under SAS. The positive contribution of fluctuations, however, depended on the sex and allele where invasion occurred. This study contributes to the growing body of work that shows the importance of non-constant environments on the maintenance of genetic diversity.

## 2 Introduction

The question of how genetic variation is maintained despite the effects of selection and drift is central within evolutionary biology (Walsh & Lynch, 2018). Classical explanations include overdominance (heterozygote advantage) or frequency-dependent selection (Hedrick, 2007), but in the modern era of genomic data, all patterns of variation that exceed the expected variation under neutrality tend to be categorized broadly as balancing selection, regardless of the evolutionary mechanism (Mitchell-Olds *et al.*, 2007). In species with separate sexes, balancing selection can arise due to sexually antagonistic selection (Connallon & Clark, 2014), which occurs when the direction of natural selection on traits or loci differs between the sexes (Lande, 1980; Arnqvist & Rowe, 2013).

Sexually antagonistic selection can maintain polymorphisms of otherwise disadvantageous alleles in a population (Gavrilets, 2014), which in turn can result in phenotypically distinct sexes that express different morphological, physiological, and behavioral traits (Mori *et al.*, 2017; Connallon & Hall, 2018). Nonetheless, the extent to which sexually antagonistic selection can maintain polymorphism in a population is thought to be limited (Connallon & Clark, 2012). This is because theoretical studies have found that the necessary parameter conditions that give rise to balancing selection are often highly restrictive (Kidwell *et al.*, 1977; Pamilo, 1979; Hedrick, 1999; Curtsinger *et al.*, 1994; Patten *et al.*, 2010; Jordan & Charlesworth, 2012). Importantly, the effect of sexually antagonistic selection generally has been studied under strong simplifying assumptions such as constant population sizes and homogeneous environments (Kidwell *et al.*, 1977; Pamilo, 1979; Immler

*et al.*, 2012; Jordan & Charlesworth, 2012). Studies that have explored the effect of sexually antagonistic selection with more realistic assumptions, such as temporal fluctuations in selection (Connallon *et al.*, 2018) or demographic fluctuations (Connallon & Clark, 2012) have found that polymorphism can be maintained in a much wider set of conditions than classical studies predict. These results suggest that environmental fluctuations are essential to fully understand the effects of sexually antagonistic selection.

The contribution of environmental fluctuations to genetic diversity remains a debated issue in evolutionary biology. Classic theoretical models predict that temporal fluctuations in environmental conditions are unlikely to maintain a genetic polymorphism in haploid populations (Dempster, 1955; Hedrick, 1974; 1986). However, other studies have found that fluctuating selection can maintain genetic variance when populations experience density dependence (Dean, 2005), on sex-linked traits (Reinhold, 2000), or in populations where generations overlap (Ellner & Hairston Jr, 1994; Ellner & Sasaki, 1996). Similarly, temporal changes in population sizes have been shown to mitigate the effect of genetic drift in small populations (Pemberton *et al.*, 1996) and in annual plant systems (Nunney, 2002). Importantly, progress requires more than just identifying if environmental fluctuations can maintain genetic diversity in a population, but to quantify how exactly they contribute to its maintenance (Ellner *et al.*, 2016).

Temporal variability in the environment has been shown to promote diversity maintenance in ecological contexts (Levins, 1979; Armstrong & McGehee, 1980; Chesson, 2000a; Barabás *et al.*, 2018). Note that from an ecological perspective, polymorphism of sexually antagonistic alleles is equivalent to the coexistence of species, and the fixation of either

one of the alleles in a population is equivalent to competitive exclusion. Allelic polymorphism, thus, can be examined through the same lens as the coexistence of competing species. (Ellner & Hairston Jr, 1994; Ellner & Sasaki, 1996; Dean, 2005; Schreiber, 2010). The benefit of analyzing evolutionary dynamics through this lens is that the main theoretical framework used to examine how competing species coexist, often called Modern Coexistence Theory (Chesson, 2000b; 1994; Barabás *et al.*, 2018), allows the quantification of how environmental fluctuations contribute to coexistence. Despite that the use of Modern Coexistence Theory often requires complex mathematical analysis of the models describing the systems dynamics and restrictive assumptions to make them tractable (Barabás *et al.*, 2018), recent computation approaches allow the quantification of the relative importance of environmental fluctuations to coexistence using simulations (Ellner *et al.*, 2016; 2019; Shoemaker *et al.*, 2020).

Here, we seek to explicitly quantify how temporal environmental fluctuations contribute to the maintenance of polymorphism under sexually antagonistic selection by using simulations. We examined how fluctuations in selection values, fluctuations in population sizes, and their interactions can further or hinder polymorphism. In particular, we examined i) Can fluctuations in population sizes and selection values allow sexually antagonistic alleles to coexist when differences in their fitness would typically not allow them to? and ii) What are the relative contributions of different types of fluctuations that allow two sexually antagonistic alleles to be maintained in a population? Our study provides the tools to analyze sexual antagonism from a novel perspective and contributes to answering long-lasting questions regarding the effect of non-constant environments on

90 genetic diversity.

### 91 **3 Methods**

92 We first present a model that describes the evolutionary dynamics of sexually antago-  
93 nistic alleles. We then show how we simulated different scenarios of alleles invading a  
94 population, where we allowed population sizes, selection, both, or neither to vary. Fi-  
95 nally, we detail how we examined the relative contribution of each type of fluctuation to  
96 the maintenance of polymorphism.

#### 97 **Population dynamics of sexually antagonistic alleles**

98 Our model examined evolution at a single, biallelic locus. We examined the dynammics  
99 of two sexually antagonistic alleles,  $j$  and  $k$ , that affect fitness in the haploid state. The  
100 frequencies of each allele in each sex at the beginning of a life-cycle at time  $t$  are given by:

$$p_{jm,t} = \frac{n_{jm,t}}{N_{m,t}} \quad (1)$$

$$p_{jf,t} = \frac{n_{jf,t}}{N_{f,t}} \quad (2)$$

$$p_{km,t} = \frac{N_{m,t} - n_{jm,t}}{N_{m,t}} \quad (3)$$

$$p_{kf,t} = \frac{N_{f,t} - n_{jf,t}}{N_{f,t}} \quad (4)$$

104 where  $N_{m,t}$  and  $N_{f,t}$  are the total numbers of males and females in the population at time  
105  $t$ ,  $n_{jf,t}$  is the number of females  $f$  with allele  $j$ , and  $n_{jm,t}$  is the number of males  $m$  with

106 allele  $j$  at time  $t$ , respectively.

107 The individuals in the population mate at random before selection occurs, and there-  
108 fore the frequency of offspring with allele  $j$  after mating,  $p'_{j,t}$  can be expressed as:

$$p'_{j,t} = \frac{n_{jf}}{N_f} \frac{n_{jm}}{N_m} + \frac{1}{2} \frac{n_{jf}}{N_f} \frac{(N_m - n_{jm})}{N_m} + \frac{1}{2} \frac{(N_f - n_{jf})}{N_f} \frac{n_{jm}}{N_m}, \quad (5)$$

109 which upon rearranging and simplifying gives:

$$p'_{j,t} = \frac{N_{m,t}n_{jf,t} + N_{f,t}n_{jm,t}}{2N_fN_m}. \quad (6)$$

110 Selection acts upon these offspring in order to determine the allelic frequencies in females  
111 and males in the next generation,  $t + 1$ . As an example, the frequency of females with  
112 allele  $j$  after selection is given by:

$$p_{jf,t+1} = \frac{n_{jf,t+1}}{N_{f,t+1}} = \frac{p'_{j,t}w_{jf}}{p'_{t,j}w_{jf} + (1 - p'_{t,j})w_{kf}} \quad (7)$$

113 The logarithmic per capita growth rate of allele  $j$  in females is therefore given by the  
114 number of females carrying allele  $j$  after selection divided by the original number of fe-  
115 males carrying allele  $j$ :

$$r_{jf,t} = \ln \left( \frac{n_{jf,t+1}}{n_{jf,t}} \right) \quad (8)$$

116 An equivalent expression for the logarithmic per capita growth rate of allele  $j$  in males  
117  $m$  can be obtained by exchanging  $f$  for  $m$  across the various subscripts in Eqn. 7.

Polymorphism in a sexual population, however, is ultimately influenced by growth and establishment of an allele across both sexes. Therefore, the growth rate of allele  $j$  across the entire population of females *and* males is given by:

$$r_{j,t} = \ln \left( \frac{n_{jf,t+1} + n_{jm,t+1}}{n_{jf,t} + n_{jm,t}} \right) \quad (9)$$

An equivalent expression describes  $r_{k,t}$ , the growth rate of allele  $k$ .

Our model further assumed allele  $j$  always has a high fitness in females ( $w_{jf} = 1$ ) but variable fitness in males ( $w_{jm} < 1$ ); and allele  $k$  has a high fitness in males ( $w_{km} = 1$ ) but variable fitness in females ( $w_{kf} < 1$ ). The selection against allele  $j$  in males is therefore  $S_m = 1 - w_{jm}$ , and the selection against allele  $k$  in females is  $S_f = 1 - w_{kf}$ . When population sizes and selection values are constant, selection maintains both alleles in the population, under the condition that:

$$\frac{S_m}{1 + S_m} < S_f < \frac{S_m}{1 - S_m} \quad (10)$$

(Kidwell *et al.*, 1977; Pamilo, 1979; Patten *et al.*, 2010; Connallon *et al.*, 2018). Thus, the maintenance of polymorphism of sexually antagonistic alleles is solely determined by the values of  $S_m$  and  $S_f$ . Note that in our model, the values  $S_m$  and  $S_f$  are bounded from 0 to 1. Therefore the parameter space of sexually antagonistic selection is within the range  $0 < S_m, S_f < 1$ . Classic theoretical models predict that, in constant environments, polymorphism is maintained in  $\approx 0.38$  of the parameter space (Kidwell *et al.*, 1977; Pamilo, 1979; Connallon *et al.*, 2018). Nonetheless it is unrealistic to assume population sizes and



135 selection are constant through time. Temporal changes in population densities are ubiqui-  
136 tous in nature (Connallon & Clark, 2012; Reinhold, 2000). Similarly, the effect of sexual  
137 selection has been shown to vary through space and time (Kasumovic *et al.*, 2008). If fluctu-  
138 ations in population sizes or selection values have an effect on the coexistence of sexually  
139 antagonistic alleles, it would be reflected in increases or decreases of the proportion of the  
140 parameter space of selection where polymorphism is maintained.

## References

- Armstrong, R.A. & McGehee, R. (1980). Competitive exclusion. *The American Naturalist*, 115, 151–170.
- Arnqvist, G. & Rowe, L. (2013). *Sexual conflict*. Princeton University Press.
- Barabás, G., D’Andrea, R. & Stump, S.M. (2018). Chesson’s coexistence theory. *Ecological Monographs*, 88, 277–303.
- Chesson, P. (1994). Multispecies competition in variable environments. *Theoretical population biology*, 45, 227–276.
- Chesson, P. (2000a). General theory of competitive coexistence in spatially-varying environments. *Theoretical Population Biology*, 58, 211–237.
- Chesson, P. (2000b). Mechanisms of maintenance of species diversity. *Annual review of Ecology and Systematics*, 31, 343–366.
- Connallon, T. & Clark, A.G. (2012). A general population genetic framework for antagonistic selection that accounts for demography and recurrent mutation. *Genetics*, 190, 1477–1489.
- Connallon, T. & Clark, A.G. (2014). Balancing selection in species with separate sexes: insights from fisher’s geometric model. *Genetics*, 197, 991–1006.
- Connallon, T. & Hall, M.D. (2018). Environmental changes and sexually antagonistic selection. *eLS*, pp. 1–7.

- 160 Connallon, T., Sharma, S. & Olito, C. (2018). Evolutionary Consequences of Sex-Specific  
161 Selection in Variable Environments: Four Simple Models Reveal Diverse Evolutionary  
162 Outcomes. *The American Naturalist*, 193, 93–105.
- 163 Curtsinger, J.W., Service, P.M. & Prout, T. (1994). Antagonistic pleiotropy, reversal of  
164 dominance, and genetic polymorphism. *The American Naturalist*, 144, 210–228.
- 165 Dean, A.M. (2005). Protecting haploid polymorphisms in temporally variable environ-  
166 ments. *Genetics*, 169, 1147–1156.
- 167 Dempster, E.R. (1955). Maintenance of genetic heterogeneity. In: *Cold Spring Harbor Sym-*  
168 *posia on Quantitative Biology*. Cold Spring Harbor Laboratory Press, vol. 20, pp. 25–32.
- 169 Ellner, S. & Hairston Jr, N.G. (1994). Role of overlapping generations in maintaining  
170 genetic variation in a fluctuating environment. *The American Naturalist*, 143, 403–417.
- 171 Ellner, S. & Sasaki, A. (1996). Patterns of genetic polymorphism maintained by fluctuating  
172 selection with overlapping generations. *theoretical population biology*, 50, 31–65.
- 173 Ellner, S.P., Snyder, R.E. & Adler, P.B. (2016). How to quantify the temporal storage effect  
174 using simulations instead of math. *Ecology letters*, 19, 1333–1342.
- 175 Ellner, S.P., Snyder, R.E., Adler, P.B. & Hooker, G. (2019). An expanded modern coexis-  
176 tence theory for empirical applications. *Ecology Letters*, 22, 3–18.
- 177 Gavrillets, S. (2014). Is sexual conflict an “engine of speciation”? *Cold Spring Harbor*  
178 *perspectives in biology*, 6, a017723.

179 Hedrick, P.W. (1974). Genetic variation in a heterogeneous environment. i. temporal het-  
 180 erogeneity and the absolute dominance model. *Genetics*, 78, 757–770.

181 Hedrick, P.W. (1986). Genetic polymorphism in heterogeneous environments: a decade  
 182 later. *Annual review of ecology and systematics*, 17, 535–566.

183 Hedrick, P.W. (1999). Antagonistic pleiotropy and genetic polymorphism: a perspective.  
 184 *Heredity*, 82, 126–133.

185 Hedrick, P.W. (2007). Balancing selection. *Current Biology*, 17, R230–R231.

186 Immler, S., Arnqvist, G. & Otto, S.P. (2012). Ploidally antagonistic selection maintains  
 187 stable genetic polymorphism. *Evolution: International Journal of Organic Evolution*, 66,  
 188 55–65.

189 Jordan, C.Y. & Charlesworth, D. (2012). The potential for sexually antagonistic polymor-  
 190 phism in different genome regions. *Evolution: International Journal of Organic Evolution*,  
 191 66, 505–516.

192 Kasumovic, M.M., Bruce, M.J., Andrade, M.C. & Herberstein, M.E. (2008). Spatial and  
 193 temporal demographic variation drives within-season fluctuations in sexual selection.  
 194 *Evolution: International Journal of Organic Evolution*, 62, 2316–2325.

195 Kidwell, J., Clegg, M., Stewart, F. & Prout, T. (1977). Regions of stable equilibria for  
 196 models of differential selection in the two sexes under random mating. *Genetics*, 85,  
 197 171–183.

- 198 Lande, R. (1980). Sexual dimorphism, sexual selection, and adaptation in polygenic char-  
199 acters. *Evolution*, pp. 292–305.
- 200 Levins, R. (1979). Coexistence in a variable environment. *The American Naturalist*, 114,  
201 765–783.
- 202 Mitchell-Olds, T., Willis, J.H. & Goldstein, D.B. (2007). Which evolutionary processes  
203 influence natural genetic variation for phenotypic traits? *Nature Reviews Genetics*, 8,  
204 845–856.
- 205 Mori, E., Mazza, G. & Lovari, S. (2017). Sexual dimorphism. *Encyclopedia of Animal Cogni-  
206 tion and Behavior* (J. Vonk, and T. Shakelford, Eds). Springer International Publishing, Switzer-  
207 land, pp. 1–7.
- 208 Nunney, L. (2002). The effective size of annual plant populations: the interaction of a seed  
209 bank with fluctuating population size in maintaining genetic variation. *The American  
210 Naturalist*, 160, 195–204.
- 211 Pamilo, P. (1979). Genic variation at sex-linked loci: Quantification of regular selection  
212 models. *Hereditas*, 91, 129–133.
- 213 Patten, M.M., Haig, D. & Ubeda, F. (2010). Fitness variation due to sexual antagonism  
214 and linkage disequilibrium. *Evolution: International Journal of Organic Evolution*, 64,  
215 3638–3642.
- 216 Pemberton, J., Smith, J., Coulson, T.N., Marshall, T.C., Slate, J., Paterson, S., Albon, S.,  
217 Clutton-Brock, T.H. & Sneath, P.H.A. (1996). The maintenance of genetic polymorphism

- 218 in small island populations: large mammals in the hebrides. *Philosophical Transactions*  
219 *of the Royal Society of London. Series B: Biological Sciences*, 351, 745–752.
- 220 Reinhold, K. (2000). Maintenance of a genetic polymorphism by fluctuating selection on  
221 sex-limited traits. *Journal of Evolutionary Biology*, 13, 1009–1014.
- 222 Schreiber, S.J. (2010). Interactive effects of temporal correlations, spatial heterogeneity  
223 and dispersal on population persistence. *Proceedings of the Royal Society B: Biological*  
224 *Sciences*, 277, 1907–1914.
- 225 Shoemaker, L.G., Barner, A.K., Bittleston, L.S. & Teufel, A.I. (2020). Quantifying the rela-  
226 tive importance of variation in predation and the environment for species coexistence.  
227 *Ecology letters*, 23, 939–950.
- 228 Walsh, B. & Lynch, M. (2018). *Evolution and Selection of Quantitative Traits*. OUP Oxford.