

# General Introduction

Alba Cervantes-Loreto

## Models of biotic interactions

Interactions between organisms underpin the persistence of almost all life forms on Earth (Lawton, 1999). Furthermore, a large body of work has shown that biotic interactions determine emergent properties of natural systems, such as stability (May, 1972; Wootton & Stouffer, 2016; Song & Saavedra, 2018), resilience (Capdevila *et al.*, 2021), ecosystem functioning (Turnbull *et al.*, 2013; Godoy *et al.*, 2020), and the coexistence of multiple species (Chesson, 2000; Saavedra *et al.*, 2017). Unsurprisingly, numerous ecological and evolutionary concepts revolve around the effects that organisms exert on each other (Gause, 1934; MacArthur & Levins, 1967; Thompson, 1999; HilleRisLambers *et al.*, 2012; Chase & Leibold, 2009).

Describing the effect of biotic interactions often requires the use of mathematical models to represent them (Maynard-Smith, 1978; Rossberg *et al.*, 2019). Mathematical descriptions of interactions are “useful fictions” (Box *et al.*, 2011) in a twofold manner. First, they create a description of how organisms that coincide in space and time affect each other. Almost all known types of interactions can be described in the form of mathematical ex-

pressions that reproduce the observed data faithfully (Volterra, 1926; Holling, 1959; Holt, 1977; Adler *et al.*, 2018; Wood & Thomas, 1999; Holland *et al.*, 2002; Vázquez *et al.*, 2005; Stouffer & Novak, 2021) . For example, the effect neighboring plants have on each other can be accurately described in various natural systems with individual fitness models that include solely the densities of the interacting species embedded in a function of negative density dependence (Adler *et al.*, 2018; Hart *et al.*, 2018). Second, models are practical tools with which to make predictions beyond the phenomena they describe and thus, provide general insights into how natural systems operate (Evans *et al.*, 2012; Stouffer, 2019). For instance, models that describe competitive interactions between plants have been extensively used to demonstrate the mechanisms that maintain diversity when species compete for the same pool of resources (Levine & HilleRisLambers, 2009; Godoy *et al.*, 2014; Godoy & Levine, 2014; Stouffer *et al.*, 2018; Bimler *et al.*, 2018)

## **Are simple models always better?**

Models that capture the effect of biotic interactions are abstractions of reality, and abstractions always reflect choices (Levins, 2006). Building models that include all aspects of reality is not only impractical but also unfeasible. Therefore, ecologists and evolutionary biologists have to continuously make choices regarding which variables to include in a model and which to omit (Evans *et al.*, 2012; Rossberg *et al.*, 2019). A common assumption when building models is that to achieve general insights, we should favor simple models (Evans *et al.*, 2013). Indeed there is a general belief in ecology and evolution that a good model should include as little as possible (Evans *et al.*, 2013; Orzack, 2012). This belief

is often rooted in an implicit philosophical stance that one can not maximize generality (i.e., models that apply to more than one system) and realism (i.e., models that produce accurate predictions for a system) (Levins, 2006; Evans *et al.*, 2012).

Inevitably, model building in biology leads to a key question that will, in turn, modify the outcomes achieved by any model: when is a model “realistic” enough (Stouffer, 2019)? The answer to this question will depend on the purpose for which a model is built. Models that capture the effect of biotic interactions tend to fall into the category of “demonstration models”. These types of models are often based on phenomenological descriptions of processes and have the general aim to show that the modeled principles are sufficient to produce the phenomena of interest (Evans *et al.*, 2013). Demonstration models however, do not help decide whether the modelled principles are *necessary* (Evans *et al.*, 2013). The task to decide the necessary principles and thus the answer to the question of when a model is realistic enough becomes the modeler’s responsibility. In many cases, the answer to this question can appear arbitrary or solely determined by the predominant paradigm regarding the studied system (Holland *et al.*, 2006; Bascompte *et al.*, 2006; Kokko & López-Sepulcre, 2007; AlAdwani & Saavedra, 2019; Mayfield & Stouffer, 2017; Martyn *et al.*, 2021).

Always favoring simple models in ecological and evolutionary studies can be problematic from two perspectives. First, the assumption that more complex models do not lead to general insights is seldomly tested. For example, most models that capture competitive interactions between plants have the implicit assumption that competitive effects between individuals are always additive and direct (Schoener, 1974; Freckleton & Watkin-

son, 2001; Kraft *et al.*, 2015). However, when models and data collection were set up to capture non-additive effects of interactions between individuals of co-occurring species, the evidence overwhelmingly showed that including these levels of biotic complexity was necessary to capture plant interactions accurately (Mayfield & Stouffer, 2017; Martyn *et al.*, 2021; Lai *et al.*, 2021). Thus, in some cases, increasing complexity increases rather than hampers the general insights obtained from models of biotic interactions.

Second, failing to include necessary levels of complexity can hinder our ability to predict how natural communities will react to novel conditions. Predictions of how natural systems will behave in the future are inherently challenging (Sutherland, 2006). Nevertheless, ignoring heterogeneities at various levels can further complicate rather than simplify predictions (Evans *et al.*, 2012). For instance, demographic models tend to treat ecological and evolutionary dynamics separately, despite the general understanding that both processes are often intertwined (MacArthur, 1962; Kokko & López-Sepulcre, 2007). Ignoring eco-evolutionary feedbacks leads to predictions that are inconsistent with empirical data and produce counterintuitive results in novel conditions (Kokko & López-Sepulcre, 2007). Thus, the implicit assumption that good models should include as little as possible should be treated with caution in ecological and evolutionary contexts (Evans *et al.*, 2013; Kokko & López-Sepulcre, 2007; Abrams, 2001).

## Thesis aims

Despite arguments in favor of increasing realism in models of biotic interactions, doing so remains a challenge in many ecological and evolutionary studies. Part of these diffi-

culties arises from the lack of appropriate theoretical frameworks that allow the inclusion of different levels of complexity (Mayfield & Stouffer, 2017; Martyn *et al.*, 2021). Thus, it remains unclear whether increasing model realism is warranted for many natural systems. In this thesis, I propose theoretical, statistical, and empirical frameworks to increase realism in models of biotic interactions to understand when higher levels of complexity are justified. Furthermore, I also explore the consequences of increasing model realism in predictions related to diversity maintenance at ecological and evolutionary scales. The individual chapters of this thesis are thematically broad, but all address in a different way the challenges and consequences of incorporating biotic, abiotic, and mathematical complexity in models of biotic interactions.

## **Including biotic, abiotic and mathematical complexity**

Perhaps one of the most challenging aspects of building more realistic models of biotic interactions is the lack of theoretical frameworks that allow incorporating complexities observed in empirical contexts (Abrams, 1983). Such is the case of competition between pollinators that forage for the same resources (Thomson & Page, 2020). Despite an overwhelming amount of empirical evidence showing that pollinators modify their foraging behavior in the presence of other foraging species (Morse, 1977; Inouye, 1978; Thompson *et al.*, 2006; Brosi & Briggs, 2013; Briggs, 2016), incorporating these behavioral changes into population dynamics remains elusive (Thomson & Page, 2020).

## References

- Abrams, P.A. (1983). Arguments in favor of higher order interactions. *The American Naturalist*, 121, 887–891.
- Abrams, P.A. (2001). Describing and quantifying interspecific interactions: a commentary on recent approaches. *Oikos*, 94, 209–218.
- Adler, P.B., Smull, D., Beard, K.H., Choi, R.T., Furniss, T., Kulmatiski, A., Meiners, J.M., Tredennick, A.T. & Veblen, K.E. (2018). Competition and coexistence in plant communities: intraspecific competition is stronger than interspecific competition. *Ecology letters*, 21, 1319–1329.
- AlAdwani, M. & Saavedra, S. (2019). Is the addition of higher-order interactions in ecological models increasing the understanding of ecological dynamics? *Mathematical Biosciences*, 315, 108222.
- Bascompte, J., Jordano, P. & Olesen, J.M. (2006). Response to comment on “asymmetric coevolutionary networks facilitate biodiversity maintenance”. *Science*, 313, 1887–1887.
- Bimler, M.D., Stouffer, D.B., Lai, H.R. & Mayfield, M.M. (2018). Accurate predictions of coexistence in natural systems require the inclusion of facilitative interactions and environmental dependency. *Journal of Ecology*, 106, 1839–1852.
- Box, G.E., Luceño, A. & del Carmen Paniagua-Quinones, M. (2011). *Statistical control by monitoring and adjustment*. vol. 700. John Wiley & Sons.

- Briggs, H.M. (2016). *Competitive context drives pollinator behavior: linking foraging plasticity, natural pollen deposition, and plant reproduction*. Ph.D. thesis, UC Santa Cruz.
- Brosi, B.J. & Briggs, H.M. (2013). Single pollinator species losses reduce floral fidelity and plant reproductive function. *Proceedings of the National Academy of Sciences*, 110, 13044–13048.
- Capdevila, P., Stott, I., Oliveras Menor, I., Stouffer, D.B., Raimundo, R.L., White, H., Barbour, M. & Salguero-Gómez, R. (2021). Reconciling resilience across ecological systems, species and subdisciplines.
- Chase, J.M. & Leibold, M.A. (2009). *Ecological niches*. University of Chicago Press.
- Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of Ecology and Systematics*, 31, 343–366.
- Evans, M.R., Grimm, V., Johst, K., Knuuttila, T., De Langhe, R., Lessells, C.M., Merz, M., O'Malley, M.A., Orzack, S.H., Weisberg, M. *et al.* (2013). Do simple models lead to generality in ecology? *Trends in ecology & evolution*, 28, 578–583.
- Evans, M.R., Norris, K.J. & Benton, T.G. (2012). Predictive ecology: systems approaches.
- Freckleton, R. & Watkinson, A. (2001). Predicting competition coefficients for plant mixtures: reciprocity, transitivity and correlations with life-history traits. *Ecology Letters*, 4, 348–357.
- Gause, G.F. (1934). Experimental analysis of vito volterra's mathematical theory of the struggle for existence. 79, 16–17.

- Godoy, O., Gómez-Aparicio, L., Matías, L., Pérez-Ramos, I.M. & Allan, E. (2020). An excess of niche differences maximizes ecosystem functioning. *Nature communications*, 11, 1–10.
- Godoy, O., Kraft, N.J.B. & Levine, J.M. (2014). Phylogenetic relatedness and the determinants of competitive outcomes. *Ecology Letters*, 17, 836–844.
- Godoy, O. & Levine, J.M. (2014). Phenology effects on invasion success: insights from coupling field experiments to coexistence theory. *Ecology*, 95, 726–736.
- Hart, S.P., Freckleton, R.P. & Levine, J.M. (2018). How to quantify competitive ability. *Journal of Ecology*, 106, 1902–1909.
- HilleRisLambers, J., Adler, P.B., Harpole, W.S., Levine, J.M. & Mayfield, M.M. (2012). Rethinking community assembly through the lens of coexistence theory. *Annual review of ecology, evolution, and systematics*, 43, 227–248.
- Holland, J.N., DeAngelis, D.L. & Bronstein, J.L. (2002). Population dynamics and mutualism: functional responses of benefits and costs. *The American Naturalist*, 159, 231–244.
- Holland, J.N., Okuyama, T. & DeAngelis, D.L. (2006). Comment on “asymmetric coevolutionary networks facilitate biodiversity maintenance”. *Science*, 313, 1887–1887.
- Holling, C.S. (1959). Some characteristics of simple types of predation and parasitism. *The Canadian Entomologist*, 91, 385–398.
- Holt, R.D. (1977). Predation, apparent competition, and the structure of prey communities. *Theoretical population biology*, 12, 197–229.



- 160 Inouye, D.W. (1978). Resource Partitioning in Bumblebees: Experimental Studies of For-  
161 aging Behavior. *Ecology*, 59, 672–678.
- 162 Kokko, H. & López-Sepulcre, A. (2007). The ecogenetic link between demography and  
163 evolution: can we bridge the gap between theory and data? *Ecology Letters*, 10, 773–782.
- 164 Kraft, N.J., Godoy, O. & Levine, J.M. (2015). Plant functional traits and the multidimen-  
165 sional nature of species coexistence. *Proceedings of the National Academy of Sciences*, 112,  
166 797–802.
- 167 Lai, H.R., Chong, K.Y., Yee, A.T.K., Mayfield, M.M. & Stouffer, D.B. (2021). Non-additive  
168 biotic interactions improve predictions of tropical tree growth and impact community  
169 size structure. *bioRxiv*, pp. 2020–09.
- 170 Lawton, J.H. (1999). Are there general laws in ecology? *Oikos*, pp. 177–192.
- 171 Levine, J.M. & HilleRisLambers, J. (2009). The importance of niches for the maintenance  
172 of species diversity. *Nature*, 461, 254–257.
- 173 Levins, R. (2006). Strategies of abstraction. *Biology and Philosophy*, 21, 741–755.
- 174 MacArthur, R. & Levins, R. (1967). The limiting similarity, convergence, and divergence  
175 of coexisting species. *The american naturalist*, 101, 377–385.
- 176 MacArthur, R.H. (1962). Some generalized theorems of natural selection. *Proceedings of*  
177 *the National Academy of Sciences of the United States of America*, 48, 1893.

- Martyn, T.E., Stouffer, D.B., Godoy, O., Bartomeus, I., Pastore, A.I. & Mayfield, M.M. (2021). Identifying “useful” fitness models: Balancing the benefits of added complexity with realistic data requirements in models of individual plant fitness. *The American Naturalist*, 197, 415–433.
- May, R.M. (1972). Will a large complex system be stable? *Nature*, 238, 413–414.
- Mayfield, M.M. & Stouffer, D.B. (2017). Higher-order interactions capture unexplained complexity in diverse communities. *Nature ecology & evolution*, 1, 1–7.
- Maynard-Smith, J. (1978). *Models in ecology*. CUP Archive.
- Morse, D.H. (1977). Resource Partitioning in Bumble Bees: The Role of Behavioral Factors. *Science*, 197, 678–680.
- Orzack, S.H. (2012). The philosophy of modelling or does the philosophy of biology have any use? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 170–180.
- Rossberg, A.G., Barabás, G., Possingham, H.P., Pascual, M., Marquet, P.A., Hui, C., Evans, M.R. & Meszéna, G. (2019). Let’s train more theoretical ecologists—here is why. *Trends in ecology & evolution*, 34, 759–762.
- Saavedra, S., Rohr, R.P., Bascompte, J., Godoy, O., Kraft, N.J. & Levine, J.M. (2017). A structural approach for understanding multispecies coexistence. *Ecological Monographs*, 87, 470–486.
- Schoener, T.W. (1974). Some methods for calculating competition coefficients from resource-utilization spectra. *The American Naturalist*, 108, 332–340.

- 198 Song, C. & Saavedra, S. (2018). Will a small randomly assembled community be feasible  
199 and stable? *Ecology*, 99, 743–751.
- 200 Stouffer, D.B. (2019). All ecological models are wrong, but some are useful. *Journal of*  
201 *Animal Ecology*, 88, 192–195.
- 202 Stouffer, D.B. & Novak, M. (2021). Hidden layers of density dependence in consumer  
203 feeding rates. *Ecology Letters*, 24, 520–532.
- 204 Stouffer, D.B., Wainwright, C.E., Flanagan, T. & Mayfield, M.M. (2018). Cyclic population  
205 dynamics and density-dependent intransitivity as pathways to coexistence between co-  
206 occurring annual plants. *Journal of Ecology*, 106, 838–851.
- 207 Sutherland, W.J. (2006). Predicting the ecological consequences of environmental change:  
208 a review of the methods. *Journal of Applied Ecology*, 43, 599–616.
- 209 Thompson, A.R., Nisbet, R.M. & Schmitt, R.J. (2006). Dynamics of mutualist populations  
210 that are demographically open. *Journal of Animal Ecology*, 75, 1239–1251.
- 211 Thompson, J.N. (1999). The evolution of species interactions. *Science*, 284, 2116–2118.
- 212 Thomson, D.M. & Page, M.L. (2020). The importance of competition between insect pol-  
213 linators in the Anthropocene. *Current Opinion in Insect Science*, 38, 55–62.
- 214 Turnbull, L.A., Levine, J.M., Loreau, M. & Hector, A. (2013). Coexistence, niches and  
215 biodiversity effects on ecosystem functioning. *Ecology letters*, 16, 116–127.

- 216 Vázquez, D.P., Morris, W.F. & Jordano, P. (2005). Interaction frequency as a surrogate for  
217 the total effect of animal mutualists on plants. *Ecology letters*, 8, 1088–1094.
- 218 Volterra, V. (1926). Fluctuations in the abundance of a species considered mathematically.  
219 *Nature*, 118, 558–560.
- 220 Wood, S.N. & Thomas, M.B. (1999). Super-sensitivity to structure in biological models.  
221 *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 266, 565–570.
- 222 Wootton, K. & Stouffer, D. (2016). Many weak interactions and few strong; food-web  
223 feasibility depends on the combination of the strength of species' interactions and their  
224 correct arrangement. *Theoretical Ecology*, 9, 185–195.