

Increasing realism in models of biotic interactions: ecological and evolutionary consequences

Alba Cervantes-Loreto

Abstract

Interactions between organisms give rise to emergent properties of natural systems. This underpins the ubiquity of biotic interactions in the study of ecological and evolutionary dynamics. The representation of biotic interactions often requires models and simplifying assumptions since it is impossible to account for all aspects of the world in a single model. Critical choices, such as the number of species that can alter the interaction between a focal pair or which abiotic variables constitute the environment, are necessary when building ecological and evolutionary models. Such simplifying assumptions inevitably lead to the omission of heterogeneities at various levels. Complexity that is unaccounted for can, in turn, make the relationships between organisms appear noisy and fundamentally change model-based predictions. Despite this, ecological and evolutionary studies often lack appropriate frameworks that allow the inclusion of different levels of complexity in representations of biotic interactions. Thus, it is unclear whether including more realistic assumptions is warranted for the vast majority of natural systems. In this thesis, I explore

how incorporating complexity as abiotic and biotic modifiers, as well as different sources of uncertainty, reveals potential explanatory generalities in natural systems. I also explore how accounting for these variables changes predictions related to the maintenance of diversity at ecological and evolutionary scales. Throughout this thesis, I focus on different types of interactions and organisms and propose mathematical and statistical frameworks that can be used beyond the studied systems.

In **Chapter 2**, I explore how the presence of multiple species and different environmental contexts change the strength of plant-pollinator interactions. I propose a framework for using pollinator functional responses to examine the role of pollinator-pollinator interactions and abiotic conditions in altering the time between floral visits of a focal pollinator. I show that both density dependent responses and abiotic conditions are necessary to explain the number of visits a pollinator makes. In **Chapter 3**, I explore how incorporating different sources of uncertainty changes predictions of species coexistence. I do this by simultaneously exploring how different model formulations, environmental contexts, and parameter uncertainty change the probability of predicting coexistence in an experimental system. I provide direct evidence that predictions of species coexistence are likely to change given the models used to quantify density-dependence. I also provide a theoretical framework to explore predictions made with different models. Finally, in **Chapter 4**, I adopt an ecological framework to examine the evolutionary dynamics of sexually antagonistic alleles. I show that incorporating environmental fluctuations can substantially increase the amount of genetic diversity in a population under sexually antagonistic selection. Overall, the results of my thesis show that the assumptions adopted

40 by some ecological and evolutionary models tend to be oversimplifying. Here, I provide
41 tools for ecologists and evolutionary biologists to explore more realistic representations
42 of biotic interactions as well as their consequences for diversity maintenance.

Preface

My thesis has been prepared as a collection of three standalone scientific articles. Each chapter is a standalone piece of research and, therefore, I only provide a general Introduction and Conclusion chapters linking the three chapters together. In **Chapter 1**, I focus on describing how my three chapters are connected. In **Chapter 5**, I focus on summarising the results from each of my thesis chapters and their combined implications both in both how we study interactions and their consequences for diversity maintenance. Finally, I further expand on new ideas beyond those presented in the different chapters to discuss about the future steps moving forward.

At the time of thesis submission, each of these three articles are in different stages of the publication process.

Chapter 2: “The context dependency of pollinator interference: how environmental conditions and co-foraging species impact floral visitation” was published in May 2021 in the journal *Ecology Letters* in volume 24, no. 7, pages 1443–1454.

Chapter 3: “The interplay of environmental conditions, parameter sensitivity and structural sensitivity in predictions of species coexistence” is in preparation for submission to *Ecology Letters*.

Chapter 4: “Quantifying the relative contributions of environmental fluctuations to the maintenance of a sexually antagonistic polymorphism” is in preparation for submission to *The American Naturalist*.

Chapter 1: General Introduction

Models of biotic interactions

Interactions between organisms underpin the persistence of almost all life forms on Earth (Lawton, 1999). Furthermore, a large body of work has shown that biotic interactions determine emergent properties of natural systems, such as stability (May, 1972; Wootton & Stouffer, 2016; Song & Saavedra, 2018), resilience (Capdevila *et al.*, 2021), ecosystem functioning (Turnbull *et al.*, 2013; Godoy *et al.*, 2020), and the coexistence of multiple species (Chesson, 2000; Saavedra *et al.*, 2017). Unsurprisingly, numerous ecological and evolutionary concepts revolve around the effects that organisms exert on each other (Gause, 1934; MacArthur & Levins, 1967; Thompson, 1999; HilleRisLambers *et al.*, 2012; Chase & Leibold, 2009).

From their origin as natural sciences, the disciplines of ecology and evolution have shifted from a descriptive towards a more predictive and quantitative approach. This shift brought with it the use of mathematical models to describe natural phenomena (Maynard-Smith, 1978; Rossberg *et al.*, 2019). Mathematical descriptions of interactions are “useful fictions” (Box *et al.*, 2011) in a twofold manner. First, they create a description of how organisms that coincide in space and time affect each other. Almost all known types of interactions can be described in the form of mathematical expressions that reproduce the observed data faithfully (Volterra, 1926; Holling, 1959; Holt, 1977; Adler *et al.*, 2018; Wood & Thomas, 1999; Holland *et al.*, 2002; Vázquez *et al.*, 2005; Stouffer & Novak, 2021) Second, models are practical tools with which to make predictions beyond the

84 phenomena they describe and thus, provide general insights into how natural systems
85 operate (Evans *et al.*, 2012; Stouffer, 2019; Rossberg *et al.*, 2019).

86 **The perils of simple models**

87 Models that capture the effect of biotic interactions are abstractions of reality, and abstrac-
88 tions always reflect choices (Levins, 2006). Building models that include all aspects of
89 reality is not only impractical but also unfeasible. Therefore, ecologists and evolutionary
90 biologists have to continuously make choices regarding which variables to include in a
91 model and which to omit (Evans *et al.*, 2012; Rossberg *et al.*, 2019). A common assumption
92 when building models is that to achieve general insights, we should favor simple models
93 (Evans *et al.*, 2013). Indeed there is a general belief in ecology and evolution that a good
94 model should include as little as possible (Evans *et al.*, 2013; Orzack, 2012). This belief
95 is often rooted in an implicit philosophical stance that one can not maximize generality
96 (i.e., models that apply to more than one system) and realism (i.e., models that produce
97 accurate predictions for a system) (Levins, 2006; Evans *et al.*, 2012).

98 Inevitably, model building in biology leads to a key question that will, in turn, mod-
99 ify the outcomes achieved by any model: when is a model “realistic” enough (Stouffer,
100 2019)? The answer to this question will depend on the purpose for which a model is
101 built. Models that capture the effect of biotic interactions tend to fall into the category of
102 “demonstration models”. These types of models are often based on phenomenological
103 descriptions of processes and have the general aim to show that the modeled principles
104 are sufficient to produce the phenomena of interest (Evans *et al.*, 2013). Demonstration

models however, do not help decide whether the modelled principles are *necessary* (Evans *et al.*, 2013). The task to decide the necessary principles and thus the answer to the question of when a model is realistic enough becomes the modeler's responsibility. In many cases, the answer to this question can appear arbitrary or solely determined by the predominant paradigm regarding the studied system (Holland *et al.*, 2006; Bascompte *et al.*, 2006; Kokko & López-Sepulcre, 2007; AlAdwani & Saavedra, 2019; Mayfield & Stouffer, 2017; Martyn *et al.*, 2021).

Always favoring simple models in ecological and evolutionary studies can be problematic from two perspectives. First, the assumption that more complex models do not lead to general insights is seldomly tested. For example, most models that capture competitive interactions between plants have the implicit assumption that competitive effects between individuals are always additive and direct (Schoener, 1974; Freckleton & Watkinson, 2001; Kraft *et al.*, 2015). However, when models and data collection were set up to capture non-additive effects of interactions between individuals of co-occurring species, the evidence overwhelmingly showed that including these levels of biotic complexity was necessary to capture plant interactions accurately (Mayfield & Stouffer, 2017; Martyn *et al.*, 2021; Lai *et al.*, 2021). Thus, in some cases, increasing complexity increases rather than hampers the general insights obtained from models of biotic interactions.

Second, failing to include necessary levels of complexity can hinder our ability to predict how natural communities will react to novel conditions. Predictions of how natural systems will behave in the future are inherently challenging (Sutherland, 2006). Nevertheless, ignoring heterogeneities at various levels can further complicate rather than simplify

predictions (Evans *et al.*, 2012). For instance, demographic models tend to treat ecological and evolutionary dynamics separately, despite the general understanding that both processes are often intertwined (MacArthur, 1962; Kokko & López-Sepulcre, 2007). Ignoring eco-evolutionary feedbacks leads to predictions that are inconsistent with empirical data and produce counterintuitive results in novel conditions (Kokko & López-Sepulcre, 2007). Thus, the implicit assumption that good models should include as little as possible should be treated with caution in ecological and evolutionary contexts (Evans *et al.*, 2013; Kokko & López-Sepulcre, 2007; Abrams, 2001).

Challenges and consequences of increasing realism

Despite arguments in favor of increasing realism in models of biotic interactions, doing so remains a challenge in many ecological and evolutionary studies. One of those challenges arises from the lack of theoretical frameworks that allow incorporating intricate empirical observations into models (Abrams, 1983). Such is the case of competition between pollinators that forage for the same resources (Thomson & Page, 2020). An overwhelmingly amount of empirical evidence shows that pollinators modify their foraging behavior in the presence of other foraging species (Morse, 1977; Inouye, 1978; Thompson *et al.*, 2006; Brosi & Briggs, 2013; Briggs, 2016), however, incorporating these behavioral changes into population dynamics remains elusive (Thomson & Page, 2020). Furthermore, density-dependent responses could themselves depend on the abiotic conditions pollinators experience, as many studies have shown that environmental conditions can drastically change how pollinators behave while foraging (Heinrich, 1976; Thomson *et al.*, 1987; Cnaani *et al.*,

2006; Westphal *et al.*, 2006; Briggs *et al.*, 2018; Classen *et al.*, 2020). A coherent framework with which to incorporate both abiotic and biotic drivers into plant–pollinator interactions was lacking. To this end, in **Chapter 2** I develop a general framework to show how pollinator functional responses can be used to incorporate biotic and abiotic drivers into floral visitation rates. Furthermore, I show the empirical relevance of this framework by parameterizing different models that incorporated pollinator–pollinator interactions and environmental conditions when predicting observed data from a highly controlled foraging chamber experiment. Results from this chapter have substantial implications related to our understanding of how species loss and environmental change might affect mutualistic communities.

Another theoretical challenge emerges when alternative models to represent biotic interactions are used interchangeably. Such is the case of phenomenological models of plant competition, where more than one mathematical form can faithfully reproduce empirical data (Levine & HilleRisLambers, 2009; Godoy & Levine, 2014; Godoy *et al.*, 2014; Mayfield & Stouffer, 2017; Bimler *et al.*, 2018). The effect biotic and abiotic drivers have in model based predictions can be dramatically different due to uncertainty associated with phenomenological models (Jørgensen & Bendoricchio, 2001; Flora *et al.*, 2011; Aldebert Clement & Stouffer Daniel B., 2018). To understand the interplay between uncertainty and abiotic complexity, in **Chapter 3** I introduce a mathematical and statistical framework to simultaneously explore how different phenomenological models of plant competition, environmental contexts, and parameter uncertainty determine predictions of species coexistence. Additionally, I use this framework to make predictions around a pairwise com-

petition experiment between annual plants, where I show that the effect of abiotic conditions in predictions of species coexistence is not independent of the model formulation used to describe species interactions.

Finally, it is not only important to understand whether increasing realism changes predictions, but it is also essential to understand how. For instance, theoretical studies have shown that environmental fluctuations can substantially increase the levels of genetic diversity in populations that experience sexually antagonistic selection (Connallon & Clark, 2012; Connallon *et al.*, 2019; Patten *et al.*, 2010; Jordan & Charlesworth, 2012). However, there are no approaches that directly quantify how environmental fluctuations promote genetic diversity in populations that experience sexual conflict. Hence, in **Chapter 4** I adopt an ecological framework that allows the quantification of the relative contributions of environmental fluctuations using simulations to capture the effect of environmental fluctuations in evolutionary dynamics. In this chapter I show that incorporating environmental heterogeneity is essential to fully understand the effect of when and how sexually antagonistic selection can maintain genetic diversity.

Concluding remarks

In this thesis, I propose theoretical and statistical frameworks that allow increasing realism in models of biotic interactions with the aim to understand when higher levels of complexity are justified. Furthermore, I also explore the consequences of increasing model realism in predictions related to diversity maintenance at ecological and evolutionary scales. The individual chapters of this thesis are thematically broad as they are

191 focused on different types of interactions and organisms, but all address in a different
192 way the challenges and consequences of incorporating biotic and abiotic complexity in
193 the study of biotic interactions.

References

- Abrams, P.A. (1983). Arguments in favor of higher order interactions. *The American Naturalist*, 121, 887–891.
- Abrams, P.A. (2001). Describing and quantifying interspecific interactions: a commentary on recent approaches. *Oikos*, 94, 209–218.
- Adler, P.B., Smull, D., Beard, K.H., Choi, R.T., Furniss, T., Kulmatiski, A., Meiners, J.M., Tredennick, A.T. & Veblen, K.E. (2018). Competition and coexistence in plant communities: intraspecific competition is stronger than interspecific competition. *Ecology letters*, 21, 1319–1329.
- AlAdwani, M. & Saavedra, S. (2019). Is the addition of higher-order interactions in ecological models increasing the understanding of ecological dynamics? *Mathematical Biosciences*, 315, 108222.
- Aldebert Clement & Stouffer Daniel B. (2018). Community dynamics and sensitivity to model structure: towards a probabilistic view of process-based model predictions. *Journal of The Royal Society Interface*, 15, 20180741.
- Bascompte, J., Jordano, P. & Olesen, J.M. (2006). Response to comment on “asymmetric coevolutionary networks facilitate biodiversity maintenance”. *Science*, 313, 1887–1887.
- Bimler, M.D., Stouffer, D.B., Lai, H.R. & Mayfield, M.M. (2018). Accurate predictions of coexistence in natural systems require the inclusion of facilitative interactions and environmental dependency. *Journal of Ecology*, 106, 1839–1852.

- 214 Box, G.E., Luceño, A. & del Carmen Paniagua-Quinones, M. (2011). *Statistical control by*
215 *monitoring and adjustment*. vol. 700. John Wiley & Sons.
- 216 Briggs, H.M. (2016). *Competitive context drives pollinator behavior: linking foraging plasticity,*
217 *natural pollen deposition, and plant reproduction*. Ph.D. thesis, UC Santa Cruz.
- 218 Briggs, H.M., Graham, S., Switzer, C.M. & Hopkins, R. (2018). Variation in context-
219 dependent foraging behavior across pollinators. *Ecology and evolution*, 8, 7964–7973.
- 220 Brosi, B.J. & Briggs, H.M. (2013). Single pollinator species losses reduce floral fidelity
221 and plant reproductive function. *Proceedings of the National Academy of Sciences*, 110,
222 13044–13048.
- 223 Capdevila, P., Stott, I., Oliveras Menor, I., Stouffer, D.B., Raimundo, R.L., White, H., Bar-
224 bour, M. & Salguero-Gómez, R. (2021). Reconciling resilience across ecological systems,
225 species and subdisciplines.
- 226 Chase, J.M. & Leibold, M.A. (2009). *Ecological niches*. University of Chicago Press.
- 227 Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of*
228 *Ecology and Systematics*, 31, 343–366.
- 229 Classen, A., Eardley, C.D., Hemp, A., Peters, M.K., Peters, R.S., Ssymank, A. & Steffan-
230 Dewenter, I. (2020). Specialization of plant–pollinator interactions increases with tem-
231 perature at mt. kilimanjaro. *Ecology and evolution*, 10, 2182–2195.
- 232 Cnaani, J., Thomson, J.D. & Papaj, D.R. (2006). Flower Choice and Learning in Foraging

- 233 Bumblebees: Effects of Variation in Nectar Volume and Concentration. *Ethology*, 112,
234 278–285.
- 235 Connallon, T. & Clark, A.G. (2012). A general population genetic framework for antag-
236 onistic selection that accounts for demography and recurrent mutation. *Genetics*, 190,
237 1477–1489.
- 238 Connallon, T., Sharma, S. & Olito, C. (2019). Evolutionary Consequences of Sex-Specific
239 Selection in Variable Environments: Four Simple Models Reveal Diverse Evolutionary
240 Outcomes. *The American Naturalist*, 193, 93–105.
- 241 Evans, M.R., Grimm, V., Johst, K., Knuuttila, T., De Langhe, R., Lessells, C.M., Merz, M.,
242 O'Malley, M.A., Orzack, S.H., Weisberg, M. *et al.* (2013). Do simple models lead to
243 generality in ecology? *Trends in ecology & evolution*, 28, 578–583.
- 244 Evans, M.R., Norris, K.J. & Benton, T.G. (2012). Predictive ecology: systems approaches.
- 245 Flora, C., David, N., Mathias, G., Andrew, M. & Jean-Christophe, P. (2011). Structural
246 sensitivity of biological models revisited. *Journal of Theoretical Biology*, 283, 82–91.
- 247 Freckleton, R. & Watkinson, A. (2001). Predicting competition coefficients for plant mix-
248 tures: reciprocity, transitivity and correlations with life-history traits. *Ecology Letters*, 4,
249 348–357.
- 250 Gause, G.F. (1934). Experimental analysis of vito volterra's mathematical theory of the
251 struggle for existence. 79, 16–17.

252 Godoy, O., Gómez-Aparicio, L., Matías, L., Pérez-Ramos, I.M. & Allan, E. (2020). An
 253 excess of niche differences maximizes ecosystem functioning. *Nature communications*,
 254 11, 1–10.

255 Godoy, O., Kraft, N.J.B. & Levine, J.M. (2014). Phylogenetic relatedness and the determi-
 256 nants of competitive outcomes. *Ecology Letters*, 17, 836–844.

257 Godoy, O. & Levine, J.M. (2014). Phenology effects on invasion success: insights from
 258 coupling field experiments to coexistence theory. *Ecology*, 95, 726–736.

259 Heinrich, B. (1976). Resource Partitioning Among Some Eusocial Insects: Bumblebees.
 260 *Ecology*, 57, 874–889.

261 HilleRisLambers, J., Adler, P.B., Harpole, W.S., Levine, J.M. & Mayfield, M.M. (2012).
 262 Rethinking community assembly through the lens of coexistence theory. *Annual review*
 263 *of ecology, evolution, and systematics*, 43, 227–248.

264 Holland, J.N., DeAngelis, D.L. & Bronstein, J.L. (2002). Population dynamics and mutu-
 265 alism: functional responses of benefits and costs. *The American Naturalist*, 159, 231–244.

266 Holland, J.N., Okuyama, T. & DeAngelis, D.L. (2006). Comment on “asymmetric coevo-
 267 lutionary networks facilitate biodiversity maintenance”. *Science*, 313, 1887–1887.

268 Holling, C.S. (1959). Some characteristics of simple types of predation and parasitism1.
 269 *The Canadian Entomologist*, 91, 385–398.

270 Holt, R.D. (1977). Predation, apparent competition, and the structure of prey communi-
 271 ties. *Theoretical population biology*, 12, 197–229.

- 272 Inouye, D.W. (1978). Resource Partitioning in Bumblebees: Experimental Studies of For-
273 aging Behavior. *Ecology*, 59, 672–678.
- 274 Jordan, C.Y. & Charlesworth, D. (2012). The potential for sexually antagonistic polymor-
275 phism in different genome regions. *Evolution: International Journal of Organic Evolution*,
276 66, 505–516.
- 277 Jørgensen, S.E. & Bendoricchio, G. (2001). *Fundamentals of Ecological Modelling*. vol. 21.
278 Elsevier.
- 279 Kokko, H. & López-Sepulcre, A. (2007). The ecogenetic link between demography and
280 evolution: can we bridge the gap between theory and data? *Ecology Letters*, 10, 773–782.
- 281 Kraft, N.J., Godoy, O. & Levine, J.M. (2015). Plant functional traits and the multidimen-
282 sional nature of species coexistence. *Proceedings of the National Academy of Sciences*, 112,
283 797–802.
- 284 Lai, H.R., Chong, K.Y., Yee, A.T.K., Mayfield, M.M. & Stouffer, D.B. (2021). Non-additive
285 biotic interactions improve predictions of tropical tree growth and impact community
286 size structure. *bioRxiv*, pp. 2020–09.
- 287 Lawton, J.H. (1999). Are there general laws in ecology? *Oikos*, pp. 177–192.
- 288 Levine, J.M. & HilleRisLambers, J. (2009). The importance of niches for the maintenance
289 of species diversity. *Nature*, 461, 254–257.
- 290 Levins, R. (2006). Strategies of abstraction. *Biology and Philosophy*, 21, 741–755.

- 291 MacArthur, R. & Levins, R. (1967). The limiting similarity, convergence, and divergence
292 of coexisting species. *The american naturalist*, 101, 377–385.
- 293 MacArthur, R.H. (1962). Some generalized theorems of natural selection. *Proceedings of*
294 *the National Academy of Sciences of the United States of America*, 48, 1893.
- 295 Martyn, T.E., Stouffer, D.B., Godoy, O., Bartomeus, I., Pastore, A.I. & Mayfield, M.M.
296 (2021). Identifying “useful” fitness models: Balancing the benefits of added complexity
297 with realistic data requirements in models of individual plant fitness. *The American*
298 *Naturalist*, 197, 415–433.
- 299 May, R.M. (1972). Will a large complex system be stable? *Nature*, 238, 413–414.
- 300 Mayfield, M.M. & Stouffer, D.B. (2017). Higher-order interactions capture unexplained
301 complexity in diverse communities. *Nature ecology & evolution*, 1, 1–7.
- 302 Maynard-Smith, J. (1978). *Models in ecology*. CUP Archive.
- 303 Morse, D.H. (1977). Resource Partitioning in Bumble Bees: The Role of Behavioral Factors.
304 *Science*, 197, 678–680.
- 305 Orzack, S.H. (2012). The philosophy of modelling or does the philosophy of biology have
306 any use? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 170–180.
- 307 Patten, M.M., Haig, D. & Ubeda, F. (2010). Fitness variation due to sexual antagonism
308 and linkage disequilibrium. *Evolution: International Journal of Organic Evolution*, 64,
309 3638–3642.

- Rossberg, A.G., Barabás, G., Possingham, H.P., Pascual, M., Marquet, P.A., Hui, C., Evans, M.R. & Meszéna, G. (2019). Let's train more theoretical ecologists—here is why. *Trends in ecology & evolution*, 34, 759–762.
- Saavedra, S., Rohr, R.P., Bascompte, J., Godoy, O., Kraft, N.J. & Levine, J.M. (2017). A structural approach for understanding multispecies coexistence. *Ecological Monographs*, 87, 470–486.
- Schoener, T.W. (1974). Some methods for calculating competition coefficients from resource-utilization spectra. *The American Naturalist*, 108, 332–340.
- Song, C. & Saavedra, S. (2018). Will a small randomly assembled community be feasible and stable? *Ecology*, 99, 743–751.
- Stouffer, D.B. (2019). All ecological models are wrong, but some are useful. *Journal of Animal Ecology*, 88, 192–195.
- Stouffer, D.B. & Novak, M. (2021). Hidden layers of density dependence in consumer feeding rates. *Ecology Letters*, 24, 520–532.
- Sutherland, W.J. (2006). Predicting the ecological consequences of environmental change: a review of the methods. *Journal of Applied Ecology*, 43, 599–616.
- Thompson, A.R., Nisbet, R.M. & Schmitt, R.J. (2006). Dynamics of mutualist populations that are demographically open. *Journal of Animal Ecology*, 75, 1239–1251.
- Thompson, J.N. (1999). The evolution of species interactions. *Science*, 284, 2116–2118.

- 329 Thomson, D.M. & Page, M.L. (2020). The importance of competition between insect pol-
330 linators in the Anthropocene. *Current Opinion in Insect Science*, 38, 55–62.
- 331 Thomson, J.D., Peterson, S.C. & Harder, L.D. (1987). Response of traplining bumble bees
332 to competition experiments: shifts in feeding location and efficiency. *Oecologia*, 71, 295–
333 300.
- 334 Turnbull, L.A., Levine, J.M., Loreau, M. & Hector, A. (2013). Coexistence, niches and
335 biodiversity effects on ecosystem functioning. *Ecology letters*, 16, 116–127.
- 336 Vázquez, D.P., Morris, W.F. & Jordano, P. (2005). Interaction frequency as a surrogate for
337 the total effect of animal mutualists on plants. *Ecology letters*, 8, 1088–1094.
- 338 Volterra, V. (1926). Fluctuations in the abundance of a species considered mathematically.
339 *Nature*, 118, 558–560.
- 340 Westphal, C., Steffan-Dewenter, I. & Tscharntke, T. (2006). Bumblebees experience land-
341 scapes at different spatial scales: possible implications for coexistence. *Oecologia*, 149,
342 289–300.
- 343 Wood, S.N. & Thomas, M.B. (1999). Super-sensitivity to structure in biological models.
344 *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 266, 565–570.
- 345 Wootton, K. & Stouffer, D. (2016). Many weak interactions and few strong; food-web
346 feasibility depends on the combination of the strength of species' interactions and their
347 correct arrangement. *Theoretical Ecology*, 9, 185–195.