

INCREASING REALISM IN MODELS OF BIOTIC INTERACTIONS:

ecological and evolutionary consequences



ALBA CERVANTES LORETO

*A thesis submitted in fulfillment of the requirements
for the degree of Doctor of Philosophy*

School of Biological Sciences
College of Science
University of Canterbury

Alba Cervantes Loreto. *Increasing realism in models of biotic interactions: ecological and evolutionary consequences* ; Doctoral thesis, November 2021

SUPERVISOR:

Daniel B. Stouffer

CO-SUPERVISORS:

Margaret M. Mayfield

Andrew Letten

Christchurch, New Zealand

Con amor a mis padres, Rosalva Loreto y Francisco Cervantes

ABSTRACT

Interactions between organisms give rise to emergent properties of natural systems. This underpins the ubiquity of biotic interactions in the study of ecological and evolutionary dynamics. The representation of biotic interactions often requires models and simplifying assumptions since it is impossible to account for all aspects of the world in a single model. Critical choices, such as the number of species that can alter the interaction between a focal pair or which abiotic variables constitute the environment, are necessary when building ecological and evolutionary models. Such simplifying assumptions inevitably lead to the omission of heterogeneities at various levels. Complexity that is unaccounted for can, in turn, make the relationships between organisms appear noisy and fundamentally change model-based predictions. Despite this, ecological and evolutionary studies often lack appropriate frameworks that allow the inclusion of different levels of complexity in representations of biotic interactions. Thus, it is unclear whether including more realistic assumptions is warranted for the vast majority of natural systems. In this thesis, I explore how incorporating complexity as abiotic and biotic modifiers, as well as different sources of uncertainty, reveals potential explanatory generalities in natural systems. I also explore how accounting for these variables changes predictions related to the maintenance of diversity at ecological and evolutionary scales. Throughout this thesis, I focus on different types of interactions and organisms and propose mathematical and statistical frameworks that can be used beyond the studied systems.

In [Chapter 2](#), I explore how the presence of multiple species and different environmental contexts change the strength of plant-pollinator interactions. I propose a framework for using pollinator functional responses to examine the role of pollinator-pollinator interactions and abiotic conditions in altering the time between floral visits of a focal pollinator. I show that both density dependent responses and abiotic conditions are necessary to explain the number of visits a pollinator makes. In [Chapter 3](#), I explore how incorporating different sources of uncertainty changes predictions of species coexistence. I do this by simultaneously exploring how different model formulations, environmental contexts, and parameter uncertainty change the probability of predicting coexistence in a pairwise competition experiment of annual plants. I provide direct evidence that predictions of species coexistence are likely to change given the models used to quantify density-dependence. I also provide a theoretical framework to explore predictions made with different models. Finally, in [Chapter 4](#), I adopt an ecological framework to examine the evolutionary dynamics of sexually antagonistic alleles. I show that environmental fluctuations can substantially increase the amount of genetic diversity in a

population under sexually antagonistic selection by contributing positively to allele's invasion growth rates. Overall, the results of my thesis show that the assumptions adopted by some ecological and evolutionary models tend to be oversimplifying. Here, I provide tools for ecologists and evolutionary biologists to explore more realistic representations of biotic interactions as well as their consequences for diversity maintenance.

PREFACE

My thesis has been prepared as a collection of three standalone scientific articles. Each chapter is a standalone piece of research and, therefore, I only provide a general Introduction and Conclusion chapters linking the three chapters together. In [Chapter 1](#), I focus on describing how my three chapters are connected. In [Chapter 5](#), I focus on summarising the results from each of my thesis chapters and their combined implications both in both how we study interactions and their consequences for diversity maintenance. Finally, I further expand on new ideas beyond those presented in the different chapters to discuss about the future steps moving forward.

At the time of thesis submission, each of these three articles are in different stages of the publication process.

[Chapter 2](#): “The context dependency of pollinator interference: how environmental conditions and co-foraging species impact floral visitaion” was published in May 2021 in the journal *Ecology Letters* in volume 24, no. 7, pages 1443–1454.

[Chapter 3](#): “The interplay of environmental conditions, parameter sensitivity and structural sensitivity in predictions of species coexistence” is in preparation for submission to *Ecology Letters*.

[Chapter 4](#): “Quantifying the relative contributions of environmental fluctuations to the maintenance of a sexually antagonistic polymorphism” is in preparation for submission to *The American Naturalist*.

ACKNOWLEDGEMENTS

Here go my aknowledgments

CONTENTS

1	GENERAL INTRODUCTION	1
2	THE CONTEXT-DEPENDENCY OF POLLINATOR INTERFERENCE	15
3	THE INTERPLAY OF ENVIRONMENTAL CONDITIONS, PARAMETER SENSITIVITY AND STRUCTURAL SENSITIVITY IN PREDICTIONS OF SPECIES COEXISTENCE	17
4	COEXISTENCE OF ALLELES	19
5	CONCLUSION	21

LIST OF FIGURES

LIST OF TABLES

GENERAL INTRODUCTION

Interactions between organisms underpin the persistence of almost all life forms on Earth (Lawton, 1999). Furthermore, a large body of work has shown that biotic interactions determine emergent properties of natural systems, such as stability (May, 1972; Song and Saavedra, 2018; Wootton and Stouffer, 2016), resilience (Capdevila et al., 2021), ecosystem functioning (Godoy et al., 2020; Turnbull et al., 2013), and the coexistence of multiple species (Chesson, 2000; Saavedra et al., 2017). Unsurprisingly, numerous ecological and evolutionary concepts revolve around the effects that organisms exert on each other (Chase and Leibold, 2009; Gause, 1934; HilleRisLambers et al., 2012; MacArthur and Levins, 1967; Thompson, 1999, 2014).

From their origins as natural sciences, the disciplines of ecology and evolution have shifted from a descriptive towards a more predictive and quantitative approach (Holling, 1966; Lässig, Mustonen, and Walczak, 2017; Marquet et al., 2014; Pickett, 1980; Rossberg et al., 2019; Simberloff, 2004). This shift brought with it the use of mathematical models to describe natural phenomena, such as models that describe the effects species have on each other (Holling, 1966; Levins, 1966; Maynard-Smith, 1978; Servedio et al., 2014). Mathematical descriptions of interactions are “useful fictions” (Box, Luceño, and Carmen Paniagua-Quinones, 2011) in a twofold manner. First, they create a description of how organisms that coincide in space and time affect each other. Almost all known types of interactions can be described in the form of mathematical expressions that faithfully reproduce features of the observed data (Adler et al., 2018; Holland, DeAngelis, and Bronstein, 2002; Holling, 1959; Holt, 1977; Stouffer and Novak, 2021; Vázquez, Morris, and Jordano, 2005; Volterra, 1926; Wood and Thomas, 1999). Second, models are practical tools with which to make predictions beyond the phenomena they describe and thus, provide general insights into how natural systems operate (Stouffer, 2019; Sutherland, 2006).

Model tradeoffs

Models that capture the effect of biotic interactions are abstractions of reality, and abstractions always reflect choices (Levins, 2006). Building models that include all aspects of reality is not only impractical but also unfeasible. Therefore, ecologists and evolutionary biologists have to continuously make choices regarding which variables to include in a model and which to omit (Odenbaugh, 2005). A common assumption when building models is that to

achieve general insights, we should favor simple models (Evans et al., 2013). Indeed there is a general belief in ecology and evolution that a general model should include as little as possible (Holling, 1966; May, 2019; Roughgarden et al., 2018). This belief is often rooted in an implicit philosophical stance that one can not simultaneously maximize generality (i.e., models that apply to more than one system) and realism (i.e., models that produce accurate predictions for a system) (Levins, 1966, 1993).

Inevitably, model building in biology leads to a key question that will, in turn, modify the outcomes achieved by any model: when is a model “realistic” enough (Stouffer, 2019)? The answer to this question will depend on the purpose for which a model is built (Levins, 2006; Odenbaugh, 2005). The classification of biological models and their purposes have been and continue to be widely debated (Evans et al., 2013; Holling, 1966; Levins, 1966, 1993; Lewontin, 1963; May, 2019; Odenbaugh, 2005; Orzack and Sober, 1993; Weisberg, 2006). Overall, it is generally recognized that the purposes of different biological models fall on a continuum (Evans et al., 2013; Levins, 1993; Servedio et al., 2014). On one end of this continuum are models that aim to understand and identify general principles (called strategic models by Holling (1966) and May (2019), or a minimal model of ideas by Roughgarden et al. (2018)). On the other end are models that aim to make detailed quantitative predictions (also called tactical models by Holling (1966) or synthetic models by Roughgarden et al. (2018)). The tradeoffs between generality, realism, and precision at each end of the spectrum have sparked extensive debate among biologists (Levins, 1966, 1993; Orzack and Sober, 1993; Weisberg, 2006).

Models that capture the effect of biotic interactions tend to fall in the spectrum under the category of “demonstration models”, as first defined by (Crick, 1988) and later by (Evans et al., 2013). These types of models are often based on phenomenological descriptions of processes and have the general aim to show that the modeled principles are sufficient to reproduce some phenomena of interest (Crick, 1988; Evans et al., 2013). Demonstration models, however, do not help decide whether the modelled principles are *necessary* (Evans et al., 2013). The task to decide the necessary principles and thus the answer to the when a model is realistic enough becomes the modeler’s responsibility. In many cases, the answer to this question can appear arbitrary or solely determined by the predominant paradigm regarding the studied system. For example, mutualistic interactions between two species can be described by a simple model that assumes a linear functional response (Bascompte, Jordano, and Olesen, 2006a), or by a more realistic model that incorporates saturating effects (Holland, DeAngelis, and Bronstein, 2002). The choice between these two models has substantial implications for predictions related to the coexistence of species and the assembly of communities (Holland, DeAngelis, and Bronstein, 2002). However, there is no consensus on which representation to favor, as the choice is usually defined by the modelers’ particular school of thought

and mathematical convince (Bascompte, Jordano, and Olesen, 2006b; Holland, Okuyama, and DeAngelis, 2006).

The perils of simple models

Always favoring simple models in ecological and evolutionary studies can be problematic from two perspectives. First, the assumption that more complex models do not lead to general insights is seldom tested. For example, most models that capture competitive interactions between plants have the implicit assumption that competitive effects between individuals are always additive and direct (Freckleton and Watkinson, 2001; Kraft, Godoy, and Levine, 2015; Schoener, 1974). However, when models were set up to capture non-additive effects of interactions between individuals of co-occurring species, the evidence overwhelmingly showed that including these levels of biotic complexity was necessary to capture plant interactions accurately (Lai et al., 2021; Martyn et al., 2021; Mayfield and Stouffer, 2017). Thus, in some cases, increasing complexity increases rather than hampers the general insights obtained from models of biotic interactions.

Second, failing to include necessary levels of complexity can hinder our ability to predict how natural communities will react to novel conditions. Predictions of how natural systems will behave in the future are inherently challenging (Sutherland, 2006). Nevertheless, ignoring heterogeneities at various levels can further complicate rather than simplify predictions (D'Andrea, Ostling, and O'Dwyer, 2018). For instance, demographic models tend to treat ecological and evolutionary dynamics separately, despite the general understanding that both processes are often intertwined (Kokko and López-Sepulcre, 2007; MacArthur, 1962). Ignoring eco-evolutionary feedbacks leads to predictions that are inconsistent with empirical data and produce counterintuitive results in novel conditions (Kokko and López-Sepulcre, 2007). Thus, the implicit assumption that good models should include as little as possible should be treated with caution in ecological and evolutionary contexts.

Challenges and consequences of increasing realism

Despite arguments in favor of increasing realism in models of biotic interactions, doing so remains a challenge in many ecological and evolutionary studies. One of those challenges arises from the lack of theoretical frameworks that allow incorporating intricate empirical observations into models (Abrams, 1983, 2001). Such is the case of competition between pollinators that forage for the same resources (Thomson and Page, 2020). An overwhelming amount of empirical evidence shows that pollinators modify their foraging behavior in the presence of other foraging species (Briggs, 2016; Brosi and Briggs, 2013; Inouye,

1978; Morse, 1977; Thompson, Nisbet, and Schmitt, 2006); however, models that incorporate these behavioral changes into population dynamics remain scarce (Thomson and Page, 2020). Furthermore, density-dependent responses could themselves depend on the abiotic conditions pollinators experience, as many studies have shown that environmental conditions can drastically change how pollinators behave and interact with plant species (Briggs et al., 2018; Classen et al., 2020; Cnaani, Thomson, and Papaj, 2006; Heinrich, 1976; Thomson, Peterson, and Harder, 1987; Westphal, Steffan-Dewenter, and Tschardtke, 2006). A coherent framework with which to incorporate both abiotic and biotic drivers into plant–pollinator interactions was lacking. To this end, in [Chapter 2](#) I develop a general framework to show how pollinator functional responses can be used to incorporate biotic and abiotic drivers into models of floral visitation rates. Furthermore, I show the empirical relevance of this framework by parameterizing different models of varying complexity that incorporate pollinator–pollinator interactions and environmental conditions when predicting observed data from a highly controlled foraging chamber experiment. Results from this chapter provide important insights related to our understanding of how species loss and environmental change might affect mutualistic communities.

Another theoretical challenge emerges when alternative models to represent biotic interactions are used interchangeably. Such is the case of phenomenological models of plant competition, where more than one mathematical form can faithfully reproduce empirical data (Bimler et al., 2018; Godoy, Kraft, and Levine, 2014; Godoy and Levine, 2014; Levine and HilleRisLambers, 2009; Mayfield and Stouffer, 2017). The effect biotic and abiotic drivers have on model based predictions can be dramatically different due to uncertainty associated with phenomenological models (Aldebert and Stouffer, 2018; Flora et al., 2011; Jørgensen and Bendricchio, 2001). To understand the interplay between uncertainty and abiotic complexity, in [Chapter 3](#) I introduce a mathematical and statistical framework to simultaneously explore how different phenomenological models of plant competition, environmental context, and parameter uncertainty impact predictions of species coexistence. Additionally, I use this framework to make predictions around a pairwise competition experiment between annual plants, where I show that the effect of abiotic conditions in predictions of coexistence outcomes is not independent of the model formulation used to describe species interactions.

Finally, even when existing studies show that increasing model realism is warranted, understanding exactly how the addition of complexity changes predictions remains a challenge. For instance, theoretical and empirical studies have shown that environmental fluctuations can substantially increase the levels of genetic diversity in populations that experience sexually antagonistic selection (Connallon and Clark, 2012; Connallon, Sharma, and Olito, 2019; Glaser-Schmitt et al., 2021). However, there are no approaches that directly quantify *how* abiotic heterogeneity promotes the maintenance of genetic di-

versity in populations that experience sexual conflict. Hence, in [Chapter 4](#) I adopt an ecological framework to explicitly quantify the contributions of fluctuations in population sizes and selection to allele's growth rates when rare using simulations. I show that environmental fluctuations can help maintain genetic variance in a population by allowing disadvantageous alleles to have positive invasion growth rates, but that their effect depends on the pathway by which each allele is introduced to the population.

Concluding remarks

In this thesis, I propose theoretical and statistical frameworks that allow increasing realism in models of biotic interactions with the aim to understand when higher levels of complexity are justified. Furthermore, I also explore the consequences of increasing model realism in predictions related to diversity maintenance at ecological and evolutionary scales. The individual chapters of this thesis are thematically broad as they are focused on different types of interactions and organisms, but all address in a different way the challenges and consequences of incorporating biotic and abiotic complexity in the study of biotic interactions.

BIBLIOGRAPHY

- Abrams, Peter A (1983). 'Arguments in favor of higher order interactions.' In: *The American Naturalist* 121.6, pp. 887–891.
- (2001). 'Describing and quantifying interspecific interactions: a commentary on recent approaches.' In: *Oikos* 94.2, pp. 209–218.
- Adler, Peter B, Danielle Smull, Karen H Beard, Ryan T Choi, Tucker Furniss, Andrew Kulmatiski, Joan M Meiners, Andrew T Tredennick, and Kari E Veblen (2018). 'Competition and coexistence in plant communities: intraspecific competition is stronger than interspecific competition.' In: *Ecology letters* 21.9, pp. 1319–1329.
- Aldebert, Clement and Daniel B Stouffer (2018). 'Community dynamics and sensitivity to model structure: towards a probabilistic view of process-based model predictions.' In: *Journal of the Royal Society Interface* 15.149, p. 20180741.
- Bascompte, Jordi, Pedro Jordano, and Jens M Olesen (2006a). 'Asymmetric coevolutionary networks facilitate biodiversity maintenance.' In: *Science* 312.5772, pp. 431–433.
- (2006b). 'Response to Comment on " Asymmetric Coevolutionary Networks Facilitate Biodiversity Maintenance".' In: *Science* 313.5795, pp. 1887–1887.
- Bimler, Malyon D., Daniel B. Stouffer, Hao Ran Lai, and Margaret M. Mayfield (2018). 'Accurate predictions of coexistence in natural systems require the inclusion of facilitative interactions and environmental dependency.' en. In: *Journal of Ecology* 106.5, pp. 1839–1852. ISSN: 1365-2745.
- Box, George EP, Alberto Luceño, and Maria del Carmen Paniagua-Quinones (2011). *Statistical control by monitoring and adjustment*. Vol. 700. John Wiley & Sons.
- Briggs, Heather M, Stuart Graham, Callin M Switzer, and Robin Hopkins (2018). 'Variation in context-dependent foraging behavior across pollinators.' In: *Ecology and evolution* 8.16, pp. 7964–7973.
- Briggs, Heather Mae (2016). 'Competitive context drives pollinator behavior: linking foraging plasticity, natural pollen deposition, and plant reproduction.' en. PhD thesis. UC Santa Cruz.

- Broisi, Berry J. and Heather M. Briggs (Aug. 2013). 'Single pollinator species losses reduce floral fidelity and plant reproductive function.' en. In: *Proceedings of the National Academy of Sciences* 110.32, pp. 13044–13048. ISSN: 0027-8424, 1091-6490.
- Capdevila, Pol, Iain Stott, Imma Oliveras Menor, Daniel B Stouffer, Rafael LG Raimundo, Hannah White, Matthew Barbour, and Roberto Salguero-Gómez (2021). *Reconciling resilience across ecological systems, species and subdisciplines*.
- Chase, Jonathan M and Mathew A Leibold (2009). *Ecological niches*. University of Chicago Press.
- Chesson, Peter (2000). 'Mechanisms of maintenance of species diversity.' In: *Annual review of Ecology and Systematics* 31.1, pp. 343–366.
- Classen, Alice, Connal D Eardley, Andreas Hemp, Marcell K Peters, Ralph S Peters, Axel Ssymank, and Ingolf Steffan-Dewenter (2020). 'Specialization of plant–pollinator interactions increases with temperature at Mt. Kilimanjaro.' In: *Ecology and evolution* 10.4, pp. 2182–2195.
- Cnaani, Jonathan, James D. Thomson, and Daniel R. Papaj (2006). 'Flower Choice and Learning in Foraging Bumblebees: Effects of Variation in Nectar Volume and Concentration.' en. In: *Ethology* 112.3, pp. 278–285. ISSN: 1439-0310.
- Connallon, Tim and Andrew G Clark (2012). 'A general population genetic framework for antagonistic selection that accounts for demography and recurrent mutation.' In: *Genetics* 190.4, pp. 1477–1489.
- Connallon, Tim, Shefali Sharma, and Colin Olito (Jan. 2019). 'Evolutionary Consequences of Sex-Specific Selection in Variable Environments: Four Simple Models Reveal Diverse Evolutionary Outcomes.' In: *The American Naturalist* 193.1, pp. 93–105. ISSN: 0003-0147.
- Crick, F (1988). 'What Mad Pursuit: A Personal View of Scientific Discovery. Basic Books, New York.' In:
- D'Andrea, Rafael, Annette Ostling, and James P O'Dwyer (2018). 'Translucent windows: How uncertainty in competitive interactions impacts detection of community pattern.' In: *Ecology Letters* 21.6, pp. 826–835.
- Evans, Matthew R, Volker Grimm, Karin Johst, Tarja Knuuttila, Rogier De Langhe, Catherine M Lessells, Martina Merz, Maureen A O'Malley, Steve H Orzack, Michael Weisberg, et al. (2013). 'Do simple models lead to generality in ecology?' In: *Trends in ecology & evolution* 28.10, pp. 578–583.
- Flora, Cordoleani, Nerini David, Gauduchon Mathias, Morozov Andrew, and Poggiale Jean-Christophe (Aug. 2011). 'Structural sensitivity of biological models revisited.' In: *Journal of Theoretical Biology* 283.1, pp. 82–91. ISSN: 0022-5193.

- Freckleton, RP and AR Watkinson (2001). 'Predicting competition coefficients for plant mixtures: reciprocity, transitivity and correlations with life-history traits.' In: *Ecology Letters* 4.4, pp. 348–357.
- Gause, Georgii Frantsevitch (1934). 'Experimental analysis of Vito Volterra's mathematical theory of the struggle for existence.' In: *Science* 79.2036, pp. 16–17.
- Glaser-Schmitt, Amanda, Meike J Wittmann, Timothy JS Ramnarine, and John Parsch (2021). 'Sexual antagonism, temporally fluctuating selection, and variable dominance affect a regulatory polymorphism in *Drosophila melanogaster*.' In: *Molecular Biology and Evolution* 38.11, pp. 4891–4907.
- Godoy, Oscar, Lorena Gómez-Aparicio, Luis Matías, Ignacio M Pérez-Ramos, and Eric Allan (2020). 'An excess of niche differences maximizes ecosystem functioning.' In: *Nature communications* 11.1, pp. 1–10.
- Godoy, Oscar, Nathan J. B. Kraft, and Jonathan M. Levine (2014). 'Phylogenetic relatedness and the determinants of competitive outcomes.' en. In: *Ecology Letters* 17.7, pp. 836–844. ISSN: 1461-0248.
- Godoy, Oscar and Jonathan M. Levine (2014). 'Phenology effects on invasion success: insights from coupling field experiments to coexistence theory.' en. In: *Ecology* 95.3, pp. 726–736. ISSN: 1939-9170.
- Heinrich, Bernd (Aug. 1976). 'Resource Partitioning Among Some Eusocial Insects: Bumblebees.' en. In: *Ecology* 57.5, pp. 874–889. ISSN: 00129658.
- HilleRisLambers, Janneke, Peter B Adler, W Stanley Harpole, Jonathan M Levine, and Margaret M Mayfield (2012). 'Rethinking community assembly through the lens of coexistence theory.' In: *Annual review of ecology, evolution, and systematics* 43, pp. 227–248.
- Holland, J Nathaniel, Donald L DeAngelis, and Judith L Bronstein (2002). 'Population dynamics and mutualism: functional responses of benefits and costs.' In: *The American Naturalist* 159.3, pp. 231–244.
- Holland, J Nathaniel, Toshinori Okuyama, and Donald L DeAngelis (2006). 'Comment on "Asymmetric Coevolutionary Networks Facilitate Biodiversity Maintenance".' In: *Science* 313.5795, pp. 1887–1887.
- Holling, Crawford S (1959). 'Some characteristics of simple types of predation and parasitism1.' In: *The Canadian Entomologist* 91.7, pp. 385–398.
- (1966). 'The strategy of building models of complex ecological systems.' In: *Systems analysis in ecology*, pp. 195–214.
- Holt, Robert D (1977). 'Predation, apparent competition, and the structure of prey communities.' In: *Theoretical population biology* 12.2, pp. 197–229.

- Inouye, David W. (1978). 'Resource Partitioning in Bumblebees: Experimental Studies of Foraging Behavior.' en. In: *Ecology* 59.4, pp. 672–678. ISSN: 1939-9170.
- Jørgensen, Sven Erik and Giuseppe Bendoricchio (2001). *Fundamentals of Ecological Modelling*. Vol. 21. Elsevier.
- Kokko, Hanna and Andrés López-Sepulcre (2007). 'The ecogenetic link between demography and evolution: can we bridge the gap between theory and data?' In: *Ecology Letters* 10.9, pp. 773–782.
- Kraft, Nathan JB, Oscar Godoy, and Jonathan M Levine (2015). 'Plant functional traits and the multidimensional nature of species coexistence.' In: *Proceedings of the National Academy of Sciences* 112.3, pp. 797–802.
- Lai, Hao Ran, Kwek Yan Chong, Alex Thiam Koon Yee, Margaret M Mayfield, and Daniel B Stouffer (2021). 'Non-additive biotic interactions improve predictions of tropical tree growth and impact community size structure.' In: *bioRxiv*, pp. 2020–09.
- Lässig, Michael, Ville Mustonen, and Aleksandra M Walczak (2017). 'Predicting evolution.' In: *Nature ecology & evolution* 1.3, pp. 1–9.
- Lawton, John H (1999). 'Are there general laws in ecology?' In: *Oikos*, pp. 177–192.
- Levine, Jonathan M and Janneke HilleRisLambers (2009). 'The importance of niches for the maintenance of species diversity.' In: *Nature* 461.7261, pp. 254–257.
- Levins, Richard (1966). 'The strategy of model building in population biology.' In: *American scientist* 54.4, pp. 421–431.
- (1993). 'A response to Orzack and Sober: formal analysis and the fluidity of science.' In: *The Quarterly Review of Biology* 68.4, pp. 547–555.
- (2006). 'Strategies of abstraction.' In: *Biology and Philosophy* 21.5, pp. 741–755.
- Lewontin, Richard C (1963). 'Models, mathematics and metaphors.' In: *Synthese*, pp. 222–244.
- MacArthur, Robert H (1962). 'Some generalized theorems of natural selection.' In: *Proceedings of the National Academy of Sciences of the United States of America* 48.11, p. 1893.
- MacArthur, Robert and Richard Levins (1967). 'The limiting similarity, convergence, and divergence of coexisting species.' In: *The american naturalist* 101.921, pp. 377–385.
- Marquet, Pablo A, Andrew P Allen, James H Brown, Jennifer A Dunne, Brian J Enquist, James F Gillooly, Patricia A Gowaty, Jessica L Green, John Harte, Steve P Hubbell, et al. (2014). 'On theory in ecology.' In: *BioScience* 64.8, pp. 701–710.

- Martyn, Trace E, Daniel B Stouffer, Oscar Godoy, Ignasi Bartomeus, Abigail I Pastore, and Margaret M Mayfield (2021). 'Identifying "Useful" Fitness Models: Balancing the Benefits of Added Complexity with Realistic Data Requirements in Models of Individual Plant Fitness.' In: *The American Naturalist* 197.4, pp. 415–433.
- May, Robert M (1972). 'Will a large complex system be stable?' In: *Nature* 238.5364, pp. 413–414.
- (2019). *Stability and complexity in model ecosystems*. Princeton university press.
- Mayfield, Margaret M and Daniel B Stouffer (2017). 'Higher-order interactions capture unexplained complexity in diverse communities.' In: *Nature ecology & evolution* 1.3, pp. 1–7.
- Maynard-Smith, John (1978). *Models in ecology*. CUP Archive.
- Morse, D. H. (Aug. 1977). 'Resource Partitioning in Bumble Bees: The Role of Behavioral Factors.' en. In: *Science* 197.4304, pp. 678–680. ISSN: 0036-8075, 1095-9203.
- Odenbaugh, Jay (2005). 'Idealized, inaccurate but successful: A pragmatic approach to evaluating models in theoretical ecology.' In: *Biology and Philosophy* 20.2-3, pp. 231–255.
- Orzack, Steven Hecht and Elliott Sober (1993). 'A critical assessment of Levins's The strategy of model building in population biology (1966).' In: *The Quarterly Review of Biology* 68.4, pp. 533–546.
- Pickett, Steward TA (1980). 'Non-equilibrium coexistence of plants.' In: *Bulletin of the Torrey Botanical Club*, pp. 238–248.
- Rossberg, Axel G, György Barabás, Hugh P Possingham, Mercedes Pascual, Pablo A Marquet, Cang Hui, Matthew R Evans, and Géza Meszén (2019). 'Let's train more theoretical ecologists—here is why.' In: *Trends in ecology & evolution* 34.9, pp. 759–762.
- Roughgarden, Jonathan, Aviv Bergman, Sharoni Shafir, and Charles Taylor (2018). 'Adaptive Computation in Ecology and.' In: *Adaptive Individuals In Evolving Populations: Models And Algorithms*, p. 25.
- Saavedra, Serguei, Rudolf P Rohr, Jordi Bascompte, Oscar Godoy, Nathan JB Kraft, and Jonathan M Levine (2017). 'A structural approach for understanding multispecies coexistence.' In: *Ecological Monographs* 87.3, pp. 470–486.
- Schoener, Thomas W (1974). 'Some methods for calculating competition coefficients from resource-utilization spectra.' In: *The American Naturalist* 108.961, pp. 332–340.
- Servedio, Maria R, Yaniv Brandvain, Sumit Dhole, Courtney L Fitzpatrick, Emma E Goldberg, Caitlin A Stern, Jeremy Van Cleve, and D Justin Yeh (2014). 'Not just a theory—the utility of mathematical models in evolutionary biology.' In: *PLoS biology* 12.12, e1002017.

- Simberloff, Daniel (2004). 'Community Ecology: Is It Time to Move On? (An American Society of Naturalists Presidential Address).' In: *The American Naturalist* 163.6, pp. 787–799.
- Song, Chuliang and Serguei Saavedra (2018). 'Will a small randomly assembled community be feasible and stable?' In: *Ecology* 99.3, pp. 743–751.
- Stouffer, Daniel B (2019). 'All ecological models are wrong, but some are useful.' In: *Journal of Animal Ecology* 88.2, pp. 192–195.
- Stouffer, Daniel B and Mark Novak (2021). 'Hidden layers of density dependence in consumer feeding rates.' In: *Ecology Letters* 24.3, pp. 520–532.
- Sutherland, William J (2006). 'Predicting the ecological consequences of environmental change: a review of the methods.' In: *Journal of Applied Ecology* 43.4, pp. 599–616.
- Thompson, Andrew R., Roger M. Nisbet, and Russell J. Schmitt (Nov. 2006). 'Dynamics of mutualist populations that are demographically open.' en. In: *Journal of Animal Ecology* 75.6, pp. 1239–1251. ISSN: 0021-8790, 1365-2656.
- Thompson, John N (1999). 'The evolution of species interactions.' In: *Science* 284.5423, pp. 2116–2118.
- (2014). *Interaction and coevolution*. University of Chicago Press.
- Thomson, Diane M and Maureen L Page (Apr. 2020). 'The importance of competition between insect pollinators in the Anthropocene.' en. In: *Current Opinion in Insect Science*. Ecology • Parasites/Parasitoids/Biological control 38, pp. 55–62. ISSN: 2214-5745.
- Thomson, J. D., S. C. Peterson, and L. D. Harder (Jan. 1987). 'Response of traplining bumble bees to competition experiments: shifts in feeding location and efficiency.' en. In: *Oecologia* 71.2, pp. 295–300. ISSN: 0029-8549, 1432-1939.
- Turnbull, Lindsay Ann, Jonathan M Levine, Michel Loreau, and Andy Hector (2013). 'Coexistence, niches and biodiversity effects on ecosystem functioning.' In: *Ecology letters* 16, pp. 116–127.
- Vázquez, Diego P, William F Morris, and Pedro Jordano (2005). 'Interaction frequency as a surrogate for the total effect of animal mutualists on plants.' In: *Ecology letters* 8.10, pp. 1088–1094.
- Volterra, Vito (1926). 'Fluctuations in the abundance of a species considered mathematically.' In: *Nature* 118.2972, pp. 558–560.
- Weisberg, Michael (2006). 'Forty years of 'the strategy': Levins on model building and idealization.' In: *Biology and Philosophy* 21.5, pp. 623–645.
- Westphal, Catrin, Ingolf Steffan-Dewenter, and Teja Tscharntke (Aug. 2006). 'Bumblebees experience landscapes at different spatial scales:

- possible implications for coexistence.' en. In: *Oecologia* 149.2, pp. 289–300. ISSN: 0029-8549, 1432-1939.
- Wood, Simon N and Matthew B Thomas (1999). 'Super-sensitivity to structure in biological models.' In: *Proceedings of the Royal Society of London. Series B: Biological Sciences* 266.1419, pp. 565–570.
- Wootton, KL and DB Stouffer (2016). 'Many weak interactions and few strong; food-web feasibility depends on the combination of the strength of species' interactions and their correct arrangement.' In: *Theoretical Ecology* 9.2, pp. 185–195.

THE CONTEXT-DEPENDENCY OF POLLINATOR INTERFERENCE

This is your bee foraging manuscript please compile

THE INTERPLAY OF ENVIRONMENTAL CONDITIONS,
PARAMETER SENSITIVITY AND STRUCTURAL
SENSITIVITY IN PREDICTIONS OF SPECIES
COEXISTENCE

This is your bayesian competition manauscript

COEXISTENCE OF ALLELES

This is your coexistence of alleles manuscript

CONCLUSION

*Where is the rest of the world?
That is the question we must always ask about any model:
where is the rest of the world?*

— Levins (2006)

In this thesis, I show how to incorporate biotic and abiotic complexity in models of biotic interactions to increase model realism. Furthermore, I provide direct evidence that many models used to describe biotic interactions are oversimplistic since they fail to capture dynamics accurately by *a priori* ignoring abiotic and biotic factors. Throughout this thesis, I also show that increasing realism in models of biotic interactions has important repercussions on our understanding and predictions about the maintenance of diversity at ecological and evolutionary scales.

SUMMARY OF RESULTS

In [Chapter 2](#) I found that the abundance of co-foragers can fundamentally change the number of visits pollinators make. These results imply that it is necessary to account for the density of species other than the focal pair to characterize plant-pollinator interactions accurately. However, results from this chapter also show that the environmental context pollinators experience mediates density-dependent responses to co-foraging species. Thus, abiotic drivers can modify the number of visits made by pollinators through both density-independent and density-dependent responses. These two types of responses can cause the same environmental context to have opposite effects on floral visits. Such is the case of high resource abundance in our foraging experiment. Additionally, in this chapter, I show that pollinators do not respond equally to all co-foraging species. Therefore the effects of biotic and abiotic drivers depend on the identity of the interacting species. Results from this chapter clearly show that including these levels of complexity in a model of floral visits is justified, despite the increasing number of parameters necessary to fit such a model. Since floral visitation is a good predictor of the strength of plant-pollinator interactions (Vázquez, Morris, and Jordano, 2005; Vázquez et al., 2012) my results demonstrate that failing to account for biotic and abiotic

CONCLUSION

complexity can result in misleading estimations of the level of interdependence of animal and plant populations.

In [Chapter 3](#) I found that accounting for the abiotic context where interactions occur can fundamentally change predictions of species coexistence. While other studies have previously shown that predictions of coexistence between plant species can be context-dependent (Bimler et al., 2018; Lanuza, Bartomeus, and Godoy, 2018), my results are the first to show that the estimated effect of the abiotic context depends on the model used to describe species interactions. Thus, my results show that incorporating abiotic complexity in models of biotic interactions is far from straightforward, as different phenomenological models can enhance or diminish its effect. Additionally, parameter uncertainty can further hinder the interpretation of the effect abiotic drivers have on predictions. For instance, predictions showed that in the *woody* environment our focal species were unlikely to coexist, however, the species predicted to be competitively excluded varied across posterior draws. Therefore, my results show that robust predictions of species coexistence need to consider the abiotic context where interactions occur *and* different sources of uncertainty associated with phenomenological models.

Finally, in [Chapter 4](#) I found that environmental fluctuations can substantially increase the level of polymorphism in populations that experience sexually antagonistic selection. Perhaps most importantly, the results of this chapter show that environmental fluctuations can maintain disadvantageous alleles in a population by contributing positively to their growth rates when rare. However, the positive contributions of fluctuations depended on the pathway by which each allele was introduced into the population. Thus, I show that abiotic heterogeneity must be coupled with aspects of the evolutionary dynamics of the populations involved to maintain genetic diversity. This chapter highlights that not all types of abiotic drivers have the same effects on the populations involved. For instance, fluctuations in selection contributed positively to allele's invasion growth rates when fluctuations were positively correlated. In contrast, fluctuations in population sizes needed to be negatively correlated to have positive contributions. Therefore, my results show the importance of not only investigating *if* environmental drivers change predictions but also *how* they do it.

GENERAL IMPLICATIONS

Increasing model realism can be achieved in multiple ways. One of them is to add independent variables to a model that represent previously ignored aspects of the real world (Evans et al., 2013; Orzack and Sober, 1993). For example, in [Chapter 2](#), I increased realism by adding variables that accounted for the densities of co-foragers to a model of floral visits. Other methods include adding a new link to variables already present or imposing bounds to some

aspects of the model (Levins, 1993). For example, in [Chapter 3](#) I assumed unlimited growth was unrealistic and imposed abundance constraints when predicting species coexistence. Importantly, whether any of these methods increase the correspondence between model and phenomena of interest can not be evaluated *a priori*. For instance, in [Chapter 2](#) it may have been the case that abiotic conditions had no effect on how a pollinator forages the presence of other species. Then, a model that included density-dependent effects to environmental conditions would have been over-parameterized. However, this assessment can only be done *after* the fit a more complex model.

In this thesis, I show that models of biotic interactions aimed to make predictions regarding diversity maintenance tend to be oversimplistic. The implicit assumption that general insights can only be achieved with simple models (Holling, 1966; May, 2019; Roughgarden et al., 2018) has led to the automatic omission of biotic and abiotic heterogeneities in many models of biotic interactions. However, scientific generality is not the same as mathematical generality (Levins, 1993). For instance, to build a simple model in mathematical terms, we should include as little as possible (Orzack, 2012). In contrast, building a simple model in biological terms might entail accounting for abiotic and biotic dependency.

A key simplifying assumption in many models of biotic interactions is that species pairs is the relevant unit of study (Levine et al., 2017). However, many theoretical and empirical studies have shown that this assumption is likely a major oversimplification (Abrams, 1983; Billick and Case, 1994; Letten and Stouffer, 2019; Mayfield and Stouffer, 2017). Various biological mechanisms can cause individuals of the same or different species to modify the interaction between a focal pair, a phenomenon broadly referred to as higher order interactions (Kleinhesselink, Kraft, and Levine, 2019). One of these mechanisms, interaction modification, arises when an intermediary species induces behavioral changes in one or both of the species in the focal pair, thereby modifying their interaction (Adler and Morris, 1994; Billick and Case, 1994). In [Chapter 2](#), I show that interaction modifications are likely to occur in mutualistic communities by behavioral responses of pollinators to other foraging species. Evidence that shows higher order interactions have non-negligible effect in natural communities continues to accumulate (Lai et al., 2020; Levine et al., 2017; Martyn et al., 2021; Mayfield and Stouffer, 2017). Therefore, ignoring biotic complexity in favor of mathematical simplicity can be detrimental to understanding how multispecies communities are assembled and maintained.

The environmental dependence of biotic interactions has broad empirical and theoretical support (Bimler et al., 2018; Callaway et al., 2002; Chamberlain, Bronstein, and Rudgers, 2014; Chesson, 2000; Lanuza, Bartomeus, and Godoy, 2018; Tylianakis et al., 2008). Furthermore, the importance of heterogeneous environments in the maintenance of diversity has been shown at ecological (Amarasekare, 2003; Kneitel and Chase, 2004) and evolutionary scales (Connallon, Sharma, and Olito, 2019; Dean, 2005; Ellner and Hairston Jr, 1994). Yet, most

CONCLUSION

models used to make predictions regarding diversity maintenance tend to keep changes in biotic interactions due to environmental effects implicit or treat them as constants (but see Bimler et al. (2018), or Connallon, Sharma, and Olito (2019)). In Chapter 3 and Chapter 4 I address two different challenges when accounting for environmental dependency in models of biotic interactions: the interplay between model uncertainty and environmental heterogeneity, and the interpretation of environmental effects in model predictions. While in natural systems, as opposed to experimental systems or simulations, it is challenging to determine exactly what variables constitute “the environment”, an inability to account for abiotic dependence prevents ecologists and evolutionary biologists to correctly identify the drivers of diversity maintenance (Connallon, Sharma, and Olito, 2019; Freckleton, Watkinson, and Rees, 2009).

FUTURE DIRECTIONS

While it might be tempting to argue that increasing model realism is a Sisyphean endeavor, there are limits to the level of complexity that can be added to a model without losing its usefulness. Where those limits are and whether a model is realistic enough will depend on the state of science at the time (Levins, 1993). In this thesis, I argue that automatically excluding biotic and abiotic factors from models of biotic interactions in favor of mathematical simplicity obstacles our understanding of how natural systems operate. The level of realism needed to describe biotic interactions accurately will undoubtedly depend on the system studied, as not all types of interactions are equally likely to be affected by the biotic and abiotic context they are embedded in (Chamberlain, Bronstein, and Rudgers, 2014). Nonetheless, scientific progress requires that we acknowledge and explore this complexity. This has been and continues to be done by studies that compare models of varying levels of complexity to understand when its inclusion is warranted (Bimler et al., 2018; Lai et al., 2020; Martyn et al., 2021; Weiss-Lehman et al., 2021), studies that investigate the biological rationale for why certain levels of complexity should be included in models (Abrams, 1983; Abrams and Ginzburg, 2000; AlAdwani and Saavedra, 2019; Letten and Stouffer, 2019; Stouffer and Novak, 2021), and studies that show how to account for unmeasured biotic and abiotic drivers (D’Andrea, Ostling, and O’Dwyer, 2018; Song et al., 2020).

Finally, a key aspect missing from most ecological and evolutionary research focused on diversity maintenance is the bidirectional link between organisms and the environment. Extensive research, including this thesis, has been done to disentangle the effects of abiotic and biotic factors on the performance of organisms. The reciprocal phenomenon, the reaction and evolution of the environment in response to the organisms embedded in it, is generally ignored in many ecological and evolutionary studies (Laland, Odling-Smee, and Feldman, 1999; Levins and Lewontin, 1980). However, including the feedback loop

between organism and environment is essential to fully understand the interplay between biotic and abiotic variables in natural communities (Callaway, 2007; Hastings et al., 2007). Additionally, theoretical studies suggest that this feedback process can fundamentally change predictions related to diversity maintenance in natural communities (Kylafis and Loreau, 2008, 2011). Thus, subsequent efforts should examine how biotic and abiotic complexity jointly modifies the responses of organisms and the environment. This thesis offers some theoretical tools for ecologists and evolutionary biologists to explore the first part of this puzzle.

BIBLIOGRAPHY

- Abrams, Peter A (1983). 'Arguments in favor of higher order interactions.' In: *The American Naturalist* 121.6, pp. 887–891.
- Abrams, Peter A. and Lev R. Ginzburg (Aug. 2000). 'The nature of predation: prey dependent, ratio dependent or neither?' In: *Trends in Ecology & Evolution* 15.8, pp. 337–341. ISSN: 0169-5347.
- Adler, Frederick R. and William F. Morris (Sept. 1994). 'A General Test for Interaction Modification.' en. In: *Ecology* 75.6, pp. 1552–1559. ISSN: 00129658.
- AlAdwani, Mohammad and Serguei Saavedra (2019). 'Is the addition of higher-order interactions in ecological models increasing the understanding of ecological dynamics?' In: *Mathematical Biosciences* 315, p. 108222.
- Amarasekare, Priyanga (2003). 'Competitive coexistence in spatially structured environments: a synthesis.' In: *Ecology letters* 6.12, pp. 1109–1122.
- Billick, Ian and Ted J. Case (1994). 'Higher Order Interactions in Ecological Communities: What Are They and How Can They be Detected?' en. In: *Ecology* 75.6, pp. 1529–1543. ISSN: 1939-9170.
- Bimler, Malyon D., Daniel B. Stouffer, Hao Ran Lai, and Margaret M. Mayfield (2018). 'Accurate predictions of coexistence in natural systems require the inclusion of facilitative interactions and environmental dependency.' en. In: *Journal of Ecology* 106.5, pp. 1839–1852. ISSN: 1365-2745.
- Callaway, Ragan M (2007). *Positive interactions and interdependence in plant communities*. Springer.
- Callaway, Ragan M. et al. (June 2002). 'Positive interactions among alpine plants increase with stress.' en. In: *Nature* 417.6891, pp. 844–848. ISSN: 1476-4687.
- Chamberlain, Scott A., Judith L. Bronstein, and Jennifer A. Rudgers (2014). 'How context dependent are species interactions?' en. In: *Ecology Letters* 17.7, pp. 881–890. ISSN: 1461-0248.
- Chesson, Peter (2000). 'Mechanisms of maintenance of species diversity.' In: *Annual review of Ecology and Systematics* 31.1, pp. 343–366.

- Connallon, Tim, Shefali Sharma, and Colin Olito (Jan. 2019). 'Evolutionary Consequences of Sex-Specific Selection in Variable Environments: Four Simple Models Reveal Diverse Evolutionary Outcomes.' In: *The American Naturalist* 193.1, pp. 93–105. ISSN: 0003-0147.
- D'Andrea, Rafael, Annette Ostling, and James P O'Dwyer (2018). 'Translucent windows: How uncertainty in competitive interactions impacts detection of community pattern.' In: *Ecology Letters* 21.6, pp. 826–835.
- Dean, Antony M (2005). 'Protecting haploid polymorphisms in temporally variable environments.' In: *Genetics* 169.2, pp. 1147–1156.
- Ellner, Stephen and Nelson G Hairston Jr (1994). 'Role of overlapping generations in maintaining genetic variation in a fluctuating environment.' In: *The American Naturalist* 143.3, pp. 403–417.
- Evans, Matthew R, Volker Grimm, Karin Johst, Tarja Knuuttila, Rogier De Langhe, Catherine M Lessells, Martina Merz, Maureen A O'Malley, Steve H Orzack, Michael Weisberg, et al. (2013). 'Do simple models lead to generality in ecology?' In: *Trends in ecology & evolution* 28.10, pp. 578–583.
- Freckleton, Robert P, Andrew R Watkinson, and Mark Rees (2009). 'Measuring the importance of competition in plant communities.' In: *Journal of ecology* 97.3, pp. 379–384.
- Hastings, Alan, James E Byers, Jeffrey A Crooks, Kim Cuddington, Clive G Jones, John G Lambrinos, Theresa S Talley, and William G Wilson (2007). 'Ecosystem engineering in space and time.' In: *Ecology letters* 10.2, pp. 153–164.
- Holling, Crawford S (1966). 'The strategy of building models of complex ecological systems.' In: *Systems analysis in ecology*, pp. 195–214.
- Kleinhesselink, Andrew R, Nathan JB Kraft, and Jonathan M Levine (2019). 'Mechanisms underlying higher order interactions: from quantitative definitions to ecological processes.' In: *bioRxiv*, p. 857920.
- Kneitel, Jamie M and Jonathan M Chase (2004). 'Trade-offs in community ecology: linking spatial scales and species coexistence.' In: *Ecology letters* 7.1, pp. 69–80.
- Kylafis, Grigoris and Michel Loreau (2008). 'Ecological and evolutionary consequences of niche construction for its agent.' In: *Ecology letters* 11.10, pp. 1072–1081.
- (Feb. 2011). 'Niche construction in the light of niche theory: Niche construction in light of niche theory.' en. In: *Ecology Letters* 14.2, pp. 82–90. ISSN: 1461023X.
- Lai, Hao Ran, Kwek Yan Chong, Margaret M Mayfield, and Daniel B Stouffer (2020). 'The role of higher-order biotic interactions on tropical tree growth.' In: *bioRxiv*.

- Laland, Kevin N, F John Odling-Smee, and Marcus W Feldman (1999). 'Evolutionary consequences of niche construction and their implications for ecology.' In: *Proceedings of the National Academy of Sciences* 96.18, pp. 10242–10247.
- Lanuza, Jose B., Ignasi Bartomeus, and Oscar Godoy (2018). 'Opposing effects of floral visitors and soil conditions on the determinants of competitive outcomes maintain species diversity in heterogeneous landscapes.' en. In: *Ecology Letters* 21.6, pp. 865–874. ISSN: 1461-0248.
- Letten, Andrew D. and Daniel B. Stouffer (2019). 'The mechanistic basis for higher-order interactions and non-additivity in competitive communities.' en. In: *Ecology Letters* 22.3, pp. 423–436. ISSN: 1461-0248.
- Levine, Jonathan M., Jordi Bascompte, Peter B. Adler, and Stefano Allesina (June 2017). 'Beyond pairwise mechanisms of species coexistence in complex communities.' en. In: *Nature* 546.7656, pp. 56–64. ISSN: 1476-4687.
- Levins, Richard (1993). 'A response to Orzack and Sober: formal analysis and the fluidity of science.' In: *The Quarterly Review of Biology* 68.4, pp. 547–555.
- (2006). 'Strategies of abstraction.' In: *Biology and Philosophy* 21.5, pp. 741–755.
- Levins, Richard and Richard Lewontin (Jan. 1980). 'Dialectics and reductionism in ecology.' en. In: *Synthese* 43.1, pp. 47–78. ISSN: 1573-0964.
- Martyn, Trace E, Daniel B Stouffer, Oscar Godoy, Ignasi Bartomeus, Abigail I Pastore, and Margaret M Mayfield (2021). 'Identifying "Useful" Fitness Models: Balancing the Benefits of Added Complexity with Realistic Data Requirements in Models of Individual Plant Fitness.' In: *The American Naturalist* 197.4, pp. 415–433.
- May, Robert M (2019). *Stability and complexity in model ecosystems*. Princeton university press.
- Mayfield, Margaret M and Daniel B Stouffer (2017). 'Higher-order interactions capture unexplained complexity in diverse communities.' In: *Nature ecology & evolution* 1.3, pp. 1–7.
- Orzack, Steven Hecht (2012). 'The philosophy of modelling or does the philosophy of biology have any use?' In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 367.1586, pp. 170–180.
- Orzack, Steven Hecht and Elliott Sober (1993). 'A critical assessment of Levins's The strategy of model building in population biology (1966).' In: *The Quarterly Review of Biology* 68.4, pp. 533–546.
- Roughgarden, Jonathan, Aviv Bergman, Sharoni Shafir, and Charles Taylor (2018). 'Adaptive Computation in Ecology and.' In: *Adaptive Individuals In Evolving Populations: Models And Algorithms*, p. 25.

- Song, Chuliang, Sarah Von Ahn, Rudolf P. Rohr, and Serguei Saavedra (Jan. 2020). 'Towards a Probabilistic Understanding About the Context-Dependency of Species Interactions.' en. In: *Trends in Ecology & Evolution*. ISSN: 0169-5347.
- Stouffer, Daniel B and Mark Novak (2021). 'Hidden layers of density dependence in consumer feeding rates.' In: *Ecology Letters* 24.3, pp. 520–532.
- Tylianakis, Jason M, Raphael K Didham, Jordi Bascompte, and David A Wardle (2008). 'Global change and species interactions in terrestrial ecosystems.' In: *Ecology letters* 11.12, pp. 1351–1363.
- Vázquez, Diego P, William F Morris, and Pedro Jordano (2005). 'Interaction frequency as a surrogate for the total effect of animal mutualists on plants.' In: *Ecology letters* 8.10, pp. 1088–1094.
- Vázquez, Diego P., Silvia B. Lomáscolo, M. Belén Maldonado, Natacha P. Chacoff, Jimena Dorado, Erica L. Stevani, and Nydia L. Vitale (2012). 'The strength of plant–pollinator interactions.' en. In: *Ecology* 93.4, pp. 719–725. ISSN: 1939-9170.
- Weiss-Lehman, Christopher P, Chhaya M Werner, Catherine H Bowler, Lauren M Hallett, Margaret M Mayfield, Oscar Godoy, Lina Aoyama, György Barabás, Chengjin Chu, Emma Ladouceur, et al. (2021). 'Disentangling key species interactions in diverse and heterogeneous communities: A Bayesian sparse modeling approach.' In: *bioRxiv*.

COLOPHON

This document was typeset using the typographical look-and-feel `classicthesis` developed by André Miede. The style was inspired by Robert Bringhurst’s seminal book on typography “*The Elements of Typographic Style*”. `classicthesis` is available for both \LaTeX and \LyX :

<https://bitbucket.org/amiede/classicthesis/>

Final Version as of November 23, 2021 (Version 1.0).