

Problem 1. Let $f : A \rightarrow B$. Then:

- a) f is injective if and only if it has a left inverse.
- b) f is surjective if and only if it has a right inverse.
- c) f is bijective if and only if it has a left and right inverse.
- d) If $|A| = |B| = n \in \mathbb{Z}_{\geq 0}$ then f is injective if and only if f is surjective if and only if f is bijective.

Proof.

- a) Suppose that f is injective. This implies that $f^{-1}(f(a)) = \{a\}$ for all $a \in A$. Define $g : B \rightarrow A$ as $g(b) = f^{-1}(a)$ where $b \in f(A)$. If $b \notin f(A)$ then send it to any arbitrary $a \in A$. Then $g \circ f(a) = f^{-1}(f(a)) = a$ which implies that $g \circ f = id_A$ and that g is a left inverse of f .

Now suppose that there exists a function $g : B \rightarrow A$ such that $g \circ f = id_A$. Let $a_0, a_1 \in A$ such that $f(a_0) = f(a_1)$. Then $g \circ f(a_0) = a_0 = a_1 = g \circ f(a_1)$ which is the definition of injectivity.

Therefore f is injective if and only if it has a left inverse.

- b) Suppose that f is surjective. Then given $b \in B$ there exists an $a \in A$ such that $f(a) = b$. Define a function $g : B \rightarrow A$ via $g(b) = a$ where a fulfills $f(a) = b$. Then $f \circ g(b) = b$ by definition which implies that g is a right inverse of f .

Now suppose that there exists a function $g : B \rightarrow A$ such that $f \circ g = id_B$. Then given $b \in B$ let $a = g(b)$. Then $f(a) = f \circ g(b) = b$. Since this holds for all elements of b f is surjective.

Therefore f is surjective if and only if it has a right inverse.

- c) Suppose that f is a bijection. Then it is both injective and surjective which by the previous statements in the proposition implies that f has both a left and right inverse.

Otherwise suppose that f has a left and right inverse. Then via the previous statements in the proposition we know that f is both injective and surjective and thus a bijection.

To show that the left and right inverse are unique let g, h be a left and right inverse for f respectively. Then

$$g = g \circ id_B = g \circ (f \circ h) = (g \circ f) \circ h = id_A \circ h = h$$

Therefore f is a bijection if and only if it has a left and right inverse. Moreover these inverses are equal.

- d) Suppose that $|A| = |B| = 1$. Then there is only one function $f : A \rightarrow B$ defined as $f(a_0) = b_0$. As such the function f is injective, surjective and bijective.

Next assume for sets of size n that a function is injective if and only if it is surjective if and only if it is bijective. Let $|A| = |B| = n + 1$, and without loss of generality let f be injective. Select a_n and $b_n = f(a_n)$ and look at $f|_{A \setminus \{a_n\}}$. This will preserve injectivity since we removed paired items. This implies that $f|_{A \setminus \{a_n\}}$ is injective, surjective, and bijective via our inductive hypothesis. Reintroduce the pair (a_n, b_n) to the function $f|_{A \setminus \{a_n\}}$. This will preserve injectivity as only a_n maps to b_n , surjectivity because b_n is mapped to by a_n , and as such f is bijective.

Therefore via induction, if $|A| = |B| = n \in \mathbb{Z}_{>0}$ then a function $f : A \rightarrow B$ is injective if and only if it is surjective if and only if it is bijective.

□

Problem 2. Let \sim be an equivalence relation of the set A . For any $a, b \in A$,

a) $a \sim b$ if and only if $\bar{a} = \bar{b}$.

b) if $\bar{a} \neq \bar{b}$, then $\bar{a} \cap \bar{b} = \emptyset$

Proof.

a) Suppose that $a \sim b$. Without loss of generality let $c \in \bar{a}$. Then $c \sim a$ which implies that $c \sim b$ by transitivity and as such $c \in \bar{b}$. Therefore if $a \sim b$ then $\bar{a} = \bar{b}$.

Now suppose that $\bar{a} = \bar{b}$. Since $a \in \bar{a}$ and $b \in \bar{b}$ by reflexivity we know that $a, b \in \bar{b}$ and as such $a \sim b$.

Therefore $a \sim b$ if and only if $\bar{a} = \bar{b}$.

b) Suppose that $\bar{a} \neq \bar{b}$ and that there existed a $c \in \bar{a} \cap \bar{b}$. Then $a \sim c$ and $c \sim b$ which would imply that $a \sim b$ by transitivity and that $\bar{a} = \bar{b}$ by the previous part of the proposition which is a contradiction.

Therefore if $\bar{a} \neq \bar{b}$ then $\bar{a} \cap \bar{b} = \emptyset$

□

Problem 3. Let n be a fixed positive integer. Then

$$(\mathbb{Z}/n\mathbb{Z})^\times = \{\bar{a} \in \mathbb{Z}/n\mathbb{Z} \mid 1 \leq a \leq n \text{ and } a, n \text{ are relatively prime}\}$$

Proof. Let $\bar{a} \in (\mathbb{Z}/n\mathbb{Z})^\times$. Then \bar{a} has a multiplicative inverse \bar{c} such that $\bar{a}\bar{c} = \bar{1}$.

□