

Problem 1 (7.14). 1. Let R be a commutative ring with $1 \neq 0$ and I_1, \dots, I_n pairwise comaximal ideals of R . Prove that

$$(R/(I_1 \dots I_n))^X \cong (R/I_1)^X \times \dots \times (R/I_n)^X$$

as groups.

2. Let m, n be relatively prime positive integers. Prove that

$$(\mathbb{Z}_{mn})^X \cong (\mathbb{Z}_m)^X \times (\mathbb{Z}_n)^X$$

as groups.

3. Solve the system of congruences:

$$\begin{aligned} x &\equiv 2 \pmod{9} \\ x &\equiv 3 \pmod{5} \\ x &\equiv 1 \pmod{7} \\ x &\equiv 5 \pmod{11} \end{aligned}$$

Proof. 1. By the Chinese Remainder Theorem we know that

$$(R/(I_1 \dots I_n)) \cong (R/I_1) \times \dots \times (R/I_n)$$

as rings. As such there is an isomorphism $\varphi : R \rightarrow S$ and since ring isomorphisms send units to units we have a bijection $\varphi|_{R^X} : R^X \rightarrow S^X$. However since φ is an isomorphism it will send identity to identity and preserve multiplication.

Thus $\varphi|_{R^X}$ is a group homomorphism and

$$(R/(I_1 \dots I_n))^X \cong (R/I_1)^X \times \dots \times (R/I_n)^X$$

are isomorphic as groups.

2. From a prior homework we have that $(n)(m) = (nm)$. Therefore $\mathbb{Z}_{mn} \cong \mathbb{Z}/m\mathbb{Z}n\mathbb{Z}$ in addition to $\mathbb{Z}_n \cong \mathbb{Z}/n\mathbb{Z}$ and $\mathbb{Z}_m \cong \mathbb{Z}/m\mathbb{Z}$. Finally note that since $\gcd(n, m) = 1$ there exist $\alpha, \beta \in \mathbb{Z}$ such that $\alpha a + \beta b = 1$ and therefore $(n) + (m) = \mathbb{Z}$. Thus we can apply part 1 of this problem to get that

$$\mathbb{Z}_{mn}^X \cong \mathbb{Z}_m^X \times \mathbb{Z}_n^X$$

3. $x \cong 533 \pmod{3465}$

□

Problem 2 (8.1). Prove that the division algorithm holds for any polynomial ring over a field.

Proof. Let $f, g \in k[x]$ where k is a field. We will prove the division algorithm holds via induction over the degree of f .

Let $\deg(f) = 0$. Then $f = 0 \cdot g + f$ and since $\deg(f) = 0 < \deg(g)$ this is a valid choice for the division algorithm.

Assume that the division algorithm holds for polynomials $f \in k[x]$ when $\deg(f) = n$. Then given $g \in k[x]$ we have $f = qg + r$ where $\deg(g) > \deg(r)$. However we can form any given a polynomial $f' = \sum_0^{n+1} a_i x^i$ let $f := \sum_0^n a_{i+1} x^i + a_0$. Then $f' = x \cdot f + a_0$. Apply the division algorithm to f and we get

$$f' = x(qg + r) = (q \cdot x)g + x \cdot r$$

which shows that the division algorithm holds for $\deg = k + 1$ if we assume it for $\deg = k$.

Therefore the division algorithm holds for any polynomial ring over a field.

□

Problem 3 (8.2). 1. Prove that $a|b$ iff $b \in (a)$ iff $(b) \subseteq (a)$.

2. If $a|b$ and $a|c$, prove that $a|(bx + cy)$ for all $x, y \in R$.

3. Suppose $b \neq 0$. If $a|b$ and $b|c$, then $a|c$.

4. If d is a greatest common divisor of a, b then du is also a greatest common divisor of a, b for any unit u of R .

Proof. 1. Suppose that $a|b$. Then there exists a c such that $ac = b$ which implies that $b \in (a)$.

Next suppose that $b \in (a)$ and let $d \in (b)$. Then $d = fb$. However $b = ca$ since $b \in (a)$ and as such $d = fca \in (a)$.

Finally suppose that $(b) \subseteq (a)$. Then $b \in (a)$ which implies that $b = ca$ for some c . This is the definition of $a|b$.

Therefore $a|b$ iff $b \in (a)$ iff $(b) \subseteq (a)$.

2. Suppose that $a|b$ and that $a|c$. Then there exist $\beta, \gamma \in R$ such that $a\beta = b$ and $a\gamma = c$. If we have $(bx + cy)$ we can substitute b, c to get

$$bx + cy = a\beta x + a\gamma c = a(\beta x + \gamma y)$$

which implies that $a|(bx + cy)$.

Therefore if $a|b$ and $a|c$ then $a|(bx + cy)$ for all $x, y \in R$.

3. Suppose that $a|b$ and $b|c$. Then there exist $\alpha, \beta \in R$ such that $\alpha a = b$ and $\beta b = c$. Thus $\beta\alpha a = c$ and therefore $a|c$.

4. Let d be a greatest common divisor of a, b . Then $d|a, d|b$ and if $d'|a, b$ then $d|d'$. Consider ud where u is a unit. Then ud divides a, b as $\alpha d = a$ and $\beta d = b$ which implies that $\alpha u^{-1}(ud) = a$ and $\beta u^{-1}(ud) = b$. Thus $ud|a$ and $ud|b$. Now suppose that $f|a$ and $f|b$. Then $d|f$ which implies that $\gamma d = f$. It then follows that $\gamma u^{-1}(ud) = f$ and thus $ud|f$.

Therefore ud is a greatest common divisor of a and b . □

Problem 4 (8.3). An element p in an integral domain R is prime if, and only if, $p|ab$ implies $p|a$ or $p|b$ for any $a, b \in R$.

Proof. Let $p \in R$ be prime. Suppose that $p|ab$. Then $\alpha p = ab$ which implies that $ab \in (p)$. However since p is prime, (p) is a prime ideal. This means that either $a \in (p)$, in which case $p|a$, or $b \in (p)$ with $p|b$.

Otherwise suppose that when $p|ab$ then $p|a$ or $p|b$. Let $ab \in (p)$. Then $p|ab$ which implies that $p|a$, in which case $a \in (p)$, or $p|b$ and $b \in (p)$ which shows that (p) is prime.

Therefore an element p in an integral domain R is prime if, and only if, $p|ab$ implies that $p|a$ or $p|b$ for any $a, b \in R$. □

Problem 5 (8.4). Let R be a UFD and $a, b \in R \setminus \{0\}$. Then a, b has a greatest common divisor in R . If a, b are relatively prime and $a|bc$ for some $c \in R$, then $a|c$.

Proof. □

Problem 6 (G1). Let H be a normal subgroup of a group G , and let K be a subgroup of H .

1. Give an example of this situation where K is not a normal subgroup of G .
2. Prove that if the normal subgroup H is cyclic, then K is normal in G .

Proof. 1. Consider S_5 and A_5 . We know that $A_5 \trianglelefteq S_5$ however the subgroup $\langle (1\ 2\ 3) \rangle$ is not normal in S_5 .

2.

□

Problem 7 (G2). Prove that every finite group of order at least three has a nontrivial automorphism.

Proof.

□

Problem 8 (R1). Let $R = \mathbb{Z}[\sqrt{-3}] = \{a + b\sqrt{-3} \mid a, b \in \mathbb{Z}\}$.

1. Why is R an integral domain?
2. What are the units in R ?
3. Is the element 2 irreducible in R ?
4. If $x, y \in R$, and $2 \mid xy$, does it follow that 2 divides either x or y ? Justify your answer.

Proof.

□

Problem 9 (R2). 1. Give an example of an integral domain with exactly 9 elements.

2. Is there an integral domain with exactly 10 elements? Justify your answer.

Proof. 1. $\mathbb{Z}_3[\sqrt{2}]$

2.

□

Problem 10 (R3). Let

$$F = \left\{ \begin{pmatrix} a & b \\ 2b & a \end{pmatrix} \mid a, b \in \mathbb{Q} \right\}$$

1. Prove that F is a field under the usual matrix operations of addition and multiplication.
2. Prove that F is isomorphic to the field $\mathbb{Q}(\sqrt{2})$.

Proof.

□