

Problem 1 (2.7.1).

Proof. First we will show that N is normal. Then given $g \in N$ for all $x \in G$ there exists an $h \in H$ such that $g = xhx^{-1}$. Let $k \in G$ then $kgk^{-1} = h \in H$ as $g \in k^{-1}Hk$. However $g \in k^{-1}xHx^{-1}k$ for any $x \in G$. It then follows that

$$h = kgk^{-1} = kk^{-1}xh'x^{-1}kk^{-1} = xh'x$$

which implies that $h \in xHx^{-1}$ for all $x \in G$ and therefore $h \in N$. Therefore N is a normal subgroup of G .

Now we will show that N is the largest normal subgroup of G contained in H . Let M be a subgroup of G such that $M \trianglelefteq G$ and $M \leq H$. Then given $g \in M$ and $x \in G$ we have $x^{-1}gx = h \in M$. However this implies that $g = xhx^{-1}$ for all x and since $M \leq H$ we have that $h \in H$ and therefore $g \in N$ and $M \leq N$.

Therefore since N is normal and normal subgroup contained in H is contained in N we have that N is the largest normal subgroup contained in H . \square

Problem 2 (2.7.2).

Proof. a) For reflexivity, since $H, K \leq G$ we have $exe = x \in HxK$.

For symmetry if $x \sim y$ then $x \in HyK$ which implies that $x = hyk$ for some $h \in H$ and $k \in K$. However since $H, K \leq G$ we have that $h^{-1}xk^{-1} = y$ which implies that $y \in HxK$ and therefore $y \sim x$.

For transitivity suppose that we have $x \sim y$ and $y \sim z$. Then as before we have $x = hyk$ and $y = h'zk'$. It then follows that $x = hh'zk'k$ which implies that $x \in HzK$ and therefore $x \sim z$.

Let \bar{x} denote the equivalence class of x . Then if $y \in HxK$ by definition $y \sim x$ and $y \in X$. Otherwise if $y \in \bar{x}$ by definition $y \sim x$ which implies that $y \in HxK$.

Therefore \sim is an equivalence relation, the equivalence classes are of the form HxK , and as such $H \setminus G/K$ forms a partition of G .

b)

c)

\square

Problem 3 (2.8).

Proof. 1. We'll start by showing that $C_G(A)$ is a subgroup. If we have $g, h \in C_G(A)$ then $gha = gah = agh$ so it is closed under the group operation. Then if $g \in C_G(A)$ we have $ga = ag$. Multiply on the left by and right by g^{-1} and we get $ag^{-1} = g^{-1}a$. Therefore $C_G(A)$ is a subgroup.

Next consider $N_G(A)$. If we have $g, h \in N_G(A)$ then $hah^{-1} = a' \in A$. This implies that $ga'g^{-1} \in A$ and therefore $ghah^{-1}g^{-1} \in A$. Next let $g \in N_G(A)$ and $a \in A$. Then **Closed under inverses**.

Let $g \in C_G(A)$ then for $a \in A$ we have $gag^{-1} = agg^{-1} = a$ which implies that $g \in N_G(A)$. Therefore $C_G(A) \subset N_G(A)$.

2. Let $a \in A$ and $n \in N_G(A)$. Then $nan^{-1} \in A$ by definition of $N_G(A)$. Therefore if $A \leq G$ then $A \trianglelefteq N_G(A)$.
3. Let $z \in Z(G)$ and let $g \in G$. Then $zgz^{-1} = gzz^{-1} = g$ which implies that $Z(G)$ is a normal subgroup of G .
4. For the group S_3 take the subgroup $\langle (1\ 2) \rangle$. Then $(1\ 2\ 3)(1\ 2)(3\ 2\ 1) = (2\ 3) \notin \langle (1\ 2) \rangle$ which implies that $\langle (1\ 2) \rangle$ is not normal.
5. If n is even then the center consists of $r^{n/2}$ and the identity. Otherwise if n is odd then the center is just the identity.
6. The subgroups of \mathcal{Q}_8 are $\langle 1 \rangle, \langle -1 \rangle, \langle i \rangle, \langle j \rangle, \langle k \rangle, \mathcal{Q}_8$. The trivial group and \mathcal{Q}_8 are both normal. For the others we have

$\langle -1 \rangle$

g	$g(-1)g$
1	-1
-1	
i	
$-i$	
j	
$-j$	
k	
$-k$	

$\langle i \rangle$

g	$g(i)g$
1	i
-1	
i	
$-i$	
j	
$-j$	
k	
$-k$	

$\langle j \rangle$

g	$g(j)g$
1	j
-1	
i	
$-i$	
j	
$-j$	
k	
$-k$	

$\langle k \rangle$

g	$g(k)g$
1	k
-1	
i	
$-i$	
j	
$-j$	
k	
$-k$	

□

Problem 4 (2.9.1).

Proof. Since $|G| = p$ where p is prime by Lagrange's Theorem (2.12 in notes) the order of any subgroup must be either 1 or p . However the only subgroups that fulfill these criterion are either the trivial group or G itself. Therefore G cannot have any non-trivial subgroups.

Let $g \in G \setminus \{e\}$. Since there is no non-trivial subgroup the element $\langle g \rangle$ must generate the whole group. Then by definition G is cyclic. □

Problem 5 (2.9.2).

Proof. □

Problem 6 (2.10).

Proof. □