

**Problem 1** (7.26). Let  $M$  and  $N$  be finitely generated  $R$ -modules over a PID  $R$ . Compute  $M \otimes_R N$ . As a special case, if  $M$  is a finite abelian group with invariant factors  $s_1, \dots, s_t$  (where as usual we assume that  $s_i$  divides  $s_{i+1}$ ), show that  $M \otimes_{\mathbb{Z}} M$  is a finite group of order  $\prod_{j=1}^t s_j^{2t-2j+1}$ .

*Proof.* By the structure theorem for finitely generated modules over a PID we have that

$$M \cong R^r \oplus R/(a_1) \oplus \cdots \oplus R/(a_m)$$

and

$$N \cong R^s \oplus R/(b_1) \oplus \cdots \oplus R/(b_n)$$

where  $a_i | a_{i+1}$  and  $b_i | b_{i+1}$ . Then by Theorem 2.18 of Adkins' book the tensor  $M \otimes_R N$  distributes over the direct sum. Together with  $R/(a_i) \otimes_R R/(b_j) \cong R/(a_i + b_j)$  we get that

$$M \otimes_R N \cong \bigoplus_{0 < i \leq m, 0 < j \leq n} (R/(a_i + b_j)) \bigoplus_{0 < i \leq m} (R/(a_i) \otimes_R R^t) \bigoplus_{0 < j \leq n} (R^s \otimes_R R/(b_j))$$

If we let  $M$  be a finite abelian group with invariant factor decomposition

$$M \cong \bigoplus_{0 < i \leq t} \mathbb{Z}_{s_i}$$

Then, from above, the tensor of  $M$  with itself is

$$M \otimes_{\mathbb{Z}} M \cong \bigoplus_{0 < i, j \leq t} \mathbb{Z}_{\gcd(s_i, s_j)}$$

However since  $s_i | s_j$  for  $i \leq j$  we have that  $\gcd(s_i, s_j) = s_i$ . This will cause  $2t - 1$  copies of  $\mathbb{Z}_{s_1}$  to appear,  $2t - 3$  for  $\mathbb{Z}_{s_2}$ , and so on until we have only a single copy of  $\mathbb{Z}_{s_t}$ . Thus  $M \otimes_{\mathbb{Z}} M$  will have order  $\prod_{j=1}^t s_j^{2t-2j+1}$ .  $\square$

**Problem 2** (7.30). (a) Let  $F$  be a field and  $K$  a field containing  $F$ . If  $f(X) \in F[X]$ , show that there is an isomorphism of  $K$ -algebras:

$$K \otimes_F (F[X]/\langle f(X) \rangle) \cong K[X]/\langle f(X) \rangle$$

(b) By choosing  $F, f(x)$ , and  $K$  appropriately, find an example of two fields  $K$  and  $L$  containing  $F$  such that the  $F$ -algebra  $K \otimes_F L$  has nilpotent elements.

*Proof.* (a) Define a map  $\varphi : K \times F[x]/\langle g(x) \rangle \rightarrow K[x]/\langle g(x) \rangle$  via

$$\varphi(k, f) = kf$$

Now briefly verify that  $\varphi$  is  $F$ -middle linear

$$\begin{aligned} \varphi(ak, fb) &= akfb &= a\varphi(k, f)b \\ \varphi(k_1 + k_2, f) &= k_1f + k_2f &= \varphi(k_1, f) + \varphi(k_2, f) \\ \varphi(k, f_1 + f_2) &= kf_1 + kf_2 &= \varphi(k, f_1) + \varphi(k, f_2) \\ \varphi(ka, f) &= kaf &= \varphi(k, af) \end{aligned}$$

Since  $\varphi$  is  $F$ -middle linear it induces a map  $\tilde{\varphi} : K \otimes_F F[x]/\langle g(x) \rangle \rightarrow K[x]/\langle g(x) \rangle$ . Now define a map  $\psi : K[x]/\langle g(x) \rangle \rightarrow K \otimes_F F[x]/\langle g(x) \rangle$  by  $\psi(kx^i) = k \otimes x^i$  and extending linearly. We will now show that  $\tilde{\varphi}$  and  $\psi$  are inverses.

First  $\tilde{\varphi} \circ \psi$

$$\begin{aligned}\tilde{\varphi} \circ \psi(\sum a_i x^i) &= \tilde{\varphi}(\sum a_i \otimes x^i) \\ &= \sum a_i x^i\end{aligned}$$

Then for  $\psi \circ \tilde{\varphi}$

$$\begin{aligned}\psi \circ \tilde{\varphi}(k \otimes \sum a_i x^i) &= \psi \circ \tilde{\varphi}(\sum k \otimes a_i x^i) \\ &= \psi(\sum \tilde{\varphi}(k \otimes a_i x^i)) \\ &= \psi(\sum k a_i x^i) \\ &= \sum \psi(k a_i x^i) \\ &= \sum k a_i \otimes x^i \\ &= \sum k \otimes a_i x^i\end{aligned}$$

Since  $\tilde{\varphi}$  has an inverse it is indeed an isomorphism. Which shows that  $K \otimes_F F[x]/\langle g(x) \rangle$  is isomorphic to  $K[x]/\langle g(x) \rangle$ .

- (b) Let  $F$  be rational functions of  $t$  with coefficients in  $\mathbb{Z}_2$ . Then let  $K$  be the splitting field of  $x^2 - t$ . Then in  $K \otimes_F K$  the element  $t^{1/2} \otimes 1 + 1 \otimes t^{1/2}$  is zero when squared making it an idempotent.

□

**Problem 3 (7.31).** Let  $F$  be a field. Show that  $F[X, Y] \cong F[X] \otimes_F F[Y]$  where the isomorphism is an isomorphism of  $F$ -algebras.

*Proof.* Define  $\varphi : F[x] \times F[y] \rightarrow F[x] \otimes F[y]$  via

$$\varphi(\sum_i a_i x^i, \sum_j b_j y^j) = \sum_{i,j} a_i b_j x^i y^j$$

Since we are in a polynomial ring over a field it follows that  $\varphi$  is  $F$ -middle linear. As such there is a map  $\tilde{\varphi} : F[x] \otimes F[y] \rightarrow F[x, y]$  that mimics  $\varphi$ .

Now define  $\psi : F[x, y] \rightarrow F[x] \otimes F[y]$  via  $\psi(cx^i y^j) = c(x^i \otimes y^j)$  and extending linearly. We will now show that  $\tilde{\varphi}$  and  $\psi$  are inverses. Since both of these maps are linear we only need to verify them on either  $cx^i y^j$  and  $a_i x^i \otimes b_j y^j$ .

First for  $\tilde{\varphi} \circ \psi$  we have

$$\begin{aligned}\tilde{\varphi} \circ \psi(cx^i y^j) &= \tilde{\varphi}(c(x^i \otimes y^j)) \\ &= \tilde{\varphi}(cx^i \otimes y^j) \\ &= cx^i y^j\end{aligned}$$

Then for  $\psi \circ \tilde{\varphi}$

$$\begin{aligned}\psi \circ \tilde{\varphi}(a_i x^i \otimes b_j y^j) &= \psi(a_i b_j x^i y^j) \\ &= a_i b_j (x^i \otimes y^j) \\ &= a_i x^i \otimes b_j y^j\end{aligned}$$

Thus  $\tilde{\varphi}$  is an isomorphism which shows that  $F[x, y]$  is isomorphic to  $F[x] \otimes F[y]$ .  $\square$

**Problem 4** (7.34). *Let  $F$  be a field,  $V$  and  $W$  finite-dimensional vector spaces over  $F$ , and let  $T \in \text{End}_F(V)$ ,  $S \in \text{End}_F(W)$ .*

(a) *If  $\alpha$  is an eigenvalue of  $S$  and  $\beta$  is an eigenvalue of  $T$ , show that the product  $\alpha\beta$  is an eigenvalue of  $S \otimes T$ .*

(b) *If  $S$  and  $T$  are diagonalizable, show that  $S \otimes T$  is diagonalizable.*

*Proof.* (a) Let  $u, v$  be eigenvectors with eigenvalues  $\alpha, \beta$  for linear transformations  $S$  and  $T$  respectively. Then

$$S \otimes T(u \otimes v) = S(u) \otimes T(v) = \alpha u \otimes \beta v = \alpha\beta(u \otimes v)$$

(b) Since  $S$  and  $T$  are diagonalizable they each have a number of eigenvalues equal to the dimension of the space they act on. By the previous problem the product of two eigenvalues gives an eigenvalue which means there will be  $\dim(V)\dim(W) = \dim(V \otimes W)$  eigenvalues for  $S \otimes T$  making it diagonalizable.  $\square$

**Problem 5** (10.4.3). *Show that  $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$  and  $\mathbb{C} \otimes_{\mathbb{C}} \mathbb{C}$  are both left  $\mathbb{R}$  modules but are not isomorphic as  $\mathbb{R}$ -modules.*

*Proof.* The complex numbers can be given the bimodule structures  $(\mathbb{R}, \mathbb{R})$ ,  $(\mathbb{R}, \mathbb{C})$ , and  $(\mathbb{C}, \mathbb{R})$  by left and right multiplication from the correct field. As such both of the above tensors can be given a left  $\mathbb{R}$ -module structure.

To show they are not isomorphic first note that from Dummit 10.4.19 we know that  $\mathbb{C} \otimes_{\mathbb{C}} \mathbb{C} \cong \mathbb{C}$  which has rank 2 as a free  $\mathbb{R}$  module. Meanwhile  $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$  has free rank 4 as a left  $\mathbb{R}$ -module (Dummit pg. 375) which implies that they cannot be isomorphic.  $\square$

**Problem 6** (10.4.5). *Let  $A$  be a finite abelian group of order  $n$  and let  $p^k$  be the largest power of a prime  $p$  dividing  $n$ . Prove that  $\mathbb{Z}/p^k\mathbb{Z} \otimes_{\mathbb{Z}} A$  is isomorphic to the Sylow  $p$ -subgroup of  $A$ .*

*Proof.* Let  $\text{Syl}_p(A)$  denote the Sylow  $p$ -subgroup of  $A$ . Then by theorem 2.18 from Adkins' we have that

$$\mathbb{Z}_{p^k} \otimes_{\mathbb{Z}} A \cong \bigoplus_{q \text{ prime}} \mathbb{Z}_{p^k} \otimes_{\mathbb{Z}} \text{Syl}_q(A)$$

Now we will show that  $\mathbb{Z}_{p^k} \otimes_{\mathbb{Z}} \text{Syl}_q(A)$  is zero when  $q \neq p$  and is isomorphic to  $\text{Syl}_p(A)$  otherwise.

Let's start when  $p \neq q$ . Then let  $q^l$  be the highest power of  $q$  that divides  $n$ . As  $p$  and  $q$  are relatively prime there exist  $\alpha$  and  $\beta$  such that  $\alpha p^k + \beta q^l = 1$ . Then given  $x \otimes a \in \mathbb{Z}_{p^k} \otimes_{\mathbb{Z}} \text{Syl}_q(A)$  we have

$$x \otimes a = x(\alpha p^k + \beta q^l) \otimes a = (x\alpha p^k + x\beta q^l) \otimes a = x \otimes q^l a = x \otimes 0 = x \otimes p^k 0 = xp^k \otimes 0 = 0 \otimes 0$$

which shows that every element of  $\mathbb{Z}_{p^k} \otimes_{\mathbb{Z}} \text{Syl}_q(A)$  is trivial making the group itself trivial.

The fact that  $\mathbb{Z}_{p^k} \otimes_{\mathbb{Z}} \text{Syl}_p(A) \cong \text{Syl}_p(A)$  follows from both constituents have size  $p^k$ . As such the inner action of  $\mathbb{Z}$  is equivalent to the corresponding action by  $\mathbb{Z}_{p^k}$ . Thus  $\mathbb{Z}_{p^k} \otimes_{\mathbb{Z}} \text{Syl}_p(A)$  is isomorphic to  $\mathbb{Z}_{p^k} \otimes_{\mathbb{Z}_{p^k}} \text{Syl}_p(A)$  which is then isomorphic to  $\text{Syl}_p(A)$ .

Therefore  $\mathbb{Z}_{p^k} \otimes_{\mathbb{Z}} A$  is isomorphic to the Sylow  $p$ -subgroup of  $A$ .  $\square$

**Problem 7** (10.4.10). Suppose  $R$  is commutative and  $N \cong R^n$  is a free  $R$ -module of rank  $n$  with  $R$ -module basis  $e_1, \dots, e_n$ .

- (a) For any nonzero  $R$ -module  $M$  show that every element of  $M \otimes N$  can be written uniquely in the form  $\sum_{i=1}^n m_i \otimes e_i$  where  $m_i \in M$ . Deduce that if  $\sum_{i=1}^n m_i \otimes e_i = 0$  in  $M \otimes N$  then  $m_i = 0$  for  $i = 1, \dots, n$ .
- (b) Show that if  $\sum m_i \otimes n_i = 0$  in  $M \otimes N$  where the  $n_i$  are merely assumed to be  $R$ -linearly independent then it is not necessarily true that all the  $m_i$  are 0. [Consider  $R = \mathbb{Z}, n = 1, M = \mathbb{Z}/2\mathbb{Z}$ , and the element  $1 \otimes 2$ .]

*Proof.* (a) Let  $m \otimes n \in M \otimes_R N$ . Then since  $N$  is free we can rewrite  $m \otimes n$  as

$$m \otimes n = m \otimes \left( \sum_{i=1}^n a_i e_i \right) = \sum_i m \otimes a_i e_i = \sum_{i=1}^n a_i m \otimes e_i$$

Since this process is entirely reversible if another element had the same decomposition then it would be equal to the original value. As such we know that  $\sum_{i=1}^n 0 \otimes e_i = 0$  and by uniqueness it follows that this is the only way of writing zero.

- (b) Let  $R = \mathbb{Z}, n = 1, M = \mathbb{Z}_2$ , and  $N = \mathbb{Z}$ . Then  $1 \otimes 2 = 1 \cdot 2 \otimes 1 = 0 \otimes 1 = 0$  which fulfills the conditions laid out above.  $\square$

**Problem 8** (10.4.24). Prove that the extension of scalars from  $\mathbb{Z}$  to the Gaussian integers  $\mathbb{Z}[i]$  of the ring  $\mathbb{R}$  is isomorphic to  $\mathbb{C}$  as a ring:  $\mathbb{Z}[i] \otimes_{\mathbb{Z}} \mathbb{R} \cong \mathbb{C}$  as rings.

*Proof.* Let  $\varphi : \mathbb{Z}[i] \times \mathbb{R} \rightarrow \mathbb{C}$  be defined via  $\varphi(k, x) = kx$ . Since it is effectively the same map that we defined before we can see that it is  $\mathbb{Z}$ -middle linear. As such there is an induced map  $\tilde{\varphi} : \mathbb{Z}[i] \otimes_{\mathbb{Z}} \mathbb{R} \rightarrow \mathbb{C}$  where  $\tilde{\varphi}(k \otimes x) = kx$ . Define  $\psi : \mathbb{C} \rightarrow \mathbb{Z}[i] \otimes_{\mathbb{Z}} \mathbb{R}$  via  $\psi(x) = 1 \otimes x$  and  $\psi(ix) = i \otimes x$  when  $x \in \mathbb{R}$  then extending linearly. We will now show that  $\tilde{\varphi}$  and  $\psi$  are inverses.

First for  $\psi \circ \tilde{\varphi}$  we have

$$\begin{aligned} \psi \circ \tilde{\varphi}((a + bi) \otimes x) &= \psi(ax + bxi) \\ &= \psi(ax) + \psi(bxi) \\ &= 1 \otimes ax + i \otimes bx \\ &= a \otimes x + bi \otimes x \\ &= (a + bi) \otimes x \end{aligned}$$

Then for  $\tilde{\varphi} \circ \psi$

$$\begin{aligned}\tilde{\varphi} \circ \psi(u + iv) &= \tilde{\varphi}(1 \otimes u + i \otimes v) \\ &= \tilde{\varphi}(1 \otimes u) + \tilde{\varphi}(i \otimes v) \\ &= u + iv\end{aligned}$$

Thus  $\psi$  and  $\tilde{\varphi}$  are inverses which makes  $\tilde{\varphi}$  an isomorphism which shows that  $\mathbb{C}$  is isomorphic to  $\mathbb{Z}[i] \otimes \mathbb{R}$ .  $\square$

**Problem 9** (10.4.27). (a) Write down a formula for the multiplication of two elements  $a \cdot 1 + b \cdot e_2 + c \cdot e_3 + d \cdot e_4$  and  $a' \cdot 1 + b' \cdot e_2 + c' \cdot e_3 + d' \cdot e_4$  in the example  $A = \mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$  following proposition 21 (where  $1 = 1 \otimes 1$  is the identity of  $A$ ).

(b) Let  $\epsilon_1 = \frac{1}{2}(1 \otimes 1 + i \otimes i)$  and  $\epsilon_2 = \frac{1}{2}(1 \otimes 1 - i \otimes i)$ . Show that  $\epsilon_1 \epsilon_2 = 0$ ,  $\epsilon_1 + \epsilon_2 = 1$ , and  $\epsilon_j^2 = \epsilon_j$  for  $j = 1, 2$  ( $\epsilon_1$  and  $\epsilon_2$  are called orthogonal idempotents in  $A$ ). Deduce that  $A$  is isomorphic as a ring to the direct product of two principle ideals:  $A \cong A\epsilon_1 \times A\epsilon_2$  (cf. Exercise 1, Section 7.6).

(c) Prove that the map  $\varphi : \mathbb{C} \otimes \mathbb{C} \rightarrow \mathbb{C} \times \mathbb{C}$  by  $\varphi(z_1, z_2) = (z_1 z_2, z_1 \bar{z}_2)$ , where  $\bar{z}_2$  denotes the complex conjugate of  $z_2$ , is an  $\mathbb{R}$ -bilinear map.

(d) Let  $\Phi$  be the  $\mathbb{R}$ -module homomorphism from  $A$  to  $\mathbb{C} \times \mathbb{C}$  obtained from  $\varphi$  in (c). Show that  $\Phi(\epsilon_1) = (0, 1)$  and  $\Phi(\epsilon_2) = (1, 0)$ . Show also that  $\Phi$  is  $\mathbb{C}$ -linear, where the action of  $\mathbb{C}$  is on the left tensor factor in  $A$  and on both factors in  $\mathbb{C} \times \mathbb{C}$ . Deduce that  $\Phi$  is surjective. Show that  $\Phi$  is a  $\mathbb{C}$ -algebra isomorphism.

*Proof.* (a) Let  $e_1 = 1 \otimes 1, e_2 = 1 \otimes i, e_3 = i \otimes 1, e_4 = i \otimes i$ . Then multiply out to get

$$\begin{aligned}aa'e_1^2 + a'be_1e_2 + ab'e_1e_2 + bb'e_2^2 + a'ce_1e_3 + ac'e_1e_3 + b'ce_2e_3 + bc'e_2e_3 + cc'e_3^2 + a'de_1e_4 + ad'e_1e_4 \\ + b'de_2e_4 + bd'e_2e_4 + c'de_3e_4 + cd'e_3e_4 + dd'e_4^2\end{aligned}$$

Evaluating the multiplication gets us

$$\begin{aligned}aa'e_1 + a'be_2 + ab'e_2 + bb'(-e_1) + a'ce_3 + ac'e_3 + b'ce_4 + bc'e_4 + cc'(-e_1) + a'de_4 + ad'e_4 + b'd(-e_3) + bd'(-e_3) \\ c'd(-e_2) + cd'(-e_2) + dd'e_1\end{aligned}$$

Which we can then simplify to

$$(aa' - bb' - cc' + dd')e_1 + (a'b + ab' - c'd - cd')e_2 + (a'c + ac' - b'd - bd')e_3 + (b'c + bc' + a'd + ad')e_4$$

(b) First adding

$$\epsilon_1 + \epsilon_2 = \frac{1}{2}(1 \otimes 1 + i \otimes i) + \frac{1}{2}(1 \otimes 1 - i \otimes i) = \frac{2}{2}(1 \otimes 1) = 1 \otimes 1 = 1$$

Then multiplying

$$\epsilon_1 \epsilon_2 = \frac{1}{2}(1 \otimes 1 + i \otimes i) \frac{1}{2}(1 \otimes 1 - i \otimes i) = \frac{1}{4}(1 \otimes 1 + (-1 \otimes -1)) = \frac{1}{4}(1 \otimes 1 - 1 \otimes 1) = 0$$

Squaring  $\epsilon_1$  gets you

$$\epsilon_1^2 = \frac{1}{4}(1 \otimes 1 + 1 \otimes 1 + i \otimes i + i \otimes i) = \frac{1}{2}(1 \otimes 1 + i \otimes i) = \epsilon_1$$

Similarly for  $\epsilon_2$

$$\epsilon_2^2 = \frac{1}{4}(1 \otimes 1 + 1 \otimes 1 - i \otimes i - i \otimes i) = \frac{1}{2}(1 \otimes 1 - i \otimes i) = \epsilon_2$$

Then from exercise 7.6.1 from Dummit we have that  $A \cong A\epsilon_1 \times A\epsilon_2$ .

- (c) Since the space  $\mathbb{C} \times \mathbb{C}$  can be treated as a vector space over  $\mathbb{R}$ . This along with the fact that we can pull real constants out of the complex conjugate implies that  $\varphi$  is  $\mathbb{R}$ -bilinear.
- (d) First we calculate  $\Phi(\epsilon_1)$  directly

$$\Phi(\epsilon_1) = \Phi\left(\frac{1}{2}(1 \otimes 1 + i \otimes i)\right) = \frac{1}{2}(\Phi(1 \otimes 1) + \Phi(i \otimes i)) = \frac{1}{2}((1, 1) + (-1, 1)) = (0, 1)$$

Similarly for  $\Phi(\epsilon_2)$  we have

$$\Phi(\epsilon_1) = \Phi\left(\frac{1}{2}(1 \otimes 1 - i \otimes i)\right) = \frac{1}{2}(\Phi(1 \otimes 1) - \Phi(i \otimes i)) = \frac{1}{2}((1, 1) - (-1, 1)) = (1, 0)$$

Next we show that  $\Phi$  is  $\mathbb{C}$ -linear.

$$\Phi(w(z_1 \otimes z_2)) = \Phi(wz_1 \otimes z_2) = (wz_1 z_2, wz_1 \bar{z}_2) = w(z_1 z_2, z_1 \bar{z}_2) = w\Phi(z_1 \otimes z_2)$$

Since  $\Phi$  is  $\mathbb{C}$ -linear and we have elements that map to  $(1, 0)$  and  $(0, 1)$  it must be surjective as this allows us to reach any element using linearity.

To see that  $\Phi$  is injective note that if  $\Phi(z_1 \otimes z_2) = 0$  then either  $z_1$  or  $z_2$  must be zero since  $\mathbb{C}$  is a field. Then

$$z_1 \otimes 0 = 0(z_1 \otimes 0) = 0 \otimes 0 = 0$$

The case for  $z_1 = 0$  follows similarly. As such the kernel of  $\Phi$  is trivial which implies that  $\Phi$  is in fact an isomorphism.

□