

Problem 1. Show that an angle of 30° and an angle of 15° cannot be trisected.

Proof. First note that if 15° can be trisected then so can 30° as we could bisect 30° , trisect 15° and then double the resulting angle. As such it will suffice to show that we cannot trisect 15° .

A number is constructible if, and only if, both its real and imaginary parts are constructible. If 15° were constructible then so would $e^{i \cdot 10^\circ}$ as it would be the intersection of the angle and the unit circle. The real part of which is

$$\alpha = \cos 10^\circ = \frac{1}{2} \sqrt{\frac{1}{2} \left(4 + 2 \cdot \left(\frac{1}{2} (1 + i\sqrt{3})^{-\frac{1}{3}} + 2^{\frac{2}{3}} (1 + i\sqrt{3}) \right) \right)}$$

We know that a number is constructible if, and only if, we have an ascending chain of fields $\mathbb{Q} = F_0 \subset \dots \subset F_n = \mathbb{Q}[\alpha]$ where all of the intermediate degrees are two. This enforces that the degree of the extension must be a power of 2. However for α at some point we will have to adjoin $(1 + i\sqrt{3})^{-\frac{1}{3}}$ for which the extension will be of degree 3. By the tower theorem this means that $3|\mathbb{Q}[\alpha]$ but this cannot occur.

Therefore α is not constructible and it then follows that neither 15° nor 30° can be trisected. \square

Problem 2. Let $\xi = e^{2\pi i/6} = \cos(2\pi/6) + i \sin(2\pi/6)$ be a primitive 6th root of unity over \mathbb{Q} . Find each of the following:

1. The minimum polynomial $f(x) \in \mathbb{Q}[x]$ of ξ over \mathbb{Q} .
 2. The splitting field F of $f(x)$ over \mathbb{Q} .
 3. $[F : \mathbb{Q}]$.
- (a) Let $f(x) = x^2 - x + 1$. This polynomial has ξ as a root. Moreover it is irreducible by the rational roots theorem as ± 1 are not roots.
 - (b) The roots of f are ξ and $-e^{2\pi i/3} = -\xi^2$. Thus $\mathbb{Q}[\xi, -\xi^2] = \mathbb{Q}[\xi]$ is the splitting field for f .
 - (c) Since the degree of f is 2 it follows that $[\mathbb{Q}[\xi] : \mathbb{Q}] = 2$.

Problem 3. Find a splitting field extension $K : \mathbb{Q}$ for each of the following polynomials over \mathbb{Q} and in each case determine the degree $[K : \mathbb{Q}]$.

$$(a) x^4 + 1 \quad (b) x^4 + 4 \quad (c) (x^4 + 1)(x^4 + 4) \quad (d) (x^4 - 1)(x^4 + 4)$$

- (a) The roots of $f(x) = x^4 + 1$ are $r := e^{\pi i/4}, r^3, r^5$, and r^7 . Since each all of the other roots can be expressed as a power of r we have that the splitting field of f is $\mathbb{Q}[r, r^3, r^5, r^7] = \mathbb{Q}[r]$ the degree of which is 4 as f is irreducible and thus the minimal polynomial. The irreducibility can be checked by shifting to $f(x + 1)$ and apply Eisenstein's Criterion with $p = 2$.
- (b) The roots of $g(x) = x^4 + 4$ are the same roots as above but with each multiplied by $\sqrt{2}$. Let $s := \sqrt{2}e^{\pi i/4}$. Then the other roots are $s^3/2, s^5/4$, and $s^7/8$. Similar to before the splitting field is then $\mathbb{Q}[s]$ and since this polynomial is irreducible we have that $[\mathbb{Q}[s] : \mathbb{Q}] = 4$.
- (c) The roots of $p(x) = fg(x) = (x^4 + 1)(x^4 + 4)$ are the roots of both part a and b . Since the roots here are cyclic if we take rs we get $\sqrt{2}e^{\pi i/2}$. Keep multiplying that by r and we can hit every root. Thus the splitting field will be $\mathbb{Q}[r]$ and the minimal polynomial will be the one from part (a) giving us that $[\mathbb{Q}[r] : \mathbb{Q}] = 4$ for the degree of our splitting field.

- (d) The roots of $q(x) = (x^4 - 1)(x^4 + 4)$ are the roots of part *b* as well as ± 1 and $\pm i$. However $s^2/2 = i$ which means that we can express all of the roots in terms of s . Similar to part (c) our splitting field is the same as b , $\mathbb{Q}[s]$. As before the degree of this splitting field is 4.

Problem 4. Let $f(x) \in \mathbb{Q}[x]$ be the minimal polynomial of $\alpha = \sqrt{2 + \sqrt{2}}$.

1. Show that $f(x) = x^4 - 4x^2 + 2$. Thus, $[\mathbb{Q}(\alpha) : \mathbb{Q}] = 4$.
 2. Show that $\mathbb{Q}(\alpha)$ is the splitting field of $f(x)$ over \mathbb{Q} .
- (a) It will follow that $[\mathbb{Q}(\alpha) : \mathbb{Q}] = 4$ if $f(x) = x^4 - 4x^2 + 2$ is irreducible since $f(\alpha) = 0$. However f is irreducible by Eisenstein's criterion using 2.
- (b) The roots of f are $\pm\sqrt{2 \pm \sqrt{2}}$

Problem 5. Let $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$ be the field with p elements, where p is a prime number. Write down all monic cubic polynomials in $\mathbb{F}_2[x]$, factor them completely into irreducible factors and construct a splitting field for each of them. Which of these fields are isomorphic?

Proof.

□

Problem 6. Let $f(x) = x^3 + 2x + 2 \in \mathbb{F}_3[x]$.

1. Show that $f(x)$ is irreducible in $\mathbb{F}_3[x]$.
2. Let α be a root of $f(x)$ in some extension field K of \mathbb{F}_3 , so that $[\mathbb{F}_3[\alpha] : \mathbb{F}_3] = \deg f(x) = 3$. Show that $\mathbb{F}_3[\alpha]$ is a splitting field of $f(x)$ over \mathbb{F}_3 .

Proof.

□

Problem 7. Suppose that $f(x) \in F[x]$ is irreducible of degree $n > 0$, and let L be the splitting field of $f(x)$ over F .

1. Suppose that $[L : F] = n!$. Prove that $f(x)$ is irreducible.
2. Show that the converse of part (a) is false.

Proof.

□