

Problem 1.

Proof. Let X be a space such that all paths having the same endpoints are fixed-endpoint homotopic. Then given a loop α based at some point x_0 this loop is homotopic to the constant map at x_0 which implies that $\pi_1(X, x_0)$ is trivial for all X . Therefore X is simply connected.

Now suppose that X is a simply connected space. Let α, β be loops from x_0 to x_1 . For a fixed $t' \in [0, 1]$ using path connectedness we can construct a loop $l_{t'}$ such that $l_{t'}(0) = l_{t'}(1) = \alpha(t')$, $l_{t'}(\frac{1}{2}) = \beta(t)$ and that the function $p(t) = l_t(\frac{1}{2})$ is a path **Why?**. Since X is simply connected there is a homotopy $L_{t'}$ from $l_{t'}$ to the constant map at $\alpha(t')$. Then we can construct a homotopy H from β to α as

$$H(t, s) = L_t(\frac{1}{2}, s)$$

which will be continuous because of the continuity of the $L_{t'}$ s and that we choose $l_{t'}$ such that as we vary over t there is a path.

Therefore a space X is simply connected if and only if all paths with the same endpoints are fixed endpoint homotopic. \square

Problem 2.

Proof. Let $f : (X, x_0) \rightarrow (Y, y_0)$ and $g : (Y, y_0) \rightarrow (Z, z_0)$. For a continuous map h we have h_* defined as $h_*([\gamma]) = [h \circ \gamma]$. Then if we consider $(g \circ f)_*$:

$$(g \circ f)_*([\gamma]) = [(g \circ f) \circ \gamma] = [g \circ (f \circ \gamma)] = g_*([f \circ \gamma]) = g_* \circ f_*([\gamma])$$

$$\text{Therefore } (g \circ f)_* = g_* \circ f_*.$$

 \square **Problem 3.**

Proof.

 \square **Problem 4.**

Proof.

 \square **Problem 5.**

Proof.

 \square **Problem 6.**

Proof.

 \square