

Problem 1 (2.2.36). Show that $H_i(X \times S^n) \cong H_i(X) \oplus H_{i-n}(X)$ for all i and n , where $H_i = 0$ for $i < 0$ by definition. Namely, show $H_i(X \times S^n) \cong H_i(X) \oplus H_i(X \times S^n, X \times \{x_0\})$ and $H_i(X \times S^n, X \times \{x_0\}) \cong H_{i-1}(X \times S^{n-1}, X \times \{x_0\})$ [For the latter isomorphism the relative Mayer-Vietoris sequence yields an easy proof].

Proof. First we show that $H_i(X \times S^n) \cong H_i(X) \oplus H_i(X \times S^n, X \times \{x_0\})$. Swap out $H_i(X)$ for $H_i(X \times \{x_0\})$. Then apply the long exact sequence for the pair to get

$$\cdots \rightarrow H_{i+1}(X \times S^n, X \times \{x_0\}) \rightarrow H_i(X \times \{x_0\}) \rightarrow H_i(X \times S^n) \rightarrow H_i(X \times S^n, X \times \{x_0\}) \rightarrow H_{i-1}(X \times \{x_0\}) \rightarrow \cdots$$

Thus by the splitting lemma we have that $H_i(X \times S^n) \cong H_i(X) \oplus H_i(X \times S^n, X \times \{x_0\})$.

For the next isomorphism $H_i(X \times S^n, X \times \{x_0\}) \cong H_{i-1}(X \times S^{n-1}, X \times \{x_0\})$ apply the relative Mayer-Vietoris sequence with $A = X \times S^n \setminus \{x_0\}$, $B = X \times S^n \setminus \{x_1\}$ ($x_1 \neq x_0$), and $C, D = X \times \{x_0\}$. Then we have

$$\begin{aligned} \cdots \rightarrow H_i(X \times S^n \setminus \{x_0\}, X \times \{x_0\}) \oplus H_i(X \times S^n \setminus \{x_0\}, X \times \{x_1\}) &\rightarrow H_i(X \times S^n, X \times \{x_0\}) \\ \rightarrow H_{i-1}(X \times S^{n-1}, X \times \{x_0\}) &\rightarrow H_{i-1}(X \times S^n \setminus \{x_0\}, X \times \{x_0\}) \oplus H_{i-1}(X \times S^n \setminus \{x_0\}, X \times \{x_1\}) \rightarrow \cdots \end{aligned}$$

However the terms on the left and the right are all zero. As such this induces an isomorphism between $H_i(X \times S^n, X \times \{x_0\}) \cong H_{i-1}(X \times S^{n-1}, X \times \{x_0\})$.

Finally apply the second isomorphism n times to the first isomorphism to get $H_i(X \times S^n) \cong H_i(X \times \{x_0\}) \oplus H_{i-n}(X \times S^0, X \times \{x_0\})$. However the relative homology will make one of the two copies of X in $X \times S^0$ null. As such $H_{i-n}(X \times S^0, X \times \{x_0\}) \cong H_{i-n}(X)$.

Therefore $H_i(X \times S^n) \cong H_i(X) \oplus H_{i-n}(X)$ for all i, n where we assume $H_i(X) = 0$ for $i < 0$. \square

Problem 2 (2.3.3). Show that if \tilde{h} is a reduced homology theory then $\tilde{h}_n(\text{point}) = 0$ for all n . Deduce that there are suspension isomorphisms $\tilde{h}_n(X) \cong \tilde{h}_{n+1}(SX)$ for all n .

Proof. First note that if we wedge the point with itself that we get a point back. Then using the wedge axiom we have $\tilde{h}_i(\bigvee_1^2 * = *)$ is isomorphic to $\tilde{h}_i(*) \oplus \tilde{h}_i(*)$. The only way this could occur would be if $\tilde{h}_i(*) \cong 0$ for all i .

For the suspension use the Mayer-Vietoris sequence along with the fact that the cone of a space is contractable to get

$$\cdots \longrightarrow \tilde{h}_{i+1}(CX) \oplus \tilde{h}_{i+1}(CX) \longrightarrow \tilde{h}_{i+1}(SX) \longrightarrow \tilde{h}_i(X) \longrightarrow \tilde{h}_i(CX) \oplus \tilde{h}_i(CX) \longrightarrow \cdots$$

Which is equivalent to

$$0 \longrightarrow \tilde{h}_{i+1}(SX) \longrightarrow \tilde{h}_i(X) \longrightarrow 0$$

This show that $\tilde{h}_i(X) \cong \tilde{h}_{i+1}(SX)$ for all i . \square

Problem 3 (2.B.3). Let $(D, S) \subset (D^n, S^{n-1})$ be a pair of subspaces homeomorphic to (D^k, S^{k-1}) , with $D \cap S^{n-1} = S$. Show that the inclusion $S^{n-1} - S \hookrightarrow D^n - D$ induces an isomorphism on homology. [Glue two copies of (D^n, D) to the two ends of $(S^{n-1} \times I, S \times I)$ to produce a k -sphere in S^n and look at the Mayer-Vietoris sequence for the complement of this k -sphere.]

Proof. Using the construction listed in the problem decompose $S^n \setminus S^k$ as $A \cup B$ where $A = (S^{n-1} \times I, S \times I)$ filled in with $D^n \setminus D$ on one side and B as the filling on the other. Then using the fact that taking the Cartesian product with the interval does not change the homology and applying the Mayer-Vietoris sequence we get:

$$\cdots \longrightarrow H_i(S^{n-1} \setminus S) \longrightarrow H_i(D^n \setminus D) \oplus H_i(D^n \setminus D) \longrightarrow H_i(S^n \setminus S^k) \longrightarrow \cdots$$

Need to show that everything is zero unless $i = n - k - 1, n - k - 2$. Otherwise make it work. \square