

**Problem 1** (22). *Prove by induction on dimension the following facts about the homology of a finite-dimensional CW complex  $X$ , using the observation that  $X^n/X^{n-1}$  is a wedge sum of  $n$ -spheres.*

- (a) *If  $X$  has dimension  $n$  then  $H_i(X) = 0$  for  $i > n$  and  $H_n(X)$  is free.*
- (b)  *$H_n(X)$  is free with basis in bijective correspondence with the  $n$ -cells if there are no cells of dimension  $n - 1$  or  $n + 1$ .*
- (c) *If  $X$  has  $k$   $n$ -cells then  $H_n(X)$  is generated by at most  $k$  elements.*

$$\cdots \rightarrow \tilde{H}_k(X^{n-1}) \rightarrow \tilde{H}_k(X^n) \rightarrow \tilde{H}_k(X^n/X^{n-1}) \rightarrow \tilde{H}_{k-1}(X^{n-1}) \rightarrow \tilde{H}_{k-1}(X^n) \rightarrow \tilde{H}_{k-1}(X^n/X^{n-1}) \rightarrow \cdots$$

*Proof.* (a) Suppose that  $X$  has dimension zero. Then  $X$  is the disjoint union of points and as such from Hatcher we have that  $\tilde{H}_0(X) \cong \bigoplus_{x \in X \setminus x_0} \mathbb{Z}$  and  $\tilde{H}_k(X) = 0$  for  $k > 0$  where  $x_0$  is just some point.

Now assume that when a finite-dimensional CW complex  $X$  has dimension  $j < n$  that  $H_j(X)$  is free and  $H_i(X) = 0$  for  $i > n$ . Let  $X$  be an  $n + 1$  dimensional CW complex. From Hatcher we have the long exact sequence

$$\cdots \rightarrow \tilde{H}_k(X^{n-1}) \rightarrow \tilde{H}_k(X^n) \rightarrow \tilde{H}_k(X^n/X^{n-1}) \rightarrow \tilde{H}_{k-1}(X^{n-1}) \rightarrow \tilde{H}_{k-1}(X^n) \rightarrow \tilde{H}_{k-1}(X^n/X^{n-1}) \rightarrow \cdots$$

If  $k > n$  by induction we have

$$0 \rightarrow \tilde{H}_k(X^n) \rightarrow 0$$

which forces  $\tilde{H}_k(X^n) = 0$ . Otherwise if  $k = n$  we have

$$0 \rightarrow \tilde{H}_n(X^n) \rightarrow \bigoplus_{\sigma \text{ an } n\text{-simplex}} \mathbb{Z} \rightarrow \cdots$$

Since the sequence is exact we have that  $\tilde{H}_n(X^n)$  is isomorphic to a subgroup of a free abelian group and as such is free abelian.

Therefore by induction if  $X$  has dimension  $n$  then  $H_i(X) = 0$  for  $i > n$  and  $H_n(X)$  is free.

- (b) Let  $n = 0$ . From Hatcher we have that  $H_0(X)$  is free abelian with a generator for each 0-cell. Now assume that for dimension  $j < n$  if there are no  $j - 1$  or  $j + 1$  simplices that  $H_j(X)$  is free with basis in bijective correspondence with the  $j$ -cells. Then let  $X$  be a finite-dimensional CW complex with no  $n$ -simplices. Then via the same exact sequence from above we get

$$\tilde{H}_n(X^{n-1}) \cong 0 \rightarrow \tilde{H}_n(X^n) \rightarrow \tilde{H}_n(X^n/X^{n-1}) \rightarrow \tilde{H}_{n-1}(X^{n-1} \cong 0)$$

Which induces an isomorphism between the remaining groups. Since  $\tilde{H}_n(X^n/X^{n-1})$  has a generator for each  $n$  simplex it follows that  $\tilde{H}_n(X^n)$  does as well.

Next suppose that there are no  $n + 1$  simplices.

(c)

□

**Problem 2** (26). *Show that  $H_1(X, A)$  is not isomorphic to  $\tilde{H}_1(X/A)$  if  $X = [0, 1]$  and  $A$  is the sequence  $1, \frac{1}{2}, \frac{1}{3}, \dots$  together with its limit 0. [See Example 1.25.]*

*Proof.* We can see that  $H_1(X, A) \cong \bigoplus_1^\infty \mathbb{Z}$  from the short exact sequence

$$0 \longrightarrow H_1(X, A) \longrightarrow H_0(A) \cong \bigoplus_0^\infty \mathbb{Z} \longrightarrow H_0(X) \cong \mathbb{Z} \longrightarrow 0$$

as this short exact sequence enforces that  $H_1(X, A) \oplus \mathbb{Z} \cong \bigoplus_0^\infty \mathbb{Z}$ . Thus  $H_1(X, A)$  has a countable number elements.

However if we examine the space  $X/A$  we can see that it is homeomorphic to the Hawaiian earring. Moreover there are a countable number of one simplices and a single zero simplex. Thus  $\tilde{H}_1(X/A)$  is not finitely generated and as such by Hatcher (49) we have that  $\tilde{H}_1(X/A)$  is uncountable.

Therefore  $H_1(X, A)$  and  $\tilde{H}_1(X/A)$  are not isomorphic.  $\square$

**Problem 3** (27). Let  $f : (X, A) \rightarrow (Y, B)$  be a map that both  $f : X \rightarrow Y$  and the restriction  $f : A \rightarrow B$  are homotopy equivalences.

- (a) Show that  $f_* : H_n(X, A) \rightarrow H_n(Y, B)$  is an isomorphism for all  $n$ .
- (b) For the inclusion  $f : (D^n, S^{n-1}) \rightarrow (D^n, D^n - \{0\})$ , show that  $f$  is not a homotopy equivalence of pairs —there is no  $g : (D^n, D^n - \{0\}) \rightarrow (D^n, S^{n-1})$  such that  $fg$  and  $gf$  are homotopic to the identity through maps of pairs. [Observe that a homotopy equivalence of pairs  $(X, A) \rightarrow (Y, B)$  is also a homotopy equivalence for the pairs obtained by replacing  $A$  and  $B$  by their closures.]

*Proof.* (a) From Hatcher we have the long exact sequences with morphisms between them like so:

$$\begin{array}{ccccccccc} \cdots & \longrightarrow & H_n(A) & \longrightarrow & H_n(X) & \longrightarrow & H_n(X, A) & \longrightarrow & H_{n-1}(A) & \longrightarrow & H_{n-1}(X) & \longrightarrow & \cdots \\ & & \downarrow f_* & & \downarrow f_* & & \downarrow f_* & & \downarrow f_* & & \downarrow f_* & & \\ \cdots & \longrightarrow & H_n(B) & \longrightarrow & H_n(Y) & \longrightarrow & H_n(Y, B) & \longrightarrow & H_{n-1}(B) & \longrightarrow & H_{n-1}(Y) & \longrightarrow & \cdots \end{array}$$

Since  $f$  is a homotopy equivalence for  $A$  and  $X$  the left 2 and the right 2  $f_*$ s are isomorphisms. Thus by the five lemma the center map  $f_*$  is an isomorphism.

- (b) Suppose otherwise. Then we have a map  $g : (D^n, D^n \setminus \{0\}) \rightarrow (D^n, S^{n-1})$  such that  $g$  is a homotopy equivalence. Then since  $g : D^n \setminus \{0\} \rightarrow S^{n-1}$  is continuous it will force  $g : D^n \rightarrow S^{n-1}$  to send 0 to  $S^{n-1}$  as well as it is continuous. Since  $g$  extends is then a map from  $D^n$  that extends to a disc it is null and therefore the zero map on homology.

However for  $k = n - 1$  we have a map

$$H_{n-1}(D^n \setminus \{0\}) \cong \mathbb{Z} \rightarrow H_n(S^{n-1}) \cong \mathbb{Z}$$

which is both the zero map and an isomorphism since it is a homotopy equivalence. This is a contradiction.

Therefore there the inclusion map is not a homotopy equivalence of pairs for  $(D^n, S^{n-1})$  and  $(D^n, D^n \setminus \{0\})$ .  $\square$