

a
Relative abundance
Number of clusters
coassembly_timepoint_all
coassembly_animal
coassembly_timepoint_cage
multicoverage_animal
multicoverage_timepoint_cage
single_coverage
multisplit_timepoint_all
multisplit_timepoint_cage
multisplit_animal
multicoverage_timepoint_all
120
180
Bacillota
Bacillota_B
Bacillota_A
Actinomycetota
Pseudomonadota
Bacteroidota
Verrucomicrobiota
No taxonomy
b
Single
Multi-coverage
Coassembly
Multi-split
animal
timepoint all
timepoint cage
Completeness
100
90
80
70
c
Contamination
10.0
7.5
5.0
2.5
0.0
d
PCoA2 (17.5%)
PCoA1 (22.6%)
0
-2
-4
2
4
Number of strategies
1
3
5
7
9
e
single_coverage
multicoverage_animal
multicoverage_timepoint_all
multicoverage_timepoint_cage
coassembly_animal
coassembly_timepoint_all
coassembly_timepoint_cage
multisplit_animal
multisplit_timepoint_all
multisplit_timepoint_cage
KEGG modules
70%
100%
f
multicoverage_timepoint_all
coassembly_timepoint_all
Average genome completeness
93.5
93.0
92.5
92.0
Average function recovery
0.95
0.96
0.97
0.98