

Inferring Lensing Mass in the Universe from Photometric Catalog Data

Thomas E. Collett^{1*}, Philip J. Marshall², Matthew W. Auger¹, Stefan Hilbert³, Sherry H. Suyu⁴, Zachary Greene⁴, Tommaso Treu^{4†}, Christopher D. Fassnacht⁵, Lèon V. E. Koopmans⁶, Roger D. Blandford³

¹*Institute of Astronomy, University of Cambridge, Madingley Rd, Cambridge, CB3 0HA, UK*

²*Dept. of Physics, University of Oxford, Keble Road, Oxford, OX1 3RH, UK*

³*Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, 452 Lomita Mall, Stanford, CA 94035, USA*

⁴*Dept. of Physics, University of California, Santa Barbara, CA 93106, USA*

⁵*Dept. of Physics, U.C. Davis, Davis, CA 95616, USA*

⁶*Kapteyn Astronomical Institute, University of Groningen, P.O.Box 800, 9700 AV Groningen, The Netherlands*

to be submitted to MNRAS

ABSTRACT

Strong gravitational lenses can provide high precision cosmological distance measurements; one of the most important sources of systematic error in these estimates is the additional mass along the line of sight to the source. Weak lensing by galaxies and groups in this beam can contribute an “external convergence” κ_{ext} that dominates the uncertainty in the inferred distances; likewise, it may also contaminate estimates of the luminosity of objects at high redshift (such as reionisation-era proto-galaxies, and type Ia supernovae standard candles). We characterise this uncertainty by marginalising over a probability density function for κ_{ext} : here, we investigate the use of a simple halo model to estimate $\text{Pr}(\kappa_{\text{ext}})$ given noisy estimates of the photometric redshift and stellar mass of galaxies observed in optical imaging surveys of the field in question. Tests using mock catalogs from the Millenium Simulation and comparing to ray-traced κ values, we find that ... The $\text{Pr}(\kappa_{\text{ext}})$ from this reconstruction process are narrower than the expected distribution of κ_{ext} values over an ensemble of time delay lenses: incorporating the photometric catalogs in a joint analysis of 100 lenses leads to an increase in precision of factor of 1.X over simple averaging. We investigate several sources of bias and determine ...

Key words: gravitational lensing – methods: statistical – galaxies: halos – gaaxies: mass function – cosmology: observations

1 INTRODUCTION

Averaged over large areas of sky, the weak gravitational lensing effect can be used to constrain the statistical properties of the matter density in the Universe: every distant object we observe has had its shape distorted, and size and total brightness magnified (or demagnified) by a compound lens constructed from all the mass distributed between us and it.

This fact makes gravitational lensing a potentially important source of systematic error for any estimate of luminosity or distance, an issue addressed for *e.g.* type Ia supernovae by ?? and for high redshift galaxies by ? among

others. In time delay lens cosmography the effect can be particularly important. ? found that the lensing effect of mass structures along the (unusually over-dense) line of sight to the quadruply-imaged radio source B1608+656, the so-called “external convergence,” was the dominant source of uncertainty in the inferred cosmological parameters.

How can we correct for the line of sight lensing error, and make accurate measurements through the Universe? As ? noted, when inferring global quantities (such as the cosmological parameters), averaging the results from a large number of independent objects will tend to reduce the error, as the ensemble-averaged convergence approaches zero for an unbiased sample (?). If the sample is biased, however, some residual systematic error will remain while the statistical uncertainty decreases – and this will be the case for

* t.collett@ast.cam.ac.uk

† Packard Research Fellow

any sample selected by the brightness of the lensed object, or any other quantity that depends on κ_{ext} . The latter class includes time delay lenses, which are detected based on the separation of their images, which increases with κ . Large samples of lenses are expected to be discovered in the next decade in ground-based optical imaging surveys (?) – but these samples will be image separation-limited, due to the surveys’ relatively poor (~ 1 arcsec) image quality. Some of the current sample of known time delay lenses were detected in surveys like this: we might expect them to be somewhat biased towards high values of κ_{ext} .

There are two broad approaches one could take at this point. First, one could seek to characterise the lensed objects’ selection function, and use this understanding to derive a suitable correction to cosmological parameters inferred assuming no κ_{ext} at all. To do this, one needs information about the unlensed population of objects, and the population of lenses, and how the objects were detected – or to infer these hierarchically during the analysis. Alternatively, one could simply accept all lenses provided, regardless of their selection, and try to use any additional information that is available to estimate the external convergence in each system on a case by case basis, and hence optimize their joint analysis to produce a better-constrained result. In this work we follow the second approach, and use measurements of the positions, magnitude and colours of galaxies close to time delay lens lines of sight to try to infer the external convergence at the lens position.

Previous attempts at reconstructing the line of sight mass distribution in strong lens fields have focused on spectroscopic surveys to ascertain group membership, and on understanding the external shear apparently required by the strong lens models. The probability density function (PDF) $\text{Pr}(\kappa_{\text{ext}})$ is the target for any reconstruction analysis, and the key quantity for any ensemble analysis, since by marginalising over it we can correctly propagate the uncertainties due to weak lensing by line of sight mass structure. **More needed here. Previous work on environments: Keeton, Zabludoff, Auger, Momcheva, Wong, others.** Most recently, ? ran Monte Carlo reconstructions of the mass in nine lens fields, and compared the resulting predicted shear distributions with the lens model external shear. Their simulations show most of the shear being generated by bright galaxies within 2 arcminutes of the lens, and that both line of sight structures and the group of galaxies in each lens plane contribute significant proportions of the shear. The significant discrepancies between the lens model shear and the Monte Carlo predictions are attributed to both the environment modelling and the strong lens modelling itself.

These observational investigations have provided very useful insight into the problem. Coming from the opposite direction, large numerical simulations have been used to predict, from first principles the distribution of likely external shears and convergences along strong lens lines of sight. **More needed here. Holder & Schechter, Dalal & Watson.** ? performed ray-tracing experiments through the Millenium Simulation, generating a predicted κ_{ext} at every position in a simulated sky. The first step towards using this excellent calibration resource was taken by ?, who selected Millenium lines of sight that a) were found to contain a strong lens and b) had comparable apparent galaxy

overdensity in a 45 arcsec radius aperture down to $i < 24$. This overdensity could be compared to the overdensity of the B1608+656 line of sight (?), and hence an estimate of $\text{Pr}(\kappa_{\text{ext}})$ could be made.

What has not been investigated before, but is the subject of this paper, is the use of simulations to *calibrate* light-cone reconstruction methods. Ray-tracing through the simulations gives a “truth” that we can attempt to recover. In this paper we use the Millenium Simulation mock galaxy catalogs and ray-traced lensing effects to test the accuracy of line of sight mass reconstructions. Probabilistically assigning mass and redshift to every observed galaxy in the field around a time delay lens, we generate Monte Carlo sample line of sight mass distributions, and estimate the external convergence due to each one. This gives us a $\text{Pr}(\kappa_{\text{ext}})$ that can be propagated into further joint inferences. We focus on time delay lens cosmography, and quantify our results in terms of the uncertainty on the mean time delay distance to a test sample of N lenses (where for clarity each lens has the same deflector and source redshift, and the same lens model and time delay uncertainty).

We aim to answer the following questions:

- Faint galaxies and other dark structures will not appear in a photometric object catalog, but they will contribute convergence at some level. How much of the total external convergence in a time delay lens system comes from visible galaxies? How does this change as a function of magnitude cut?
- Can the true (ray-traced) convergence be recovered by halo model reconstruction? What scatter and residual bias are induced in the ensemble distance estimate? Which aspects of the model dominate these uncertainties?
- When faced with a newly-detected lens, surrounded by galaxies on the sky, we have some choices to make when planning follow-up observations. Which are the most important galaxies, with regard to the external convergence produced? Can nearby groups and clusters be straightforwardly accounted for? If lenses with such massive structures nearby are discarded, what impact does that have on the distance accuracy?

This paper is organized as follows. In Section 2 we review the relevant gravitational lensing theory, before introducing our simple reconstruction model in Section 3. We then test this model in two phases: first, in Section 4, with known redshift and halo mass for every galaxy in a lens field, in order to quantify the irreducible uncertainty due to unseen mass, and second, in Section 5, with realistic observational uncertainties on the field galaxies’ stellar masses and redshifts. We then propagate all these uncertainties through to the time delay distance in a toy sample of 100 lenses, in Section 6. We discuss our results in Section 7 before concluding in Section 8.

Throughout this paper magnitudes are given in the AB system (?) and we adopt standard “concordance” parameters for our reference cosmology, *i.e.* $h = 0.7$, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$, where the symbols indicate the Hubble Constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and the matter and dark energy density of the Universe in units of the critical density (e.g. ?).

2 THEORETICAL BACKGROUND

Convergence, or Ricci focusing, occurs when a gravitational lens focuses the rays in a raybundle. This focussing can cause distant objects to appear brighter, and larger than they would if the lens were removed. The convergence from an isolated mass sheet is the ratio of the projected surface mass density ($\Sigma(D_{\text{od}}\theta)$) divided by the critical surface mass density at the redshift of the mass sheet (Σ_{cr}),

$$\kappa(\theta) = \frac{\Sigma(D_{\text{od}}\theta)}{\Sigma_{\text{cr}}(z)} \quad (1)$$

where

$$\Sigma_{\text{cr}}(z) \equiv \frac{c^2 D_{\text{os}}}{4\pi G D_{\text{od}} D_{\text{ds}}} \quad (2)$$

and the D's are the angular diameter distances between the objects referred to in the subscripts: o is the observer, d is the deflector and s is the source. If the surface mass density of the sheet exceeds Σ_{cr} , multiple images of the source can occur, otherwise only one image will be observed, although that image will still be perturbed relative to the unlensed case.

Strong lensing occurs when a massive object and background source are almost perfectly aligned along a line of sight. The light from the background source is deflected by the lens galaxy; this deflection allows multiple images of the background source to form at stationary points of the time delay function. For an isolated lens, the time delay function can be calculated from

$$\Delta t(\theta, \beta) = \frac{1}{c} \frac{D_{\text{od}} D_{\text{os}}}{D_{\text{ds}}} (1 + z_{\text{d}}) \phi(\theta, \beta), \quad (3)$$

where θ is the observed source position, β is the unlensed source position, z_{d} is the redshift of the lens, $\phi(\theta, \beta)$ is the Fermat potential. The Fermat potential is given by

$$\phi(\theta, \beta) \equiv \left[\frac{(\theta - \beta)^2}{2} - \psi(\theta) \right], \quad (4)$$

where $\psi(\theta)$ is the lens potential, derived from the projected dimensionless surface mass density, $\kappa(\theta)$, by

$$\kappa(\theta) = \frac{1}{2} \nabla^2 \psi(\theta). \quad (5)$$

Strong lenses sit in our real Universe – they are not truly isolated; there is some mass along the line of sight coming from intervening structures. These structures may be physically associated with the lens e.g. other members of a group or cluster, or they may be the result of chance alignments at any redshift. Whilst it is rare for three galaxies to line up well enough for both of the background sources to be strongly lensed (??), the shear scale of dark matter halos makes a partial alignment – with the additional structure acting as a weak lens – a near certainty, indeed ? and ? find that there are no empty lightcones in a realistic universe. These additional structures induce an external shear which can be recovered in the mass-modelling of the lens, but they can also introduce a convergence that is undetectable from the image positions, shapes and relative fluxes; *the mass sheet-degeneracy* (see e.g. ??, for details).

If the intrinsic source brightness changes, the observed brightness in each image does not change simultaneously – there is a time delay, Δt , between the brightness change in each image. Despite the invariance of the image positions

and fluxes, under the addition of an external convergence the relative Fermat potential between the images is *not* invariant, and so the time delay changes. As such it is necessary to include κ_{ext} in our lens modelling, if cosmological parameters are to be estimated accurately and precisely from observed time delays (?).

If the mass sheet is not physically associated with the lens galaxy, it can lie at any redshift. The effects of such a mass-sheet can be projected back onto the lens plane in the form of an effective convergence. We define this effective convergence in two limits.

- No significant perturbations along the line of sight, and
- A single dominant perturber, relative to which all other perturbations are small.

We will refer to these cases as the magnification case (subscript μ in formulae), and the strong lensing case (subscript SL in formulae), respectively. We will ignore the additional case, of a compound lens, with several significant perturbers.

The magnification case has no primary lens plane, and is chiefly relevant for the magnifications of distant objects. In the absence of a primary lens plane, we are free to define the effective convergence on any point; the effective convergence is the same, independent of the plane chosen. It can be shown (e.g. ?) that if there are several mass sheets at redshifts $\{z_p\}$, with convergences $\{\kappa_p = \Sigma_p / \Sigma_{\text{cr}}(z_p)\}$, the effective convergence is given by

$$\kappa_{\text{effective}, \mu} = \sum_p \kappa_p. \quad (6)$$

In the strong lensing case, the presence of a significant perturber alters the geometry of the system, and provides a primary lens plane. It is the primary lens plane upon which we need to know $\kappa_{\text{effective}}$, since this is where the mass-sheet degeneracy is defined. Following Equations 20–32 of ?**keeton2003**, it can be shown that the relevant effective convergence is given by

$$\kappa_{\text{effective}, \text{SL}} = \frac{(1 - \beta_{12})(\kappa - \beta_{12}(\kappa^2 - \gamma^2))}{(1 - \beta_{12}\kappa)^2 - (\beta_{12}\gamma)^2} \quad (7)$$

where κ and γ are, respectively, the convergence and shear caused by the perturber, in their own plane and β_{12} is a geometrical scaling factor defined by

$$\beta_{12} = \frac{D_{12} D_{\text{os}}}{D_{\text{o1}} D_{1s}} \quad (8)$$

where the D's are again angular diameter distances: the 1 (2) in the subscript refers to the lower (higher) redshift object of the lens and perturber. For the case that the mass sheet is in the lens plane, $\beta_{12} = 0$ and $\kappa_{\text{eff}} = \kappa$; if the mass-sheet is in the plane of the observer, or source, $\beta_{12} = 1$ and $\kappa_{\text{eff}} = 0$; for a mass-sheet at any other redshift $0 < \beta_{12} < 1$. In the case of several mass sheets the effective convergences sum. In the limit that both κ and γ are small, Equation 7 reduces to

$$\kappa_{\text{effective}, \text{SL}} = (1 - \beta_{12})\kappa; \quad (9)$$

it is this $\kappa_{\text{effective}, \text{SL}}$ that we use as κ_{ext} for the strong lensing time delay analysis in 6.

The time-delay distance is defined as

$$D_{\Delta t} \equiv (1 + z_{\text{d}}) D_{\text{d}} D_{\text{s}} / D_{\text{ds}}, \quad (10)$$

with which Equation 3 can be rephrased as

$$\Delta t(\boldsymbol{\theta}, \boldsymbol{\beta}) = \frac{D_{\Delta t}}{c} \phi(\boldsymbol{\theta}, \boldsymbol{\beta}), \quad (11)$$

Since the time delays are proportional to $1/(1+\kappa_{\text{ext}})$, if there is any external convergence, that is not included in the lens modeling, then the time delay distance – inferred assuming $\kappa_{\text{ext}} = 0$ – will be $(1 - \kappa_{\text{ext}})$ less than the true value of $D_{\Delta t}$.

$$D_{\Delta t}^{\text{true}} = (1 - \kappa_{\text{ext}}) D_{\Delta t}^{\kappa_{\text{ext}}=0}. \quad (12)$$

We can overcome this degeneracy if we have additional knowledge of κ_{ext} .

For mass-sheets that are physically associated with the lens galaxy, the mass sheet affects the stellar dynamics of the lens galaxy, so dynamical observations can break the internal mass-sheet degeneracy by providing an additional estimate of the lens galaxy’s mass (e.g., ?).

In this work our aim is to quantify the lensing effect of mass-sheets that are not physically associated with the lens galaxy – mass-sheets lying in front or behind the lens – and the uncertainties induced by reconstructing those mass-sheets using simple prescriptions applied to potential astronomical observables. We also aim to quantify how uncertainties in the estimations of κ_{ext} propagate into uncertainties on the cosmological parameters which can be inferred from measurements of $D_{\Delta t}$.

3 HALO MODEL AND MASS DISTRIBUTION

For real lines of sight, it is nearly impossible to know the true κ_{ext} for a line of sight, and without a knowledge of $\kappa_{\text{ext}}^{\text{true}}$ it is hard to quantify if there is a bias induced by using different κ_{ext} estimators. A source with known unlensed brightness e.g. a strongly lensed type Ia Supernova **cite some SNe papers** or a source of known unlensed size **ideas?** would break the mass-sheet degeneracy and provide a value for κ_{ext} , but **no?** such systems are currently known.

Cosmological dark matter simulations can be used to generate mock lines of sight... **Stefan’s section on how the ray tracing was done and how halos are extracted from the MS n-body particles**

With this catalogue of halos, positions, redshifts etc. it is possible to attempt a reconstruction of the convergence induced by every halo near a line of sight. Cosmological dark matter simulations have shown that dark matter halos are well approximated by NFW profiles (?), with a density profile given by

$$\rho(r) = \frac{\rho_0}{(r/r_s)(1+r/r_s)^2}, \quad (13)$$

where r_s is a characteristic radius of the cluster, representing where the density slope transitions from r^{-1} to r^{-3} . This radius is related to the virial radius of the halo by $r_s = r_{200}/c$, where r_{200} is the radius at which

$$\rho(r_{200}) = 200\rho_{\text{crit}}(z), \text{ where } \rho_{\text{crit}}(z) \equiv \frac{3H^2(z)}{8\pi G} \quad (14)$$

and c is the concentration parameter, which can be estimated from the halo’s mass, using a mass–concentration relation. Typically more massive halos are less concentrated, but there is some scatter; we use the relation of ? to estimate c from the mass enclosed within r_{200} , which we denote

as M_{200} . The best fit relation of ? is given by

$$c_{200} = 4.67(M_{200}/10^{14}h^{-1}\text{M}_{\odot})^{-0.11}. \quad (15)$$

ρ_0 is calculated from the critical density and concentration parameter:

$$\rho_0 = \rho_{\text{crit}}(z) \frac{200}{3} \frac{c^3}{\ln(1+c) - c/(1+c)}. \quad (16)$$

Disturbingly, if one integrates the density profile of an NFW profile, the total mass is divergent. Similarly, if the universe is homogeneously populated with NFW halos, the projected surface mass along any line sight will also be divergent¹. Since infinite mass is unphysical, the profile must be truncated at some point. **previous truncation results**. Several truncation profiles have been suggested (e.g. ?), but beyond several virial radii, the amount of matter associated with a halo is likely to be low; ? showed that the Universe’s expansion can place an upper limit on the radius of NFW halos – for cluster scale halos, this happens at around 25 times the NFW scale radius. In Section 4 we investigate how different truncations affect the recovered total convergence for a line of sight, using the truncated NFW profile of ?

$$\rho(r) = \frac{\rho_{\text{NFW}}(r)}{1 + (r/r_t)^2}, \quad (17)$$

which is the same as the NFW profile in the limit that r_t goes to infinity. Defining x as the dimensionless projected radial distance R/r_s and τ as the dimensionless truncation radius r_t/r_s , ? derives the projected mass density, which is given by:

$$\Sigma_{\text{BMO}}(x) = \frac{2\tau^2}{(\tau^2 + 1)^2} \left(\frac{\tau^2 + 1}{x^2 - 1} (1 - \mathcal{F}(x)) + 2\mathcal{F}(x) - \frac{\pi}{\sqrt{\tau^2 + x^2}} + \frac{(\tau^2 - 1)\mathcal{L}(x, \tau)}{\tau\sqrt{\tau^2 + x^2}} \right) \quad (18)$$

where $\mathcal{F}(x)$ and $\mathcal{L}(x, \tau)$ are defined as

$$\mathcal{F}(x) \equiv \begin{cases} \frac{\cos^{-1}(1/x)}{\sqrt{x^2 - 1}} & (x > 1) \\ \frac{4 - x^2}{3} & (x = 1) \\ \frac{\cosh^{-1}(1/x)}{\sqrt{1 - x^2}} & (x < 1) \end{cases} \quad (19)$$

$$\mathcal{L}(x, \tau) = \ln \left(\frac{x}{\sqrt{\tau^2 + x^2} + \tau} \right) \quad (20)$$

Whilst halos contribute a positive convergence to κ_{ext} , voids contribute a negative convergence. When a ray bundle passes through a region of space that is less dense than $\rho_{\text{cr}}(z)$ the rays are de-focused². Whilst the full raytraced solution takes into account the effect of voids, our reconstruction cannot – the halo catalogue does not include voids. In principle

¹ At large radius, $\Sigma_{\text{nfw}} \propto R^{-2}$, but the differential number of halos centred within an annulus of width dR is given by $dN_{\text{annulus}} \propto R dR$, so $\Sigma_{\text{total}} \propto \int_0^\infty R^{-1} dR$, which diverges logarithmically

² In principle, the convergence caused by halos, is also caused by the overdensity, but we neglect this effect since halos are typically much more dense than the intergalactic medium

the absence of a halo could be taken to infer a void, but the under-density of such a void is hard to infer. Instead, we shift our κ_{ext} values so that $\langle \kappa_{\text{ext}} \rangle = 0$. We refer to this corrective term as κ_{void} . The void correction introduces a source of error, since the inferred value of κ_{void} will depend on the properties of the sightlines reconstructed, but this error shrinks with the number of sightlines reconstructed. To avoid potential biases we calculate κ_{void} from randomly selected sightlines. The total κ_{ext} along the line of sight is given by

$$\kappa_{\text{ext}} = \left(\sum_{\text{halos}} \kappa_{\text{halo}} \right) - \kappa_{\text{void}}. \quad (21)$$

4 TESTS WITH KNOWN HALO MASS AND REDSHIFT

Given the halo mass catalogue from the Millenium Simulation, and convergence maps produced by ray-tracing through the MS's n-body particles, it is possible to test the validity of the prescription outlined in Section 3. We do this using κ_{raytrace} as a proxy for κ_{true} . The reconstruction prescription is tested given perfect knowledge of the halos' virial masses and redshifts; this provides us with a measure of the intrinsic errors induced by treating matter as truncated NFW halos. To quantify this 10000 lines of sight are chosen from random positions in the catalogue, and for each line of sight we reconstruct the convergences from every halo within 5 arcminutes and compare the reconstructed kappas with those from the ray tracing results. The standard deviation on $\kappa_{\text{raytrace}} - \kappa_{\text{reconstruction}}$ shown in Figure ?? . It is clear from Figure ?? that as long as the truncation is greater than a few virial radii, the choice of truncation radius does not broadly affect the scatter on $\kappa_{\text{raytrace}} - \kappa_{\text{reconstruction}}$; from now on we choose to truncate the halos at $10R_{200}$. Since $\sigma(\kappa_{\text{raytrace}} - \kappa_{\text{reconstruction}})$ does not go to zero, it is clear that our halo based prescription does not perfectly replicate the n-body results, we refer to this as the intrinsic error of the prescription. The intrinsic error has a standard deviation of $\sigma_{\text{intrinsic}} \sim 0.018$, although this figure is inflated by a few outliers in the tails of $\text{Pr}(\kappa_{\text{raytrace}} - \kappa_{\text{reconstruction}})$. Whilst we do not focus on shear in this work, such outliers would likely be rejected by their reconstructed shear not matching the shear of the lens model (see ? for the practicalities of this), this makes our estimate of the intrinsic error pessimistic.

this paragraph is phrased badly If $\kappa_{\text{reconstruction}}$ is to be used in a real survey it is important that it is not a biased estimator of the true κ_{ext} . Ideally one wants $\langle \kappa_{\text{reconstruction}} - \kappa_{\text{true}} \rangle$ to equal 0 and $\kappa_{\text{reconstruction}} - \kappa_{\text{true}}$ to be independent of $\kappa_{\text{reconstruction}}$, since the void correction ensures $\langle \kappa_{\text{reconstruction}} \rangle = 0$ any bias in $\langle \kappa_{\text{reconstruction}} - \kappa_{\text{raytrace}} \rangle$ comes from the variations in κ_{raytrace} . With 8.2×10^8 pixels, uniformly spaced over 196 deg^2 of simulated sky, $\langle \kappa_{\text{raytrace}} \rangle = -0.00014$. However, it is improbable that such a large number of lines of sight could be reconstructed in reality and the mean of κ_{true} for a small subset of sightlines will not necessarily equal zero. We find this error to be

$$\sigma(\kappa_{\text{void}}) \simeq 0.036/N^{\frac{1}{2}}, \quad (22)$$




Figure 1.




Figure 2.

where N is the number of lines of sight, although $\text{Pr}(\kappa_{\text{void}})$ has positive skew for small N . Care must be taken that the calibration lines of sight are not biased in κ_{ext} or the final $\text{Pr}(\kappa_{\text{void}})$ will also be biased. For the rest of this work we use 1000 randomly chosen calibration lines of sight to calculate κ_{void} , introducing an error on κ_{ext} of 0.001. 1000 calibration lines of sight is unfeasible currently, but should be reasonable in the LSST era. Since the true $\langle \kappa_{\text{raytrace}} \rangle = -0.00014$ is two orders of magnitude smaller than intrinsic error of the halo prescription, $\kappa_{\text{reconstruction}}$ can be seen as unbiased given a large enough sample of lines of sight. Figure ?? shows that there is no statistically significant evidence that $\kappa_{\text{reconstruction}}$ is a biased estimator, regardless of the value of $\kappa_{\text{reconstruction}}$.

The error from the void correction, and the intrinsic error from the halo based prescription are unavoidable errors, even with perfect knowledge of halo mass and position. Before investigating the errors caused by imperfect knowledge of halo mass and redshift, we investigate the errors induced by a magnitude limited reconstruction and a field of view limited reconstruction. The halos in our catalogue are given magnitudes by the semi-analytic model of ? and by applying magnitude cuts to our catalogue, we can investigate the scatter caused by unobserved halos. The majority of the convergence comes from halos with an i magnitude between 18 and 24. Figure 4 shows that the width of $\kappa_{\text{reconstruction}} - \kappa_{\text{raytrace}}$ decreases quickly between $i = 18$ and $i = 24$. Objects brighter than $i = 18$ are either too rare




Figure 3.

or too close to the observer to make a significant contribution to the convergence. Objects fainter than $i = 24$ are too small to be important unless they are extremely close to the line of sight where it is likely that neglecting stellar mass and using a spherical NFW prescription are too naive to adequately reconstruct κ_{ext} - it is likely that these ultra-faint halos (or substructures) are the major source of the scatter in $\kappa_{\text{reconstruction}} - \kappa_{\text{raytrace}}$, but a deeper survey will not be sufficient to decrease the size of this scatter. Since a large number of calibration lines of sight are needed to shrink the width of $\text{Pr}(\kappa_{\text{void}})$, it may prove unrealistic to reconstruct the calibration lines of sight out to 5 arcminutes. ? showed that the majority significant perturbers are within 15 arcseconds of the lens. Figure 4 shows the width of $\kappa_{\text{reconstruction}} - \kappa_{\text{raytrace}}$ as a function of reconstruction radius; in almost all cases there is no significant contribution to κ from halos that are centred more than 1 arcminute away, although large groups or clusters can sometimes still make a contribution as far out as four arcminutes. For the rest of this work we will continue to model all of the halos out to 5 arcminutes with an i magnitude greater than 26, although a reconstruction with fields going out to two arcminutes and down to $i < 24$ would do almost as well.

5 TESTS ON MOCK CATALOGS

Consider three sources of scatter in the halo model: z_{phot} , $M_{\text{star}}-z_{\text{phot}}$, and $M_{\text{halo}}-M_{\text{star}}, z_{\text{phot}}$. Mechanism for including scatter in model: draw samples from PDFs. Result is not one predicted kappa, but one $P(\text{kappa})$ for each lightcone. This is what is required by single-lens analyses.

We can draw samples from this PDF and ask:

How is the kappa inference offset and blurred by uncertainty on redshift, M_{star} and M_{halo} ?

What is the effect of using the wrong $M_{\text{star}}-M_{\text{halo}}$ relation?

6 TIME DELAY DISTANCES

Folding the kappa uncertainties into time delay distance uncertainties. Toy model: same z_l and z_s , allows simple product of pdfs to obtain estimate of $P(D)$.

What is the expected bias and scatter (centroid position and width) of $P(D)$ as a function of N lenses? How does reconstruction compare with $P(D)$ derived using $P(\text{kappa}-N45)$ for each lightcone? How does reconstruction compare with $P(D)$ derived assuming $\text{kappa}=0$ for each lightcone?

Also: compare with simple averaging. This will fail if lenses are selected to live on over-dense lines of sight. Stefan's plot in Suyu et al shows effect of selection - what is the resulting bias? Few percent? This is the target to beat, need to find it out.

Make mock sample of lenses, all wit B1608 z_d and z_s , but with different kappas. Draw kappas from Stefan's *lens* distribution. Can we correct for the selection effect?

7 DISCUSSION

Why reconstruction works or doesn't work.

Major sources of uncertainty.

Ideas for improving the procedure

Emphasize that this can be done before monitoring time is invested.

8 CONCLUSIONS

To remind you, this is what we did.

Our conclusions can be summarised as follows:

- We found this.
- And we found this too.

ACKNOWLEDGMENTS

We thank Risa Wechsler and Peter Behroozi for useful discussions and suggestions. TEC acknowledges support from STFC in the form of a research studentship. PJM was given support by the TABASGO and Kavli foundations, and the Royal Society, in the form of research fellowships. MWA SH SHS TT acknowledges support from the NSF through CAREER award NSF-0642621, and from the Packard Foundation through a Packard Research Fellowship. LVEK acknowledges the support by an NWO-VIDI programme subsidy (programme number 639.042.505). This research is supported in part by the National Science Foundation under Grant No. PHY99-07949.

APPENDIX A: TRUNCATED Λ CDMHALOS

APPENDIX B: DERIVING THE $M_* = M_{200}$ RELATION

APPENDIX C: SMOOTH COMPONENT MODELS

This paper has been typeset from a $\text{T}_{\text{E}}\text{X}/\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}$ file prepared by the author.