Chapter 1

Results

Figure 1.1 shows the performance trends for the four training tasks (*Path, Rings, Pillars* and *Exchange*), where the performance at each repetition is the average among the subjects in the assisted or control group. Apart from the *Pillars* task, the trends are increasing both for the assisted group and the unassisted group. Most significantly, the performance in the assisted group is consistently higher than the one in the control group in all training tasks.

The standard deviation of the performance score is also displayed at each repetition with error bars that denote the range $\left[\mu - \frac{1}{2}\sigma; \mu + \frac{1}{2}\sigma\right]$, with μ and σ the mean and standard deviation of the performance score, respectively. No significant difference is observed between the two groups and neither it is apparent a recognizable trend of deviation as a function of repetitions, *i.e.* the error bars do not get wider nor narrower as the repetitions increase.

The gradual removal of haptic assistance that occurred daily during the training phase does not seem to have affected the performance of the subjects in the assisted group. All performance trends increase steadily from repetition 1 to repetition 12, without any significantly visible performance plateau.

The performance on the four validation tasks (*Thymectomy, Nephrectomy, Liver Resection* and *Suturing*) recorded on the last day of the experimental phase is shown in Figure 1.2 with boxplots. For each task, the graph reports the distribution of performances collected from the 4 subjects executing 3 repetitions. Crucially, neither the subjects in the assisted group nor the ones in the control group were guided with Virtual Fixtures (VFs) on these tasks, nevertheless the performance recorded from the assisted subjects is distributed on higher values for all the tasks.

For both the assisted and unassisted group, the performance is distributed on different values




Figure 1.1: Performance Trend for the four Training Tasks. The value at each repetition is averaged across the subjects belonging to the same group. Error bars show the standard variation at each repetition




Figure 1.2: Boxplots of the average performance of assisted subjects (blue) and unassisted subjects (orange) on the four validation tasks

depending on the task, and so are the average values. This factor reflects how the tasks are different in terms of difficulty, and it is also reflected in the standard deviation of the performance. The standard deviation of distribution is, again, inconsistend when comparing the one calculated from group A and group C: in *Thymectomy* and *Nephrectomy* performance scores of assisted subjects are more spread out than the ones of unassisted subjects, while in *Liver Resection* the variation is comparable and in *Suturing* the performance of unassisted subjects has a higher deviation.

| Thymectomy | | | | |
|------------|-------|---------|--------|--|
| | Mean | STD | Median | |
| CONTROL | 0.38 | 0.01 | 0.37 | |
| ASSISTED | 0.41 | 0.04 | 0.42 | |
| % VAR | 9.83% | 193.94% | 13.07% | |

| Nephrectomy | | | | | |
|-------------|-------|---------|--------|--|--|
| | Mean | STD | Median | | |
| CONTROL | 0.68 | 0.06 | 0.71 | | |
| ASSISTED | 0.74 | 0.15 | 0.69 | | |
| % VAR | 8.50% | 167.46% | -2.71% | | |

| LiverResection | | | | | |
|----------------|--------|--------|--------|--|--|
| | Mean | STD | Median | | |
| CONTROL | 0.73 | 0.12 | 0.68 | | |
| ASSISTED | 0.90 | 0.15 | 0.83 | | |
| % VAR | 23.42% | 28.85% | 21.54% | | |

| Suturing | | | | | |
|----------|-------|---------|--------|--|--|
| | Mean | STD | Median | | |
| CONTROL | 0.84 | 0.05 | 0.85 | | |
| ASSISTED | 0.91 | 0.03 | 0.92 | | |
| % VAR | 8.38% | -25.39% | 8.44% | | |

Quantitative results are reported in the tables above: for each task, the mean, standard deviation and median values of performance are compared between the assisted and the control group. Given the data scarcity and their non-Gaussian distribution, the most meaningful conclusions will be drawn from the median values. Apart from *Nephrectomy* showing a slightly reduced median performance on assisted subject (accompanied, however, by a much larger standard deviation, as per Figure 1.2), all other tasks present an increase the median

performance, as high as +21.54%. The variation in mean performance is, instead, always positive.