



A new framework for Kolyvagin systems

Alberto Angurel

University of Nottingham

01/12/2025



Motivation: Birch and Swinnerton-Dyer conjecture

- Let E/\mathbb{Q} be an elliptic curve.



Motivation: Birch and Swinnerton-Dyer conjecture

- Let E/\mathbb{Q} be an elliptic curve.
- $E(\mathbb{Q})$ is a finitely generated abelian group, so $E(\mathbb{Q}) \cong \mathbb{Z}^r \times T$.



Motivation: Birch and Swinnerton-Dyer conjecture

- Let E/\mathbb{Q} be an elliptic curve.
- $E(\mathbb{Q})$ is a finitely generated abelian group, so $E(\mathbb{Q}) \cong \mathbb{Z}^r \times T$.
- Computing T is easy, but the rank is hard.



Motivation: Birch and Swinnerton-Dyer conjecture

- Let E/\mathbb{Q} be an elliptic curve.
- $E(\mathbb{Q})$ is a finitely generated abelian group, so $E(\mathbb{Q}) \cong \mathbb{Z}^r \times T$.
- Computing T is easy, but the rank is hard.

BSD conjecture

There exists an L -function

$$L(E, s) = P_\ell(\ell^{-s})^{-1}$$

where P_ℓ are **Euler factors**, such that

$$r = \text{ord}_{s=1} L(E, s)$$



Motivation: Birch and Swinnerton-Dyer conjecture

- Let E/\mathbb{Q} be an elliptic curve.
- $E(\mathbb{Q})$ is a finitely generated abelian group, so $E(\mathbb{Q}) \cong \mathbb{Z}^r \times T$.
- Computing T is easy, but the rank is hard.

BSD conjecture

There exists an L -function

$$L(E, s) = P_\ell(\ell^{-s})^{-1}$$

where P_ℓ are **Euler factors**, such that

$$r = \text{ord}_{s=1} L(E, s)$$

Theorem (Kolyvagin)

If $\text{ord}_{s=1} L(E, s) \leq 1$, then BSD holds true.



Idea: replace points with Galois cohomology

Fix a prime p and a natural number K .

Kummer map

There is an injection

$$E(\mathbb{Q})/p^k E(\mathbb{Q}) \hookrightarrow H^1(\mathbb{Q}, E[p^k])$$



Idea: replace points with Galois cohomology

Fix a prime p and a natural number K .

Kummer map

There is an injection

$$E(\mathbb{Q})/p^k E(\mathbb{Q}) \hookrightarrow H^1(\mathbb{Q}, E[p^k])$$

Proposition

The image of the Kummer map in the subgroup cut out by *local conditions*.



Idea: replace points with Galois cohomology

Fix a prime p and a natural number K .

Kummer map

There is an injection

$$E(\mathbb{Q})/p^k E(\mathbb{Q}) \hookrightarrow H^1(\mathbb{Q}, E[p^k])$$

Proposition


The image of the Kummer map in the subgroup cut out by *local conditions*.

Selmer group

This is known as a *Selmer group*, and it is the object that will be studied by the Euler system machinery.



General picture





Tate module

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Tate module

We will study the limit $T_p E = \varprojlim_k E[p^k]$, which is a free \mathbb{Z}_p -module of rank 2 endowed with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$.



Tate module

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Tate module

We will study the limit $T_p E = \varprojlim_k E[p^k]$, which is a free \mathbb{Z}_p -module of rank 2 endowed with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$.

Remark

The Kummer map now reads as

$$E(\mathbb{Q}) \otimes \mathbb{Z}_p \hookrightarrow H^1(\mathbb{Q}, T_p E)$$

and the image is contained in $\text{Sel}(\mathbb{Q}, T_p E)$.



Tate module

A new
framework
for
Kolyagin
systems

Introduction
and
motivation

Euler and
Kolyagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyagin
systems

Tate module

We will study the limit $T_p E = \varprojlim_k E[p^k]$, which is a free \mathbb{Z}_p -module of rank 2 endowed with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$.

Remark

The Kummer map now reads as

$$E(\mathbb{Q}) \otimes \mathbb{Z}_p \hookrightarrow H^1(\mathbb{Q}, T_p E)$$

and the image is contained in $\text{Sel}(\mathbb{Q}, T_p E)$.

Remark

We can recover the torsion points from the Tate module since

$$E[p^k] = T_p E / p^k$$



Tate module

A new
framework
for
Kolyagin
systems

Introduction
and
motivation

Euler and
Kolyagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyagin
systems

Tate module

We will study the limit $T_p E = \varprojlim_k E[p^k]$, which is a free \mathbb{Z}_p -module of rank 2 endowed with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$.

Remark

The Kummer map now reads as

$$E(\mathbb{Q}) \otimes \mathbb{Z}_p \hookrightarrow H^1(\mathbb{Q}, T_p E)$$

and the image is contained in $\text{Sel}(\mathbb{Q}, T_p E)$.

Remark

We can recover the torsion points from the Tate module since


$$E[p^k] = T_p E / p^k$$

Galois representations

We only need to assume that T is a free, finitely generated \mathbb{Z}_p -module, endowed with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$.



General picture





Euler systems

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Euler systems

An **Euler system** is a collection of classes

$$c_m \in H^1(\mathbb{Q}(\zeta_m), T)$$

satisfying **norm-compatible relations** as m changes.



Euler systems

An **Euler system** is a collection of classes

$$c_m \in H^1(\mathbb{Q}(\zeta_m), T)$$

satisfying **norm-compatible relations** as m changes.

Etymology

The norm-compatibility relations involve Euler factors.



Euler systems

An **Euler system** is a collection of classes

$$c_m \in H^1(\mathbb{Q}(\zeta_m), T)$$

satisfying **norm-compatible relations** as m changes.

Etymology

The norm-compatibility relations involve Euler factors.

Reciprocity laws

The classes c_m are related to special (algebraic) L -values.



Euler systems

An **Euler system** is a collection of classes

$$c_m \in H^1(\mathbb{Q}(\zeta_m), T)$$

satisfying **norm-compatible relations** as m changes.

Etymology

The norm-compatibility relations involve Euler factors.

Reciprocity laws


The classes c_m are related to special (algebraic) L -values.

Construction

In general, it is hard to construct Euler systems and an active research area. For the particular case of an elliptic curve, an Euler system was constructed by Kato.



General picture





Kolyvagin derivative

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Kolyvagin derivative

Fix $k \in \mathbb{N}$. Kolyvagin constructed a set of primes \mathcal{P} in which

$$\ell \equiv 1 \pmod{p^k} \quad \forall \ell \in \mathcal{P}$$

We denote by \mathcal{N} the set of square-free products of Kolyvagin primes $n = \ell_1 \cdots \ell_s$.



Kolyvagin derivative

Kolyvagin derivative

Fix $k \in \mathbb{N}$. Kolyvagin constructed a set of primes \mathcal{P} in which

$$\ell \equiv 1 \pmod{p^k} \quad \forall \ell \in \mathcal{P}$$

We denote by \mathcal{N} the set of square-free products of Kolyvagin primes $n = \ell_1 \cdots \ell_s$. For $n \in \mathcal{N}$, we can define a Kolyvagin derivative operator

$$D_n : H^1(\mathbb{Q}(\zeta_n), T) \rightarrow H^1(\mathbb{Q}, T/p^k)$$



Kolyvagin derivative

Fix $k \in \mathbb{N}$. Kolyvagin constructed a set of primes \mathcal{P} in which

$$\ell \equiv 1 \pmod{p^k} \quad \forall \ell \in \mathcal{P}$$

We denote by \mathcal{N} the set of square-free products of Kolyvagin primes $n = \ell_1 \cdots \ell_s$. For $n \in \mathcal{N}$, we can define a Kolyvagin derivative operator

$$D_n : H^1(\mathbb{Q}(\zeta_n), T) \rightarrow H^1(\mathbb{Q}, T/p^k)$$

Then the **Kolyvagin derivative class** is

$$\kappa_n := D_n c_n \in H^1(\mathbb{Q}, T/p^k)$$



Kolyvagin derivative

Fix $k \in \mathbb{N}$. Kolyvagin constructed a set of primes \mathcal{P} in which

$$\ell \equiv 1 \pmod{p^k} \quad \forall \ell \in \mathcal{P}$$

We denote by \mathcal{N} the set of square-free products of Kolyvagin primes $n = \ell_1 \cdots \ell_s$. For $n \in \mathcal{N}$, we can define a Kolyvagin derivative operator

$$D_n : H^1(\mathbb{Q}(\zeta_n), T) \rightarrow H^1(\mathbb{Q}, T/p^k)$$

Then the **Kolyvagin derivative class** is


$$\kappa_n := D_n c_n \in H^1(\mathbb{Q}, T/p^k)$$

Kolyvagin system

The collection of derivative classes $\{\kappa_n : n \in \mathcal{N}\}$ is a **Kolyvagin system**.



General picture





Kolyvagin systems

A **Kolyvagin system** is a collection of classes $\{\kappa_n : n \in \mathcal{N}\}$ such that

- $\kappa_n \in H^1(\mathbb{Q}, T/p^k)$
- κ_n is unramified at primes not dividing n or p and *interestingly* ramified at primes dividing n .
- There is a relation between κ_n and $\kappa_{n\ell}$ for all $n \in \mathcal{N}$ and $\ell \in \mathcal{P}$.



Kolyvagin systems

A **Kolyvagin system** is a collection of classes $\{\kappa_n : n \in \mathcal{N}\}$ such that

- $\kappa_n \in H^1(\mathbb{Q}, T/p^k)$
- κ_n is unramified at primes not dividing n or p and *interestingly* ramified at primes dividing n .
- There is a relation between κ_n and $\kappa_{n\ell}$ for all $n \in \mathcal{N}$ and $\ell \in \mathcal{P}$.

Remark

There are Kolyvagin systems for all $k \in \mathbb{N}$. However, we cannot define Kolyvagin systems on $H^1(\mathbb{Q}, T)$, since there is no prime $\ell \equiv 1 \pmod{p^k}$ for every k .



Assumptions

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

- T/pT is an absolutely irreducible Galois representation.



Assumptions

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

- T/pT is an absolutely irreducible Galois representation.
- The map $\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Aut}(T)$ is surjective.



Assumptions

A new
framework
for
Kolyagin
systems

Introduction
and
motivation

Euler and
Kolyagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyagin
systems

- T/pT is an absolutely irreducible Galois representation.
- The map $\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Aut}(T)$ is surjective.
- The Selmer structure is cartesian (technical assumption).



Dual Selmer group

- The dual Galois representation is $T^* = \text{Hom}(T, \mu_{p^\infty})$.



Dual Selmer group

- The dual Galois representation is $T^* = \text{Hom}(T, \mu_{p^\infty})$.
- For every Selmer group, there exists the concept of a dual Selmer group $\text{Sel}(\mathbb{Q}, T^*)$, defined in the cohomology of T^* via local Tate duality.



Dual Selmer group

- The dual Galois representation is $T^* = \text{Hom}(T, \mu_{p^\infty})$.
- For every Selmer group, there exists the concept of a dual Selmer group $\text{Sel}(\mathbb{Q}, T^*)$, defined in the cohomology of T^* via local Tate duality.
- Poitou-Tate duality relates the original Selmer group with the dual Selmer group.



Dual Selmer group

- The dual Galois representation is $T^* = \text{Hom}(T, \mu_{p^\infty})$.
- For every Selmer group, there exists the concept of a dual Selmer group $\text{Sel}(\mathbb{Q}, T^*)$, defined in the cohomology of T^* via local Tate duality.
- Poitou-Tate duality relates the original Selmer group with the dual Selmer group.

Core rank

- There exists an integer χ such that

$$\text{Sel}(\mathbb{Q}, T/p^k) \cong \text{Sel}(\mathbb{Q}, (T/p^k)^*) \oplus (\mathbb{Z}/p^k)^\chi$$

- χ is known as the **core rank** of the Selmer group.



Core rank and Kolyvagin systems

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Theorem (Mazur-Rubin, 2004)

- If $\chi = 0$, then $\text{KS}(\mathcal{T}) = 0$.

There are no Kolyvagin system to control the Selmer group. We will see a possible solution later in the talk.



Core rank and Kolyvagin systems

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Theorem (Mazur-Rubin, 2004)

- If $\chi = 0$, then $\text{KS}(T) = 0$.

There are no Kolyvagin system to control the Selmer group. We will see a possible solution later in the talk.

- If $\chi = 1$, then $\text{KS}(T/p^k) \cong \mathbb{Z}/p^k$.



Core rank and Kolyvagin systems

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Theorem (Mazur-Rubin, 2004)

- If $\chi = 0$, then $\text{KS}(T) = 0$.

There are no Kolyvagin system to control the Selmer group. We will see a possible solution later in the talk.


- If $\chi = 1$, then $\text{KS}(T/p^k) \cong \mathbb{Z}/p^k$.

- If $\chi > 1$, then $\text{KS}(T/p^k)$ is too large.

In order to compute the Selmer group, [Mazur-Rubin, 2016] and [Burns-Sakamoto-Sano, 2025] modified the definition of Kolyvagin system in (biduals of) exterior powers of the Selmer groups.



General picture





Fitting ideals

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .



Fitting ideals

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fact: Fitting ideals are well defined.



Fitting ideals

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fact: Fitting ideals are well defined.

Example

Consider $R = \mathbb{Z}_p$ and $M = \mathbb{Z}_p \times \mathbb{Z}_p/p^3 \times \mathbb{Z}_p/p^2$.



Fitting ideals

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fact: Fitting ideals are well defined.

Example

Consider $R = \mathbb{Z}_p$ and $M = \mathbb{Z}_p \times \mathbb{Z}_p/p^3 \times \mathbb{Z}_p/p^2$. A resolution is given by

$$(\mathbb{Z}_p)^3 \xrightarrow{\mu} (\mathbb{Z}_p)^3 \xrightarrow{\varepsilon} M \longrightarrow 0$$

Here ε is the natural map and μ is given by the matrix $A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & p^3 & 0 \\ 0 & 0 & p^2 \end{pmatrix}$



Fitting ideals

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fact: Fitting ideals are well defined.

Example

Consider $R = \mathbb{Z}_p$ and $M = \mathbb{Z}_p \times \mathbb{Z}_p/p^3 \times \mathbb{Z}_p/p^2$. A resolution is given by

$$(\mathbb{Z}_p)^3 \xrightarrow{\mu} (\mathbb{Z}_p)^3 \xrightarrow{\varepsilon} M \longrightarrow 0$$

Here ε is the natural map and μ is given by the matrix $A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & p^3 & 0 \\ 0 & 0 & p^2 \end{pmatrix}$

$\text{Fitt}_0(M) = (0),$	$\text{Fitt}_1(M) = (p^5),$
$\text{Fitt}_2(M) = (p^2) + (p^3) = (p^2)$	$\text{Fitt}_i(M) = (1) \forall i \geq 3$



Fitting ideals over Discrete Valuation Rings

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.



Fitting ideals over Discrete Valuation Rings

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

$$\blacksquare i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$$



Fitting ideals over Discrete Valuation Rings

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$



Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$
- $i \geq r+s \Rightarrow \text{Fitt}_i(M) = (1).$



Fitting ideals over Discrete Valuation Rings

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \dots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j}(M) = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$
- $i \geq r+s \Rightarrow \text{Fitt}_i(M) = (1)$.

Corollary

The Fitting ideals determine M up to isomorphism:



Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \dots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$
- $i \geq r+s \Rightarrow \text{Fitt}_i(M) = (1).$

Corollary

The Fitting ideals determine M up to isomorphism:

- r is the minimum i such that $\text{Fitt}_i(M) \neq 0$.



Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$
- $i \geq r+s \Rightarrow \text{Fitt}_i(M) = (1).$

Corollary

The Fitting ideals determine M up to isomorphism:

- r is the minimum i such that $\text{Fitt}_i(M) \neq 0$.
- For $i \geq 0$, $\alpha_i = \text{Fitt}_{r+i+1}(M) \text{Fitt}_{r+i}(M)^{-1}$.



Fitting ideals over the Iwasawa algebra

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Iwasawa algebra

The Iwasawa algebra can be represented as

$$\Lambda = \mathbb{Z}_p[[X]]$$



Fitting ideals over the Iwasawa algebra

Iwasawa algebra

The Iwasawa algebra can be represented as

$$\Lambda = \mathbb{Z}_p[[X]]$$

Structure theorem

Every finitely generated Λ -module is pseudo-isomorphic to

$$M \approx \Lambda^r \times \prod \frac{\Lambda}{(p)^{\alpha_i}} \times \prod \frac{\Lambda}{(f_j)^{\beta_j}}$$

where f_j are irreducible distinguished polynomials.



Fitting ideals over the Iwasawa algebra

Iwasawa algebra

The Iwasawa algebra can be represented as

$$\Lambda = \mathbb{Z}_p[[X]]$$

Structure theorem

Every finitely generated Λ -module is pseudo-isomorphic to

$$M \approx \Lambda^r \times \prod \frac{\Lambda}{(p)^{\alpha_i}} \times \prod \frac{\Lambda}{(f_j)^{\beta_j}}$$

where f_j are irreducible distinguished polynomials.

Fitting ideals

Fitting ideals can recover the structure of a finitely generated Iwasawa module up to pseudo-isomorphism.



Indices and theta-ideals

Index of an element

Let M be an R -module and let $a \in M$. Denote by $M^+ = \text{Hom}(M, R)$ the dual of M . There is a canonical map

$$\Phi : M \rightarrow M^{++} : x \mapsto (\varphi \mapsto \varphi(x))$$

Note that $\Phi(a) \in \text{Hom}(M^+, R)$. Define the index of a as

$$\text{ind}(a, M) = \text{Im}(\Phi(a))$$



Indices and theta-ideals

Index of an element

Let M be an R -module and let $a \in M$. Denote by $M^+ = \text{Hom}(M, R)$ the dual of M . There is a canonical map

$$\Phi : M \rightarrow M^{++} : x \mapsto (\varphi \mapsto \varphi(x))$$

Note that $\Phi(a) \in \text{Hom}(M^+, R)$. Define the index of a as

$$\text{ind}(a, M) = \text{Im}(\Phi(a))$$

Remark

If R is a principal local ring, with π being a generator of the maximal ideal. Then

$$\text{ind}(a, M) = \max\{j \in \mathbb{N} : j \in \pi^j M\}$$



Indices and theta-ideals

Index of an element

Let M be an R -module and let $a \in M$. Denote by $M^+ = \text{Hom}(M, R)$ the dual of M . There is a canonical map

$$\Phi : M \rightarrow M^{++} : x \mapsto (\varphi \mapsto \varphi(x))$$

Note that $\Phi(a) \in \text{Hom}(M^+, R)$. Define the index of a as

$$\text{ind}(a, M) = \text{Im}(\Phi(a))$$

Remark

If R is a principal local ring, with π being a generator of the maximal ideal. Then

$$\text{ind}(a, M) = \max\{j \in \mathbb{N} : j \in \pi^j M\}$$


Theta ideals

We denote by \mathcal{N}_i the set of square-free products of exactly i Kolyvagin primes. We define the i^{th} theta ideal of a Kolyvagin system as

$$\Theta_i(\kappa) := \sum_{n \in \mathcal{N}_i} (\kappa_n, H^1(\mathbb{Q}, T))$$



General picture





Core rank 1

Recall

When $\chi = 1$, the module of Kolyvagin system is

$$\mathrm{KS}(T/p^k) \cong \mathbb{Z}/p^k$$



Core rank 1

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Recall

When $\chi = 1$, the module of Kolyvagin system is

$$\mathrm{KS}(T/p^k) \cong \mathbb{Z}/p^k$$

Primitive Kolyvagin systems

We call a Kolyvagin system **primitive** if it generates $\mathrm{KS}(T/p^k)$.



Core rank 1

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Recall

When $\chi = 1$, the module of Kolyvagin system is

$$\mathrm{KS}(T/p^k) \cong \mathbb{Z}/p^k$$

Primitive Kolyvagin systems

We call a Kolyvagin system **primitive** if it generates $\mathrm{KS}(T/p^k)$.


Theorem (Mazur-Rubin, 2004)

When $\chi = 1$ and κ is a primitive Kolyvagin system

$$\Theta_i(\kappa) = \mathrm{Fitt}_i\left(\mathrm{Sel}\left(\mathbb{Q}, (T/p^k)^*\right)\right) = \mathrm{Fitt}_{i+1}\left(\mathrm{Sel}(\mathbb{Q}, T/p^k)\right)$$



General picture





Core rank 0: non-self-dual Galois representations

- We need to relax the local condition at one prime ℓ in order to obtain a rank one Selmer group.



Core rank 0: non-self-dual Galois representations

- We need to relax the local condition at one prime ℓ in order to obtain a rank one Selmer group.
- Fix a primitive Kolyvagin system $\kappa = (\kappa_n)_{n \in \mathcal{N}}$.
- We now consider the cohomology classes

$$\delta_n := \text{loc}_\ell(\kappa_n)$$



Core rank 0: non-self-dual Galois representations

- A new framework for Kolyvagin systems
- Introduction and motivation
- Euler and Kolyvagin systems
- Fitting ideals
- Core rank one
- Core rank 0**
- Ultra Kolyvagin systems

- We need to relax the local condition at one prime ℓ in order to obtain a rank one Selmer group.
- Fix a primitive Kolyvagin system $\kappa = (\kappa_n)_{n \in \mathcal{N}}$.
- We now consider the cohomology classes

$$\delta_n := \text{loc}_\ell(\kappa_n)$$

- We define the rank 0 theta-ideals of κ as

$$\Theta_i^{(0)}(\kappa) := \sum_{n \in \mathcal{N}_i(\mathcal{P})} \text{ind}(\delta_n)$$



Core rank 0: non-self-dual Galois representations

- We need to relax the local condition at one prime ℓ in order to obtain a rank one Selmer group.
- Fix a primitive Kolyvagin system $\kappa = (\kappa_n)_{n \in \mathcal{N}}$.
- We now consider the cohomology classes

$$\delta_n := \text{loc}_\ell(\kappa_n)$$

- We define the rank 0 theta-ideals of κ as

$$\Theta_i^{(0)}(\kappa) := \sum_{n \in \mathcal{N}_i(\mathcal{P})} \text{ind}(\delta_n)$$

Theorem (A., 2025)

If T is **not** a residually self-dual Galois representation, then

$$\Theta_i^{(0)}(\kappa) = \text{Fitt}_i(\text{Sel}(\mathbb{Q}, T/p^k))$$



Core rank 0: self-dual Galois representations

Theorem (A., 2025)

For all i , we have

$$\Theta_i^{(0)}(\kappa) \subset \text{Fitt}_i \left(\text{Sel}(\mathbb{Q}, T/p^k) \right)$$



Core rank 0: self-dual Galois representations

Theorem (A., 2025)

For all i , we have

$$\Theta_i^{(0)}(\kappa) \subset \text{Fitt}_i \left(\text{Sel}(\mathbb{Q}, T/p^k) \right)$$

The equality for some index i holds if any of the following is true:



Core rank 0: self-dual Galois representations

Theorem (A., 2025)

For all i , we have

$$\Theta_i^{(0)}(\kappa) \subset \text{Fitt}_i(\text{Sel}(\mathbb{Q}, T/p^k))$$

The equality for some index i holds if any of the following is true:

- $\Theta_{i-1}^{(0)}(\kappa) \subsetneq \text{Fitt}_{i-1}(\text{Sel}(\mathbb{Q}, T/p^k))$.



Core rank 0: self-dual Galois representations

Theorem (A., 2025)

For all i , we have

$$\Theta_i^{(0)}(\kappa) \subset \text{Fitt}_i \left(\text{Sel}(\mathbb{Q}, T/p^k) \right)$$

The equality for some index i holds if any of the following is true:

- $\Theta_{i-1}^{(0)}(\kappa) \subsetneq \text{Fitt}_{i-1} \left(\text{Sel}(\mathbb{Q}, T/p^k) \right)$.
- $\Theta_{i-1}^{(0)}(\kappa) \left(\text{Sel}(\mathbb{Q}, T/p^k) \right) = 0$.



Core rank 0: self-dual Galois representations

Theorem (A., 2025)

For all i , we have

$$\Theta_i^{(0)}(\kappa) \subset \text{Fitt}_i(\text{Sel}(\mathbb{Q}, T/p^k))$$

The equality for some index i holds if any of the following is true:

- $\Theta_{i-1}^{(0)}(\kappa) \subsetneq \text{Fitt}_{i-1}(\text{Sel}(\mathbb{Q}, T/p^k))$.
- $\Theta_{i-1}^{(0)}(\kappa) (\text{Sel}(\mathbb{Q}, T/p^k)) = 0$.

Conjecture

If T is a self-dual Galois representation, either

- $\Theta_i^{(0)}(\kappa)$ for all even i .
- $\Theta_i^{(0)}(\kappa)$ for all odd i .



Example: elliptic Curve

Limit of Selmer groups

Note that $T_p E^* = E[p^\infty]$ and that

$$\mathrm{Sel}(\mathbb{Q}, E[p^\infty]) = \varinjlim_k \mathrm{Sel}(\mathbb{Q}, E[p^k])$$



Example: elliptic Curve

Limit of Selmer groups

Note that $T_p E^* = E[p^\infty]$ and that

$$\mathrm{Sel}(\mathbb{Q}, E[p^\infty]) = \varinjlim_k \mathrm{Sel}(\mathbb{Q}, E[p^k])$$

Kurihara numbers

Kato's Euler system produces a Kolyvagin systems and $\mathrm{ind}(\delta_n) = \mathrm{ind}(\tilde{\delta}_n)$, where $\tilde{\delta}_n$ are known as the **Kurihara numbers** and are defined by the formula

$$\tilde{\delta}_n := \sum_{a \in (\mathbb{Z}/n)^\times} \left(\left[\frac{a}{n} \right]^+ + \left[\frac{a}{n} \right]^- \right) \prod_{\ell|n} \log_{\eta_\ell}(a) \in \mathbb{Z}_p$$



Example: elliptic Curve

Limit of Selmer groups

Note that $T_p E^* = E[p^\infty]$ and that

$$\mathrm{Sel}(\mathbb{Q}, E[p^\infty]) = \varinjlim_k \mathrm{Sel}(\mathbb{Q}, E[p^k])$$

Kurihara numbers

Kato's Euler system produces a Kolyvagin systems and $\mathrm{ind}(\delta_n) = \mathrm{ind}(\tilde{\delta}_n)$, where $\tilde{\delta}_n$ are known as the **Kurihara numbers** and are defined by the formula

$$\tilde{\delta}_n := \sum_{a \in (\mathbb{Z}/n)^\times} \left(\left[\frac{a}{n} \right]^+ + \left[\frac{a}{n} \right]^- \right) \prod_{\ell|n} \log_{\eta_\ell}(a) \in \mathbb{Z}_p$$

By the functional equation of the L -function, the conjecture holds true in this case.



Selmer groups with coefficients in the Iwasawa algebra

A new
framework
for
Kolyagin
systems

Introduction
and
motivation

Euler and
Kolyagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyagin
systems

\mathbb{Z}_p -extensions

In Iwasawa theory, we are interested in the Selmer group over a \mathbb{Z}_p -extension. By Shapiro's lemma, it is equivalent to study Galois representations \mathbf{T} with coefficients over the Iwasawa algebra Λ .



Selmer groups with coefficients in the Iwasawa algebra

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

\mathbb{Z}_p -extensions

In Iwasawa theory, we are interested in the Selmer group over a \mathbb{Z}_p -extension. By Shapiro's lemma, it is equivalent to study Galois representations \mathbf{T} with coefficients over the Iwasawa algebra Λ .

Selmer groups over the Iwasawa algebra

If \mathbf{T} is a (finitely generated, free) Λ -module, it is more difficult to study the Selmer group as a limit of Selmer groups with finite coefficients, since the representation theory over the finite quotients of Λ is more complicated.



Selmer groups with coefficients in the Iwasawa algebra

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

\mathbb{Z}_p -extensions

In Iwasawa theory, we are interested in the Selmer group over a \mathbb{Z}_p -extension. By Shapiro's lemma, it is equivalent to study Galois representations \mathbf{T} with coefficients over the Iwasawa algebra Λ .

Selmer groups over the Iwasawa algebra

If \mathbf{T} is a (finitely generated, free) Λ -module, it is more difficult to study the Selmer group as a limit of Selmer groups with finite coefficients, since the representation theory over the finite quotients of Λ is more complicated.

Idea

Use [patched cohomology](#) to overcome the lack of Kolyvagin primes and generalise the notion of Kolyvagin system to infinite coefficient rings.



Filters and ultrafilters

A **filter** of the natural numbers is a subset $\mathcal{U} \subset \mathbb{P}(\mathbb{N})$ such that

- $S \in \mathcal{U}, S \subset T \Rightarrow T \in \mathcal{U}$
- $S_1, S_2 \in \mathcal{U} \Rightarrow S_1 \cap S_2 \in \mathcal{U}$



Filters and ultrafilters

A **filter** of the natural numbers is a subset $\mathcal{U} \subset \mathbb{P}(\mathbb{N})$ such that

- $S \in \mathcal{U}, S \subset T \Rightarrow T \in \mathcal{U}$
- $S_1, S_2 \in \mathcal{U} \Rightarrow S_1 \cap S_2 \in \mathcal{U}$

\mathcal{U} is said to be an **ultrafilter** if

- For every $S \subset \mathbb{N}$, either $S \in \mathcal{U}$ or $(\mathbb{N} \setminus S) \in \mathcal{U}$.



Filters and ultrafilters

A **filter** of the natural numbers is a subset $\mathcal{U} \subset \mathbb{P}(\mathbb{N})$ such that

- $S \in \mathcal{U}, S \subset T \Rightarrow T \in \mathcal{U}$
- $S_1, S_2 \in \mathcal{U} \Rightarrow S_1 \cap S_2 \in \mathcal{U}$

\mathcal{U} is said to be an **ultrafilter** if

- For every $S \subset \mathbb{N}$, either $S \in \mathcal{U}$ or $(\mathbb{N} \setminus S) \in \mathcal{U}$.

Principal ultrafilters

Assume there is a finite set $S \in \mathcal{U}$. Then there is an element $a \in S$ such that

$$\mathcal{U} = \{T \subset \mathbb{N} : a \in T\}$$

In this case, \mathcal{U} is said to be a **principal ultrafilter**.



Filters and ultrafilters

A **filter** of the natural numbers is a subset $\mathcal{U} \subset \mathbb{P}(\mathbb{N})$ such that

- $S \in \mathcal{U}, S \subset T \Rightarrow T \in \mathcal{U}$
- $S_1, S_2 \in \mathcal{U} \Rightarrow S_1 \cap S_2 \in \mathcal{U}$

\mathcal{U} is said to be an **ultrafilter** if

- For every $S \subset \mathbb{N}$, either $S \in \mathcal{U}$ or $(\mathbb{N} \setminus S) \in \mathcal{U}$.

Principal ultrafilters

Assume there is a finite set $S \in \mathcal{U}$. Then there is an element $a \in S$ such that

$$\mathcal{U} = \{T \subset \mathbb{N} : a \in T\}$$

In this case, \mathcal{U} is said to be a **principal ultrafilter**.

Fix a non-principal ultrafilter \mathcal{U} .



Patching

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Patching (Sweeting 2021)

Let $(M_n)_{n \in \mathbb{N}}$ be a sequence of sets/groups/rings, we define the **patching** via the ultrafilter \mathcal{U} as

$$\mathcal{U}(M_n) = \prod_{n \in \mathcal{N}} M_n / \sim$$

where two sequences $(\alpha_n)_{n \in \mathbb{N}}$ and $(\beta_n)_{n \in \mathbb{N}}$ are said to be equivalent if $\alpha_n = \beta_n$ for \mathcal{U} -many n .



Patching

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Patching (Sweeting 2021)

Let $(M_n)_{n \in \mathbb{N}}$ be a sequence of sets/groups/rings, we define the **patching** via the ultrafilter \mathcal{U} as

$$\mathcal{U}(M_n) = \prod_{n \in \mathcal{N}} M_n / \sim$$

where two sequences $(\alpha_n)_{n \in \mathbb{N}}$ and $(\beta_n)_{n \in \mathbb{N}}$ are said to be equivalent if $\alpha_n = \beta_n$ for \mathcal{U} -many n .

Proposition

\mathcal{U} is an exact functor.



Patching (Sweeting 2021)

Let $(M_n)_{n \in \mathbb{N}}$ be a sequence of sets/groups/rings, we define the **patching** via the ultrafilter \mathcal{U} as

$$\mathcal{U}(M_n) = \prod_{n \in \mathcal{N}} M_n / \sim$$

where two sequences $(\alpha_n)_{n \in \mathbb{N}}$ and $(\beta_n)_{n \in \mathbb{N}}$ are said to be equivalent if $\alpha_n = \beta_n$ for \mathcal{U} -many n .

Proposition

\mathcal{U} is an exact functor.

Constant patching of finite groups

If M is a finite group and $M_n = M$ for all $n \in \mathbb{N}$, then

$$\mathcal{U}(M_n) = M$$



Ultraprimers

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Ultraprimers

An **ultraprime** is an element of $\mathcal{U}(\{\text{primes}\})$, so it can be represented by a sequence

$$u = (\ell_1, \dots, \ell_n, \dots)$$



Ultraprimeness

An **ultraprime** is an element of $\mathcal{U}(\{\text{primes}\})$, so it can be represented by a sequence

$$u = (\ell_1, \dots, \ell_n, \dots)$$

Kolyagin ultraprimes

An ultraprime $u = (\ell_i)_{i \in \mathbb{N}}$ is said to be a **Kolyagin ultraprime** if, for every finite quotient of Λ , ℓ_i is a Kolyagin prime for \mathcal{U} -many i .



Patched cohomology (Sweeting, 2021)

Finite coefficients

Assume T is a finite group endowed with an action a sequence of groups $G = (G_n)$. We define the patched cohomology as

$$\mathbf{H}^1(G, T) = \mathcal{U}(H^1(G, T))$$



Patched cohomology (Sweeting, 2021)

Finite coefficients

Assume T is a finite group endowed with an action a sequence of groups $G = (G_n)$. We define the patched cohomology as

$$\mathbf{H}^1(G, T) = \mathcal{U}(H^1(G, T))$$

Profinite coefficients

Assume T is profinite, the patched cohomology is defined as

$$\mathbf{H}^1(G, T) = \varprojlim_{T \twoheadrightarrow T'} \mathbf{H}^1(G, T')$$

where T' runs through the finite quotients of T .



Patched cohomology (Sweeting, 2021)

Finite coefficients

Assume T is a finite group endowed with an action a sequence of groups $G = (G_n)$. We define the patched cohomology as

$$\mathbf{H}^1(G, T) = \mathcal{U}(H^1(G, T))$$

Profinite coefficients

Assume T is profinite, the patched cohomology is defined as

$$\mathbf{H}^1(G, T) = \varprojlim_{T \twoheadrightarrow T'} \mathbf{H}^1(G, T')$$

where T' runs through the finite quotients of T .

Ind-finite coefficients

If T is ind-finite, then

$$\mathbf{H}^1(G, T) = \varinjlim_{T' \hookrightarrow T} \mathbf{H}^1(G, T')$$

where T' runs through the finite submodules of T .



Local and global patched cohomology

Local patched cohomology

Let T be a Galois representation, i.e., T is endowed with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Let $u = (\ell_n)$ be an ultraprime. Since $\text{Gal}(\overline{\mathbb{Q}_{\ell_i}}/\mathbb{Q}_{\ell_i})$ is contained in $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, it also acts on T .

Then the patched local cohomology

$$H^1(\mathbb{Q}_u, T)$$

is the patching of the sequence of local Galois groups with coefficients in T . In particular, when T is finite,

$$H^1(\mathbb{Q}_u, T) = \mathcal{U}(H^1(\text{Gal}(\overline{\mathbb{Q}_{\ell_i}}/\mathbb{Q}_{\ell_i}), T))$$



Local and global patched cohomology

Local patched cohomology

Let T be a Galois representation, i.e., T is endowed with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Let $u = (\ell_n)$ be an ultraprime. Since $\text{Gal}(\overline{\mathbb{Q}_{\ell_i}}/\mathbb{Q}_{\ell_i})$ is contained in $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, it also acts on T .

Then the patched local cohomology

$$H^1(\mathbb{Q}_u, T)$$

is the patching of the sequence of local Galois groups with coefficients in T . In particular, when T is finite,

$$H^1(\mathbb{Q}_u, T) = \mathcal{U}(H^1(\text{Gal}(\overline{\mathbb{Q}_{\ell_i}}/\mathbb{Q}_{\ell_i}), T))$$

Global patched cohomology

There is also a notion of patched global cohomology unramified outside the square-free (formal) product of ultraprimes $n = u_1 \cdots u_s$, denoted by

$$H^1(\mathbb{Q}_{\Sigma_n}/\mathbb{Q}, T)$$



Local and global patched cohomology

Local patched cohomology

Let T be a Galois representation, i.e., T is endowed with an action of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Let $\mathfrak{u} = (\ell_n)$ be an ultraprime. Since $\text{Gal}(\overline{\mathbb{Q}_{\ell_i}}/\mathbb{Q}_{\ell_i})$ is contained in $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, it also acts on T .

Then the patched local cohomology

$$H^1(\mathbb{Q}_{\mathfrak{u}}, T)$$

is the patching of the sequence of local Galois groups with coefficients in T . In particular, when T is finite,

$$H^1(\mathbb{Q}_{\mathfrak{u}}, T) = \mathcal{U}(H^1(\text{Gal}(\overline{\mathbb{Q}_{\ell_i}}/\mathbb{Q}_{\ell_i}), T))$$

Global patched cohomology

There is also a notion of patched global cohomology unramified outside the square-free (formal) product of ultraprimes $n = \mathfrak{u}_1 \cdots \mathfrak{u}_s$, denoted by

$$H^1(\mathbb{Q}_{\Sigma_n}/\mathbb{Q}, T)$$

Selmer groups

We can extend the notion of Selmer groups to this setting and define local conditions on the local cohomology groups. We recover the classical Selmer groups when



Ultra-Kolyvagin systems

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Notation

- \mathcal{P} : set of Kolyvagin ultraprimes.
- $\mathcal{N}(\mathcal{P})$: set of square-free products of Kolyvagin ultraprimesprimes
- $\mathcal{N}_i(\mathcal{P})$: set of square-free products of i Kolyvagin ultraprimes.



Ultra-Kolyvagin systems

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Notation

- \mathcal{P} : set of Kolyvagin ultraprimes.
- $\mathcal{N}(\mathcal{P})$: set of square-free products of Kolyvagin ultraprimes
- $\mathcal{N}_i(\mathcal{P})$: set of square-free products of i Kolyvagin ultraprimes.

Ultra-Kolyvagin systems

A Kolyvagin system is a collection $\{\kappa_n : n \in \mathcal{N}\}$ such that

- $\kappa_n \in \text{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, \mathbf{T}) \subset \mathbf{H}^1(\mathbb{Q}_{\Sigma_n}/\mathbb{Q}, \mathbf{T})$
- κ_n and $\kappa_{n\ell}$ satisfy a Kolyvagin relation for all $n \in \mathcal{N}$ and



Ultra-Kolyvagin systems

A new
framework
for
Kolyvagin
systems

Introduction
and
motivation

Euler and
Kolyvagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyvagin
systems

Notation

- \mathcal{P} : set of Kolyvagin ultraprimes.
- $\mathcal{N}(\mathcal{P})$: set of square-free products of Kolyvagin ultraprimes
- $\mathcal{N}_i(\mathcal{P})$: set of square-free products of i Kolyvagin ultraprimes.

Ultra-Kolyvagin systems

A Kolyvagin system is a collection $\{\kappa_n : n \in \mathcal{N}\}$ such that

- $\kappa_n \in \text{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, \mathbf{T}) \subset \mathbf{H}^1(\mathbb{Q}_{\Sigma_n}/\mathbb{Q}, \mathbf{T})$
- κ_n and $\kappa_{n\ell}$ satisfy a Kolyvagin relation for all $n \in \mathcal{N}$ and

Theta ideals

$$\Theta_i(\kappa) := \sum_{n \in \mathcal{N}_i(\mathcal{P})} \text{ind}(\kappa_n)$$



Selmer group over cyclotomic \mathbb{Z}_p -extensions

A new
framework
for
Kolyagin
systems

Introduction
and
motivation

Euler and
Kolyagin
systems

Fitting
ideals

Core rank
one

Core rank
0

Ultra
Kolyagin
systems

Assumptions

- T is residually irreducible.
- $\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Aut}(T)$ is surjective.
- The Selmer structure is cartesian.
- $\frac{H^1(\mathbb{Q}_\mu, T \otimes \Lambda)}{H^1_{\mathcal{F}}(\mathbb{Q}_\mu, T \otimes \Lambda)}$ is Λ -torsion free



Selmer group over cyclotomic \mathbb{Z}_p -extensions

Assumptions

- T is residually irreducible.
- $\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Aut}(T)$ is surjective.
- The Selmer structure is cartesian.
- $\frac{H^1(\mathbb{Q}_\mu, T \otimes \Lambda)}{H^1_{\mathcal{F}}(\mathbb{Q}_\mu, T \otimes \Lambda)}$ is Λ -torsion free

Theorem (A., in progress)

The module of ultra-Kolyagin systems $\text{KS}(T \otimes \Lambda)$ is free of rank one over Λ .
Moreover, if we choose a primitive ultra-Kolyagin system κ , then

$$\Theta_i(\kappa) =_{f.i.} \text{Fitt}_\Lambda^i(\text{Sel}(\mathbb{Q}, \mathbf{T}^*)^\vee)$$



Selmer group over cyclotomic \mathbb{Z}_p -extensions

Assumptions

- T is residually irreducible.
- $\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{Aut}(T)$ is surjective.
- The Selmer structure is cartesian.
- $\frac{H^1(\mathbb{Q}_\mu, T \otimes \Lambda)}{H^1_{\mathcal{F}}(\mathbb{Q}_\mu, T \otimes \Lambda)}$ is Λ -torsion free

Theorem (A., in progress)

The module of ultra-Kolyagin systems $\text{KS}(T \otimes \Lambda)$ is free of rank one over Λ . Moreover, if we choose a primitive ultra-Kolyagin system κ , then

$$\Theta_i(\kappa) =_{f.i.} \text{Fitt}_\Lambda^i(\text{Sel}(\mathbb{Q}, \mathbf{T}^*)^\vee)$$

Corollary

The ideals $\Theta_i(\kappa)$ determine the the structure of the Selmer group up to pseudo-isomorphism.

