

Kolyvagin systems and Fitting ideals of Selmer group of rank 0

Alberto Angurel Andrés

University of Nottingham

30/09/2025

Kolyvagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems


Structure
of Selmer
groups

Euler
system

Elliptic
Curves


Proofs

General picture




For today

- We are going to focus on



For today


- We are going to focus on



- We cannot apply the theory of Kolyvagin systems directly, because
 - The classical Selmer group is self-dual, so its core rank is zero.
 - There are no non-zero Kolyvagin systems for this Selmer group.

For today


- We are going to focus on



- We cannot apply the theory of Kolyvagin systems directly, because
 - The classical Selmer group is self-dual, so its core rank is zero.
 - There are no non-zero Kolyvagin systems for this Selmer group.
- The general theory of Kolyvagin systems only describes the structure of the Selmer group *restricted at p* .

For today

- We are going to focus on



- We cannot apply the theory of Kolyvagin systems directly, because
 - The classical Selmer group is self-dual, so its core rank is zero.
 - There are no non-zero Kolyvagin systems for this Selmer group.
- The general theory of Kolyvagin systems only describes the structure of the Selmer group *restricted at p* .
- We extend this theory to Selmer groups of rank zero by considering Kolyvagin systems over an auxiliary Selmer structure.

Setting and assumptions

- (H0) Let $p \geq 5$ and let \mathbf{T} be free \mathbb{Z}_p -module of finite rank endowed with a continuous action of $G_{\mathbb{Q}}$, ramifying only at a finite amount of primes.
- (H1) $\rho : G_{\mathbb{Q}} \rightarrow \text{Aut}(\mathbf{T})$ is surjective.
- (H2) (will appear later)

- Selmer groups are formed by the elements of the global cohomology groups $H^1(\mathbb{Q}, \mathbf{T})$ that satisfy *local conditions*.

Selmer pre-structures

- Selmer groups are formed by the elements of the global cohomology groups $H^1(\mathbb{Q}, \mathbf{T})$ that satisfy *local conditions*.
- What is a local condition? A **local condition** for a prime ℓ is a choice of a subgroup

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T}) \subset H^1(\mathbb{Q}_{\ell}, \mathbf{T})$$

Selmer pre-structures

- Selmer groups are formed by the elements of the global cohomology groups $H^1(\mathbb{Q}, \mathbf{T})$ that satisfy *local conditions*.
- What is a local condition? A **local condition** for a prime ℓ is a choice of a subgroup

$$H^1_{\mathcal{F}}(\mathbb{Q}_{\ell}, \mathbf{T}) \subset H^1(\mathbb{Q}_{\ell}, \mathbf{T})$$

Definition (Selmer pre-structure)

A **Selmer pre-structure** \mathcal{F} is a choice of a local condition for every prime (including the archimedean one).

- Selmer groups are formed by the elements of the global cohomology groups $H^1(\mathbb{Q}, \mathbf{T})$ that satisfy *local conditions*.
- What is a local condition? A **local condition** for a prime ℓ is a choice of a subgroup

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T}) \subset H^1(\mathbb{Q}_{\ell}, \mathbf{T})$$

Definition (Selmer pre-structure)

A **Selmer pre-structure** \mathcal{F} is a choice of a local condition for every prime (including the archimedean one).

Definition (Selmer group)

The **Selmer group** for \mathcal{F} is defined as

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T}) := \ker \left(H^1(\mathbb{Q}, \mathbf{T}) \rightarrow \bigoplus_{\ell} \frac{H^1(\mathbb{Q}_{\ell}, \mathbf{T})}{H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T})} \right)$$

Definition (finite cohomology)

$$H_f^1(\mathbb{Q}_\ell, \mathbf{T}) := \ker (H^1(\mathbb{Q}, \mathbf{T}) \rightarrow H^1(I_\ell, \mathbf{T} \otimes \mathbb{Q}_p))$$

Definition (finite cohomology)

$$H_f^1(\mathbb{Q}_\ell, \mathbf{T}) := \ker (H^1(\mathbb{Q}, \mathbf{T}) \rightarrow H^1(I_\ell, \mathbf{T} \otimes \mathbb{Q}_p))$$

Definition (Selmer structure)

A **Selmer structure** is a Selmer pre-structure such that there is a finite set of primes Σ such that

$$H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T}) = H_f^1(\mathbb{Q}_\ell, \mathbf{T}) \quad \forall \ell \notin \Sigma$$

Definition (finite cohomology)

$$H_f^1(\mathbb{Q}_\ell, \mathbf{T}) := \ker (H^1(\mathbb{Q}, \mathbf{T}) \rightarrow H^1(I_\ell, \mathbf{T} \otimes \mathbb{Q}_p))$$

Definition (Selmer structure)

A **Selmer structure** is a Selmer pre-structure such that there is a finite set of primes Σ such that

$$H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T}) = H_f^1(\mathbb{Q}_\ell, \mathbf{T}) \quad \forall \ell \notin \Sigma$$

Proposition (Selmer groups)

If \mathbb{Q}_Σ denotes the maximal extension of \mathbb{Q} unramified outside Σ , we have that

$$\text{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T}) = \ker \left(H^1(\mathbb{Q}_\Sigma/\mathbb{Q}, \mathbf{T}) \rightarrow \prod_{\ell \in \Sigma} \frac{H^1(\mathbb{Q}_\ell, \mathbf{T})}{H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T})} \right)$$

Definition (finite cohomology)

$$H_f^1(\mathbb{Q}_\ell, \mathbf{T}) := \ker (H^1(\mathbb{Q}, \mathbf{T}) \rightarrow H^1(I_\ell, \mathbf{T} \otimes \mathbb{Q}_p))$$

Definition (Selmer structure)

A **Selmer structure** is a Selmer pre-structure such that there is a finite set of primes Σ such that

$$H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T}) = H_f^1(\mathbb{Q}_\ell, \mathbf{T}) \quad \forall \ell \notin \Sigma$$

Proposition (Selmer groups)

If \mathbb{Q}_Σ denotes the maximal extension of \mathbb{Q} unramified outside Σ , we have that

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T}) = \ker \left(H^1(\mathbb{Q}_\Sigma/\mathbb{Q}, \mathbf{T}) \rightarrow \prod_{\ell \in \Sigma} \frac{H^1(\mathbb{Q}_\ell, \mathbf{T})}{H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T})} \right)$$

Corollary

$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T}) \subset H^1(\mathbb{Q}_\Sigma/\mathbb{Q}, \mathbf{T})$ is a finitely generated \mathbb{Z}_p -module.

Fitting ideals

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fitting ideals

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fact: Fitting ideals are well defined.

Fitting ideals

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fact: Fitting ideals are well defined.

Example

Consider $R = \mathbb{Z}_p$ and $M = \mathbb{Z}_p \times \mathbb{Z}_p/p^3 \times \mathbb{Z}_p/p^2$.

Fitting ideals

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fact: Fitting ideals are well defined.

Example

Consider $R = \mathbb{Z}_p$ and $M = \mathbb{Z}_p \times \mathbb{Z}_p/p^3 \times \mathbb{Z}_p/p^2$. A resolution is given by

$$(\mathbb{Z}_p)^3 \xrightarrow{\mu} (\mathbb{Z}_p)^3 \xrightarrow{\varepsilon} M \longrightarrow 0$$

Here ε is the natural map and μ is given by the matrix $A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & p^3 & 0 \\ 0 & 0 & p^2 \end{pmatrix}$

Fitting ideals

Definition (Fitting ideal)

Let M be a finitely generated R -module. Choose a resolution

$$R^n \xrightarrow{A} R^m \longrightarrow M \longrightarrow 0$$

$\text{Fitt}_i^R(M)$ is the ideal generated by the minors of size $(m - i)$ of A .

Fact: Fitting ideals are well defined.

Example

Consider $R = \mathbb{Z}_p$ and $M = \mathbb{Z}_p \times \mathbb{Z}_p/p^3 \times \mathbb{Z}_p/p^2$. A resolution is given by

$$(\mathbb{Z}_p)^3 \xrightarrow{\mu} (\mathbb{Z}_p)^3 \xrightarrow{\varepsilon} M \longrightarrow 0$$

Here ε is the natural map and μ is given by the matrix $A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & p^3 & 0 \\ 0 & 0 & p^2 \end{pmatrix}$

$\text{Fitt}_0(M) = (0),$	$\text{Fitt}_1(M) = (p^5),$
$\text{Fitt}_2(M) = (p^2) + (p^3) = (p^2)$	$\text{Fitt}_i(M) = (1) \forall i \geq 3$

Fitting ideals over Discrete Valuation Rings

Kolyvagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

$$\blacksquare i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$$

Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$

Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$
- $i \geq r+s \Rightarrow \text{Fitt}_i(M) = (1)$.

Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$
- $i \geq r+s \Rightarrow \text{Fitt}_i(M) = (1)$.

Corollary

The Fitting ideals determine i up to isomorphism:

Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$
- $i \geq r+s \Rightarrow \text{Fitt}_i(M) = (1)$.

Corollary

The Fitting ideals determine i up to isomorphism:

- r is the minimum i such that $\text{Fitt}_i(M) \neq 0$.

Fitting ideals over Discrete Valuation Rings

Let R be a DVR (with maximal ideal \mathfrak{m} and residue field κ). Then

$$M \cong R^r \times R/\mathfrak{m}^{\alpha_1} \times \cdots \times R/\mathfrak{m}^{\alpha_s}$$

for some non-negative integers $r, s, \alpha_1 \geq \cdots \geq \alpha_s$.

Proposition

- $i \in \{0, \dots, r-1\} \Rightarrow \text{Fitt}_i(M) = (0)$
- $j \in \{0, \dots, s-1\} \Rightarrow \text{Fitt}_{r+j} = \prod_{k=j+1}^s \mathfrak{m}^{i_k} = \mathfrak{m}^{\sum_{k=j+1}^s i_k}$
- $i \geq r+s \Rightarrow \text{Fitt}_i(M) = (1).$

Corollary

The Fitting ideals determine i up to isomorphism:

- r is the minimum i such that $\text{Fitt}_i(M) \neq 0$.
- For $i \geq 0$, $\alpha_i = \text{Fitt}_{r+i+1}(M) \text{Fitt}_{r+i}(M)^{-1}$.

Definition (dual Galois modules)

- Pontryagin dual: $\mathbf{T}^\vee = \text{Hom}(\mathbf{T}, \mathbb{Q}_p/\mathbb{Z}_p)$.
- Cartier dual: $\mathbf{T}^* = \text{Hom}(\mathbf{T}, \mu_{p^\infty})$.

Definition (dual Galois modules)

- Pontryagin dual: $\mathbf{T}^\vee = \text{Hom}(\mathbf{T}, \mathbb{Q}_p/\mathbb{Z}_p)$.
- Cartier dual: $\mathbf{T}^* = \text{Hom}(\mathbf{T}, \mu_{p^\infty})$.

Proposition (local duality)

The cup-product induces a non-degenerate pairing

$$H^1(\mathbb{Q}_\ell, \mathbf{T}) \times H^1(\mathbb{Q}_\ell, \mathbf{T}^*) \rightarrow H^2(\mathbb{Q}_\ell, \mu_{p^\infty}) \cong \mathbb{Q}_p/\mathbb{Z}_p$$

Moreover, $H_f^1(\mathbb{Q}_\ell, \mathbf{T})$ and $H_f^1(\mathbb{Q}_\ell, \mathbf{T}^*)$ are exact annihilators of each other.

Definition (dual Galois modules)

- Pontryagin dual: $\mathbf{T}^\vee = \text{Hom}(\mathbf{T}, \mathbb{Q}_p/\mathbb{Z}_p)$.
- Cartier dual: $\mathbf{T}^* = \text{Hom}(\mathbf{T}, \mu_{p^\infty})$.

Proposition (local duality)

The cup-product induces a non-degenerate pairing

$$H^1(\mathbb{Q}_\ell, \mathbf{T}) \times H^1(\mathbb{Q}_\ell, \mathbf{T}^*) \rightarrow H^2(\mathbb{Q}_\ell, \mu_{p^\infty}) \cong \mathbb{Q}_p/\mathbb{Z}_p$$

Moreover, $H_f^1(\mathbb{Q}_\ell, \mathbf{T})$ and $H_f^1(\mathbb{Q}_\ell, \mathbf{T}^*)$ are exact annihilators of each other.

Corollary

$$H^1(\mathbb{Q}_\ell, \mathbf{T})^\vee \cong H^1(\mathbb{Q}_\ell, \mathbf{T}^*)$$

Definition (dual Galois modules)

- Pontryagin dual: $\mathbf{T}^\vee = \text{Hom}(\mathbf{T}, \mathbb{Q}_p/\mathbb{Z}_p)$.
- Cartier dual: $\mathbf{T}^* = \text{Hom}(\mathbf{T}, \mu_{p^\infty})$.

Proposition (local duality)

The cup-product induces a non-degenerate pairing

$$H^1(\mathbb{Q}_\ell, \mathbf{T}) \times H^1(\mathbb{Q}_\ell, \mathbf{T}^*) \rightarrow H^2(\mathbb{Q}_\ell, \mu_{p^\infty}) \cong \mathbb{Q}_p/\mathbb{Z}_p$$

Moreover, $H_f^1(\mathbb{Q}_\ell, \mathbf{T})$ and $H_f^1(\mathbb{Q}_\ell, \mathbf{T}^*)$ are exact annihilators of each other.

Corollary

$$H^1(\mathbb{Q}_\ell, \mathbf{T})^\vee \cong H^1(\mathbb{Q}_\ell, \mathbf{T}^*)$$

$$H_f^1(\mathbb{Q}_\ell, \mathbf{T})^\vee \cong \frac{H^1(\mathbb{Q}_\ell, \mathbf{T}^*)}{H_f^1(\mathbb{Q}_\ell, \mathbf{T}^*)}$$

Definition (dual Selmer structure)

The **dual Selmer structure** \mathcal{F}^* is defined by the local conditions

$$H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T}^*) := \text{Ann}(H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T})) \subset H^1(\mathbb{Q}_\ell, \mathbf{T}^*)$$

These are the elements of $H^1(\mathbb{Q}_\ell, \mathbf{T}^*)$ which annihilate $H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T)$ under the local duality pairing.

Dual Selmer structure

Definition (dual Selmer structure)

The **dual Selmer structure** \mathcal{F}^* is defined by the local conditions

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T}^*) := \text{Ann}(H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T})) \subset H^1(\mathbb{Q}_{\ell}, \mathbf{T}^*)$$

These are the elements of $H^1(\mathbb{Q}_{\ell}, \mathbf{T}^*)$ which annihilate $H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, T)$ under the local duality pairing.

Remark (well defined)

The dual Selmer structure is well defined since

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T}^*) := \text{Ann}(H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T}))$$

Let \mathcal{F} and \mathcal{G} be Selmer structures such that

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, T) \subset H_{\mathcal{G}}^1(\mathbb{Q}_{\ell}, T) \quad \forall \ell$$

Global duality

Let \mathcal{F} and \mathcal{G} be Selmer structures such that

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, T) \subset H_{\mathcal{G}}^1(\mathbb{Q}_{\ell}, T) \quad \forall \ell$$

Then the dual local conditions satisfy the opposite relations

$$H_{\mathcal{G}^*}^1(\mathbb{Q}_{\ell}, T^*) \subset H_{\mathcal{F}^*}^1(\mathbb{Q}_{\ell}, T^*) \quad \forall \ell$$

Let \mathcal{F} and \mathcal{G} be Selmer structures such that

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, T) \subset H_{\mathcal{G}}^1(\mathbb{Q}_{\ell}, T) \quad \forall \ell$$

Then the dual local conditions satisfy the opposite relations

$$H_{\mathcal{G}^*}^1(\mathbb{Q}_{\ell}, T^*) \subset H_{\mathcal{F}^*}^1(\mathbb{Q}_{\ell}, T^*) \quad \forall \ell$$

Clearly,

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T}) \subset \mathrm{Sel}_{\mathcal{G}}(\mathbb{Q}, \mathbf{T}), \quad \mathrm{Sel}_{\mathcal{G}^*}(\mathbb{Q}, \mathbf{T}^*) \subset \mathrm{Sel}_{\mathcal{F}^*}(\mathbb{Q}, \mathbf{T}^*)$$

Global duality

Let \mathcal{F} and \mathcal{G} be Selmer structures such that

$$H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T) \subset H_{\mathcal{G}}^1(\mathbb{Q}_\ell, T) \quad \forall \ell$$

Then the dual local conditions satisfy the opposite relations

$$H_{\mathcal{G}^*}^1(\mathbb{Q}_\ell, T^*) \subset H_{\mathcal{F}^*}^1(\mathbb{Q}_\ell, T^*) \quad \forall \ell$$

Clearly,

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T) \subset \mathrm{Sel}_{\mathcal{G}}(\mathbb{Q}, T), \quad \mathrm{Sel}_{\mathcal{G}^*}(\mathbb{Q}, T^*) \subset \mathrm{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)$$

Global duality

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T) & \longrightarrow & \mathrm{Sel}_{\mathcal{G}}(\mathbb{Q}, T) & \longrightarrow & \prod_{\ell} \frac{H_{\mathcal{G}}^1(\mathbb{Q}_\ell, T)}{H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T)} \\
 & & & & & & \searrow \\
 & & & & & & \mathrm{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)^{\vee} \longrightarrow \mathrm{Sel}_{\mathcal{G}^*}(\mathbb{Q}, T^*)^{\vee} \longrightarrow 0
 \end{array}$$

Assumptions

Kolyvagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

- (H0) Let $p \geq 5$ and let \mathbf{T} be free \mathbb{Z}_p -module of finite rank endowed with a continuous action of $G_{\mathbb{Q}}$, ramifying only at a finite amount of primes.
- (H1) $\rho : G_{\mathbb{Q}} \rightarrow \text{Aut}(\mathbf{T})$ is surjective.
- (H2) $H^1(\mathbb{Q}_{\ell}, \mathbf{T})/H^1_{\mathcal{F}}(\mathbb{Q}_{\ell}, \mathbf{T})$ is a torsion-free \mathbb{Z}_p -module.

Propagation to positive characteristic

Fix $k \in \mathbb{N}$ and let $T = \mathbf{T}/p^k$. Denote $\pi : \mathbf{T} \rightarrow T$ to the canonical projection.

Kolyvagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

Propagation to positive characteristic

Fix $k \in \mathbb{N}$ and let $T = \mathbf{T}/p^k$. Denote $\pi : \mathbf{T} \rightarrow T$ to the canonical projection.

Definition (propagated local condition)

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, T) = \pi(H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T}))$$

Propagation to positive characteristic

Fix $k \in \mathbb{N}$ and let $T = \mathbf{T}/p^k$. Denote $\pi : \mathbf{T} \rightarrow T$ to the canonical projection.

Definition (propagated local condition)

$$H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, T) = \pi(H_{\mathcal{F}}^1(\mathbb{Q}_{\ell}, \mathbf{T}))$$

Proposition

Under assumptions (H0), (H1) and (H2), the following equalities hold true.

Propagation to positive characteristic

Fix $k \in \mathbb{N}$ and let $T = \mathbf{T}/p^k$. Denote $\pi : \mathbf{T} \rightarrow T$ to the canonical projection.

Definition (propagated local condition)

$$H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T) = \pi(H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T}))$$

Proposition

Under assumptions (H0), (H1) and (H2), the following equalities hold true.

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T) = \mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T})/p^k$$

Propagation to positive characteristic

Fix $k \in \mathbb{N}$ and let $T = \mathbf{T}/p^k$. Denote $\pi : \mathbf{T} \rightarrow T$ to the canonical projection.

Definition (propagated local condition)

$$H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T) = \pi(H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T}))$$

Proposition

Under assumptions (H0), (H1) and (H2), the following equalities hold true.

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T) = \mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T})/p^k$$

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T}^*)[p^k]$$

Propagation to positive characteristic

Fix $k \in \mathbb{N}$ and let $T = \mathbf{T}/p^k$. Denote $\pi : \mathbf{T} \rightarrow T$ to the canonical projection.

Definition (propagated local condition)

$$H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T) = \pi(H_{\mathcal{F}}^1(\mathbb{Q}_\ell, \mathbf{T}))$$

Proposition

Under assumptions (H0), (H1) and (H2), the following equalities hold true.

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T) = \mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T})/p^k$$

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T}^*)[p^k]$$

Remark

A study of $\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T}/p^k)$ for all k will determine $\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, \mathbf{T})$.

Definition

A prime ℓ is a **Kolyvagin prime** if

Definition

A prime ℓ is a **Kolyvagin prime** if

- $\ell \equiv 1 \pmod{p^k}.$

Definition

A prime ℓ is a **Kolyvagin prime** if

- $\ell \equiv 1 \pmod{p^k}$.
- $P_\ell(1) = \det(1 - \text{Frob}_\ell | T) = 0$.

Definition

A prime ℓ is a **Kolyvagin prime** if

- $\ell \equiv 1 \pmod{p^k}$.
- $P_\ell(1) = \det(1 - \text{Frob}_\ell | T) = 0$.

Notation

\mathcal{P} denotes the set of Kolyvagin primes.

Definition

A prime ℓ is a **Kolyvagin prime** if

- $\ell \equiv 1 \pmod{p^k}$.
- $P_\ell(1) = \det(1 - \text{Frob}_\ell | T) = 0$.

Notation

\mathcal{P} denotes the set of Kolyvagin primes.

$\mathcal{N}(\mathcal{P})$ denotes the set of square free products of Kolyvagin primes.

Definition

A prime ℓ is a **Kolyvagin prime** if

- $\ell \equiv 1 \pmod{p^k}$.
- $P_\ell(1) = \det(1 - \text{Frob}_\ell | T) = 0$.

Notation

\mathcal{P} denotes the set of Kolyvagin primes.

$\mathcal{N}(\mathcal{P})$ denotes the set of square free products of Kolyvagin primes.

$\mathcal{N}_i(\mathcal{P})$ denotes the set of square free products of exactly i Kolyvagin primes.

Transverse local condition and finite-singular map

Definition (transverse local condition)

$$H_{tr}^1(\mathbb{Q}_\ell, T) := \text{Im}(H^1(\mathbb{Q}_\ell(\mu_\ell)/\mathbb{Q}_\ell, T) \rightarrow H^1(\mathbb{Q}_\ell, T))$$

Transverse local condition and finite-singular map

Definition (transverse local condition)

$$H_{tr}^1(\mathbb{Q}_\ell, T) := \text{Im}(H^1(\mathbb{Q}_\ell(\mu_\ell)/\mathbb{Q}_\ell, T) \rightarrow H^1(\mathbb{Q}_\ell, T))$$

Proposition (split of the local cohomology group)

If ℓ is Kolyvagin prime, then

$$H^1(\mathbb{Q}_\ell, T) = H_f^1(\mathbb{Q}_\ell, T) \oplus H_{tr}^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k \oplus \mathbb{Z}/p^k$$

Transverse local condition and finite-singular map

Definition (transverse local condition)

$$H_{tr}^1(\mathbb{Q}_\ell, T) := \text{Im}(H^1(\mathbb{Q}_\ell(\mu_\ell)/\mathbb{Q}_\ell, T) \rightarrow H^1(\mathbb{Q}_\ell, T))$$

Proposition (split of the local cohomology group)

If ℓ is Kolyvagin prime, then

$$H^1(\mathbb{Q}_\ell, T) = H_f^1(\mathbb{Q}_\ell, T) \oplus H_{tr}^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k \oplus \mathbb{Z}/p^k$$

Definition (finite-singular map)

There is a canonical isomorphism

$$\phi_{fs} : H_f^1(\mathbb{Q}_\ell, T) \cong H_{tr}^1(\mathbb{Q}_\ell, T)$$

Modified Selmer structures

Kolyvagin
systems
and
fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

Let $a, b, c \in \mathbb{N}$ be such that abc is square free.

Assume all primes dividing a , b and c are Kolyvagin primes.

We can define a new Selmer structure $\mathcal{F}_a^b(c)$ by

Modified Selmer structures

Let $a, b, c \in \mathbb{N}$ be such that abc is square free.

Assume all primes dividing a , b and c are Kolyvagin primes.

We can define a new Selmer structure $\mathcal{F}_a^b(c)$ by

$$\blacksquare H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) := H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T) \text{ if } \ell \nmid abc.$$

Modified Selmer structures

Let $a, b, c \in \mathbb{N}$ be such that abc is square free.

Assume all primes dividing a , b and c are Kolyvagin primes.

We can define a new Selmer structure $\mathcal{F}_a^b(c)$ by

- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) := H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T)$ if $\ell \nmid abc$.
- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) = 0$ if $\ell \mid a$.

Modified Selmer structures

Let $a, b, c \in \mathbb{N}$ be such that abc is square free.

Assume all primes dividing a , b and c are Kolyvagin primes.

We can define a new Selmer structure $\mathcal{F}_a^b(c)$ by

- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) := H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T)$ if $\ell \nmid abc$.
- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) = 0$ if $\ell \mid a$.
- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) = H^1(\mathbb{Q}_\ell, T)$ if $\ell \mid b$.

Modified Selmer structures

Let $a, b, c \in \mathbb{N}$ be such that abc is square free.

Assume all primes dividing a , b and c are Kolyvagin primes.

We can define a new Selmer structure $\mathcal{F}_a^b(c)$ by

- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) := H_{\mathcal{F}}^1(\mathbb{Q}_\ell, T)$ if $\ell \nmid abc$.
- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) = 0$ if $\ell \mid a$.
- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) = H^1(\mathbb{Q}_\ell, T)$ if $\ell \mid b$.
- $H_{\mathcal{F}_a^b(c)}^1(\mathbb{Q}_\ell, T) = H_{\text{tr}}^1(\mathbb{Q}_\ell, T)$ if $\ell \mid c$.

Definition (Kolyvagin system)

A **Kolyvagin system** is a collection of elements

$$\kappa_n \in \mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \subset H^1(\mathbb{Q}, T)$$

for every $n \in \mathcal{N}(\mathcal{P})$, satisfying the following Kolyvagin conditions.

Definition (Kolyvagin system)

A **Kolyvagin system** is a collection of elements

$$\kappa_n \in \text{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \subset H^1(\mathbb{Q}, T)$$

for every $n \in \mathcal{N}(\mathcal{P})$, satisfying the following Kolyvagin conditions.

For every $n \in \mathcal{N}(\mathcal{P})$ and $\ell \in \mathcal{P}$ such that $\ell \nmid n$, consider the localization maps at ℓ .

$$\text{loc}_{\ell}(\kappa_n) \in H^1_{\mathcal{F}(n)}(\mathbb{Q}_{\ell}, T) = H^1_f(\mathbb{Q}_{\ell}, T)$$

$$\text{loc}_{\ell}(\kappa_{n\ell}) \in H^1_{\mathcal{F}(n\ell)}(\mathbb{Q}, T) = H^1_{tr}(\mathbb{Q}_{\ell}, T)$$

Definition (Kolyvagin system)

A **Kolyvagin system** is a collection of elements

$$\kappa_n \in \text{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \subset H^1(\mathbb{Q}, T)$$

for every $n \in \mathcal{N}(\mathcal{P})$, satisfying the following Kolyvagin conditions.

For every $n \in \mathcal{N}(\mathcal{P})$ and $\ell \in \mathcal{P}$ such that $\ell \nmid n$, consider the localization maps at ℓ .

$$\text{loc}_{\ell}(\kappa_n) \in H_{\mathcal{F}(n)}^1(\mathbb{Q}_{\ell}, T) = H_f^1(\mathbb{Q}_{\ell}, T)$$

$$\text{loc}_{\ell}(\kappa_{n\ell}) \in H_{\mathcal{F}(n\ell)}^1(\mathbb{Q}, T) = H_{tr}^1(\mathbb{Q}_{\ell}, T)$$

The **Kolyvagin condition** for $n \in \mathcal{N}(\mathcal{P})$ and $\ell \in \mathcal{P}$ is

$$\phi_{fs}(\kappa_n) = \kappa_{n\ell}$$

Definition (Kolyvagin system)

A **Kolyvagin system** is a collection of elements

$$\kappa_n \in \text{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \subset H^1(\mathbb{Q}, T)$$

for every $n \in \mathcal{N}(\mathcal{P})$, satisfying the following Kolyvagin conditions.

For every $n \in \mathcal{N}(\mathcal{P})$ and $\ell \in \mathcal{P}$ such that $\ell \nmid n$, consider the localization maps at ℓ .

$$\text{loc}_{\ell}(\kappa_n) \in H_{\mathcal{F}(n)}^1(\mathbb{Q}_{\ell}, T) = H_f^1(\mathbb{Q}_{\ell}, T)$$

$$\text{loc}_{\ell}(\kappa_{n\ell}) \in H_{\mathcal{F}(n\ell)}^1(\mathbb{Q}, T) = H_{tr}^1(\mathbb{Q}_{\ell}, T)$$

The **Kolyvagin condition** for $n \in \mathcal{N}(\mathcal{P})$ and $\ell \in \mathcal{P}$ is

$$\phi_{fs}(\kappa_n) = \kappa_{n\ell}$$

Notation

The module of Kolyvagin systems will be denoted by $\text{KS}(T, \mathcal{F})$.

Definition/proposition (core rank)

There exists a non-negative integer $\chi(\mathcal{F})$ and a non-canonical homomorphism

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T) \cong \left(\mathbb{Z}/p^k\right)^{\chi(T)} \oplus \mathrm{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)$$

(possibly after swapping the roles of T and T^* .)

The integer $\chi(T)$ is called the **core rank** of T .

Definition/proposition (core rank)

There exists a non-negative integer $\chi(\mathcal{F})$ and a non-canonical homomorphism

$$\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T) \cong \left(\mathbb{Z}/p^k\right)^{\chi(T)} \oplus \mathrm{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)$$

(possibly after swapping the roles of T and T^* .)

The integer $\chi(T)$ is called the **core rank** of T .

Proposition (Sakamoto, 2021)

$$\chi(\mathcal{F}_a^b(c)) = \chi(\mathcal{F}) + \nu(b) - \nu(a)$$

where $\nu(b)$ and $\nu(a)$ are the number of primes dividing b and a , respectively.

Core rank and Kolyvagin systems

Theorem (Mazur-Rubin, 2004)

- If $\chi(\mathcal{F}) = 0$, then $\text{KS}(T, \mathcal{F}) = 0$.

There are no Kolyvagin system to control the Selmer group. We will see a possible solution later in the talk.

Core rank and Kolyvagin systems

Theorem (Mazur-Rubin, 2004)

- If $\chi(\mathcal{F}) = 0$, then $\text{KS}(T, \mathcal{F}) = 0$.

There are no Kolyvagin system to control the Selmer group. We will see a possible solution later in the talk.

- If $\chi(\mathcal{F}) = 1$, then $\text{KS}(T, \mathcal{F}) \cong \mathbb{Z}/p^k$.

A generator of $\text{KS}(T, \mathcal{F})$ is called a **primitive Kolyvagin system**. We will see next that they carry information to compute all the Fitting ideals of the Selmer group $\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)$.

Core rank and Kolyvagin systems

Theorem (Mazur-Rubin, 2004)

- If $\chi(\mathcal{F}) = 0$, then $\text{KS}(T, \mathcal{F}) = 0$.

There are no Kolyvagin system to control the Selmer group. We will see a possible solution later in the talk.

- If $\chi(\mathcal{F}) = 1$, then $\text{KS}(T, \mathcal{F}) \cong \mathbb{Z}/p^k$.

A generator of $\text{KS}(T, \mathcal{F})$ is called a **primitive Kolyvagin system**. We will see next that they carry information to compute all the Fitting ideals of the Selmer group $\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)$.

- If $\chi(\mathcal{F}) > 1$, then $\text{KS}(T, \mathcal{F})$ is too large.

In order to compute the Selmer group, [Mazur-Rubin, 2016] and [Burns-Sakamoto-Sano, 2025] modified the definition of Kolyvagin system in (biduals of) exterior powers of the Selmer groups.

Selmer groups of core rank 1

Definition (order of a Kolyagin element)

$$\text{ord}(\kappa_n) := \max \left\{ j \in \{0, \dots, k\} : \kappa_n \in p^j H_{\mathcal{F}(n)}^1(\mathbb{Q}, T) \right\}$$

Selmer groups of core rank 1

Definition (order of a Kolyvagin element)

$$\text{ord}(\kappa_n) := \max \left\{ j \in \{0, \dots, k\} : \kappa_n \in p^j H_{\mathcal{F}(n)}^1(\mathbb{Q}, T) \right\}$$

Proposition

If κ is a primitive Kolyvagin system

$$\text{ord}(\kappa_n) = \min \left\{ k, \text{length} \left(H_{(\mathcal{F}^*)_{(n)}}^1(\mathbb{Q}, T^*) \right) \right\}$$

Selmer groups of core rank 1

Definition (order of a Kolyagin element)

$$\text{ord}(\kappa_n) := \max \left\{ j \in \{0, \dots, k\} : \kappa_n \in p^j H_{\mathcal{F}(n)}^1(\mathbb{Q}, T) \right\}$$

Proposition

If κ is a primitive Kolyagin system

$$\text{ord}(\kappa_n) = \min \left\{ k, \text{length} \left(H_{(\mathcal{F}^*)_{(n)}}^1(\mathbb{Q}, T^*) \right) \right\}$$

Definition

$$\Theta_i := (p)^{\min\{\text{ord}(\kappa_n) : n \in \mathcal{N}_i\}}$$

Selmer groups of core rank 1

Definition (order of a Kolyvagin element)

$$\text{ord}(\kappa_n) := \max \left\{ j \in \{0, \dots, k\} : \kappa_n \in p^j H_{\mathcal{F}(n)}^1(\mathbb{Q}, T) \right\}$$

Proposition

If κ is a primitive Kolyvagin system

$$\text{ord}(\kappa_n) = \min \left\{ k, \text{length} \left(H_{(\mathcal{F}^*)_{(n)}}^1(\mathbb{Q}, T^*) \right) \right\}$$

Definition

$$\Theta_i := (p)^{\min\{\text{ord}(\kappa_n) : n \in \mathcal{N}_i\}}$$

Theorem (Mazur-Rubin, 2004)

When $\chi(\mathcal{F}) = 1$ and κ is a primitive Kolyvagin system

$$\Theta_i = \text{Fitt}_i \left(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T)^* \right)$$

Core rank 0

- We have seen that there are no non-zero Kolyvagin systems.
- Choose a prime ℓ such that $H_f^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k$.
- Note that all Kolyvagin primes satisfy the above condition, but we do not restrict to them.
- Then \mathcal{F}^ℓ is cartesian and $\chi(\mathcal{F}^\ell) = 1$.

Core rank 0

- We have seen that there are no non-zero Kolyvagin systems.
- Choose a prime ℓ such that $H_f^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k$.
- Note that all Kolyvagin primes satisfy the above condition, but we do not restrict to them.
- Then \mathcal{F}^ℓ is cartesian and $\chi(\mathcal{F}^\ell) = 1$.

Definition

Let $\kappa \in \text{KS}(T, \mathcal{F}^\ell)$. Define

$$\delta_n = \delta_n(\kappa) := \text{loc}_\ell(\kappa_n) \in H^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k$$

Core rank 0

- We have seen that there are no non-zero Kolyvagin systems.
- Choose a prime ℓ such that $H_f^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k$.
- Note that all Kolyvagin primes satisfy the above condition, but we do not restrict to them.
- Then \mathcal{F}^ℓ is cartesian and $\chi(\mathcal{F}^\ell) = 1$.

Definition

Let $\kappa \in \text{KS}(T, \mathcal{F}^\ell)$. Define

$$\delta_n = \delta_n(\kappa) := \text{loc}_\ell(\kappa_n) \in H^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k$$

Definition (order)

$$\text{ord}(\delta_n) = \max \left\{ j \in \{0, \dots, k\} : \delta^n \in (p^j) \right\}$$

Core rank 0

- We have seen that there are no non-zero Kolyvagin systems.
- Choose a prime ℓ such that $H_f^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k$.
- Note that all Kolyvagin primes satisfy the above condition, but we do not restrict to them.
- Then \mathcal{F}^ℓ is cartesian and $\chi(\mathcal{F}^\ell) = 1$.

Definition

Let $\kappa \in \text{KS}(T, \mathcal{F}^\ell)$. Define

$$\delta_n = \delta_n(\kappa) := \text{loc}_\ell(\kappa_n) \in H^1(\mathbb{Q}_\ell, T) \cong \mathbb{Z}/p^k$$

Definition (order)

$$\text{ord}(\delta_n) = \max \left\{ j \in \{0, \dots, k\} : \delta^n \in (p^j) \right\}$$

Proposition (Kim, 2025)

$$\text{ord}(\delta_n) = \min \left\{ k, \text{length} \left(H_{(\mathcal{F}^*)_{(n)}}^1(\mathbb{Q}, T^*) \right) \right\}$$

Fitting ideals of Selmer groups of rank 0

Definition

$$\Theta_i := (p)^{\min\{\text{ord}(\delta_n) : n \in \mathcal{N}_i(\mathcal{P})\}} = \langle \{\delta_n : n \in \mathcal{N}_i(\mathcal{P})\} \rangle \subset \mathbb{Z}/p^k$$

Fitting ideals of Selmer groups of rank 0

Definition

$$\Theta_i := (p)^{\min\{\text{ord}(\delta_n) : n \in \mathcal{N}_i(\mathcal{P})\}} = \langle \{\delta_n : n \in \mathcal{N}_i(\mathcal{P})\} \rangle \subset \mathbb{Z}/p^k$$

Theorem (A., 2025)

For all i , we have

$$\Theta_i \subset \text{Fitt}_i(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*))$$

Fitting ideals of Selmer groups of rank 0

Definition

$$\Theta_i := (p)^{\min\{\text{ord}(\delta_n) : n \in \mathcal{N}_i(\mathcal{P})\}} = \langle \{\delta_n : n \in \mathcal{N}_i(\mathcal{P})\} \rangle \subset \mathbb{Z}/p^k$$

Theorem (A., 2025)

For all i , we have

$$\Theta_i \subset \text{Fitt}_i(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*))$$

The equality for some index i holds if any of the following is true:

Fitting ideals of Selmer groups of rank 0

Definition

$$\Theta_i := (p)^{\min\{\text{ord}(\delta_n) : n \in \mathcal{N}_i(\mathcal{P})\}} = \langle \{\delta_n : n \in \mathcal{N}_i(\mathcal{P})\} \rangle \subset \mathbb{Z}/p^k$$

Theorem (A., 2025)

For all i , we have

$$\Theta_i \subset \text{Fitt}_i(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*))$$

The equality for some index i holds if any of the following is true:

- $\Theta_{i-1} \subsetneq \text{Fitt}_{i-1}(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*)).$

Fitting ideals of Selmer groups of rank 0

Definition

$$\Theta_i := (p)^{\min\{\text{ord}(\delta_n) : n \in \mathcal{N}_i(\mathcal{P})\}} = \langle \{\delta_n : n \in \mathcal{N}_i(\mathcal{P})\} \rangle \subset \mathbb{Z}/p^k$$

Theorem (A., 2025)

For all i , we have

$$\Theta_i \subset \text{Fitt}_i(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*))$$

The equality for some index i holds if any of the following is true:

- $\Theta_{i-1} \subsetneq \text{Fitt}_{i-1}(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*)).$
- $\Theta_{i-1} = \text{Fitt}_{i-1}(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*)) = 0.$

Fitting ideals of Selmer groups of rank 0

Definition

$$\Theta_i := (p)^{\min\{\text{ord}(\delta_n) : n \in \mathcal{N}_i(\mathcal{P})\}} = \langle \{\delta_n : n \in \mathcal{N}_i(\mathcal{P})\} \rangle \subset \mathbb{Z}/p^k$$

Theorem (A., 2025)

For all i , we have

$$\Theta_i \subset \text{Fitt}_i(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*))$$

The equality for some index i holds if any of the following is true:

- $\Theta_{i-1} \subsetneq \text{Fitt}_{i-1}(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*))$.
- $\Theta_{i-1} = \text{Fitt}_{i-1}(H_{\mathcal{F}^*}^1(\mathbb{Q}, T^*)) = 0$.

Remark

When $T = \mathbf{T}/p^k$ for some \mathbb{Z}_p -module \mathbf{T} and k is large enough, the ideals Θ_i determine $H_{\mathcal{F}^*}^1(\mathbb{Q}, \mathbf{T}^*)$ up to isomorphism

Galois representations which are not residually self-dual

Kolyvagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

Theorem (A., 2025)

Under the following assumption on non self-duality,

- (N1) $T/p \not\cong T^*[p]$.

Galois representations which are not residually self-dual

Kolyvagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

Theorem (A., 2025)

Under the following assumption on non self-duality,

- (N1) $T/p \not\cong T^*[p]$.

Then for all $i \in \mathbb{Z}_{\geq 0}$, we have the equality

$$\Theta_i \subset \text{Fitt}_i(H_{\mathcal{F}}^1(\mathbb{Q}, T^*))$$

Connection to Euler systems

- Assume that we have an Euler system z .

Connection to Euler systems

- Assume that we have an Euler system \mathbf{z} .
- The Kolyvagin derivative operator produces a Kolyvagin system for \mathbf{T}/p^k for all k and the *canonical Selmer structure*, defined as

$$\begin{cases} H_{\mathcal{F}^{\text{can}}}^1(\mathbb{Q}_\ell, \mathbf{T}/p^k) = H_f^1(\mathbb{Q}_\ell, \mathbf{T}/p^k) \text{ if } \ell \neq p, \infty \\ H_{\mathcal{F}^{\text{can}}}^1(\mathbb{Q}_p, \mathbf{T}/p^k) = H^1(\mathbb{Q}_p, \mathbf{T}/p^k) \\ H_{\mathcal{F}^{\text{can}}}^1(\mathbb{R}, \mathbf{T}/p^k) = H^1(\mathbb{R}, \mathbf{T}/p^k) \end{cases}$$

This is also known as *relaxed at p* . Its dual Selmer structure will be called *restricted at p* .

- Assume that we have an Euler system \mathbf{z} .
- The Kolyvagin derivative operator produces a Kolyvagin system for \mathbf{T}/p^k for all k and the *canonical Selmer structure*, defined as

$$\begin{cases} H_{\mathcal{F}^{\text{can}}}^1(\mathbb{Q}_\ell, \mathbf{T}/p^k) = H_f^1(\mathbb{Q}_\ell, \mathbf{T}/p^k) \text{ if } \ell \neq p, \infty \\ H_{\mathcal{F}^{\text{can}}}^1(\mathbb{Q}_p, \mathbf{T}/p^k) = H^1(\mathbb{Q}_p, \mathbf{T}/p^k) \\ H_{\mathcal{F}^{\text{can}}}^1(\mathbb{R}, \mathbf{T}/p^k) = H^1(\mathbb{R}, \mathbf{T}/p^k) \end{cases}$$

This is also known as *relaxed at p* . Its dual Selmer structure will be called *restricted at p* .

- We call $\text{ord}(\mathbf{z}_{\mathbb{Q}}) = \sup \{j \in \mathbb{N} : \mathbf{z}_{\mathbb{Q}} \in p^j H^1(\mathbb{Q}, \mathbf{T})_{/\text{tors}}\}$.

Theorem (Kolyvagin, 1995)

$$\text{ord}(\mathbf{z}_{\mathbb{Q}}) \geq \text{ord}(\kappa_1) \geq \text{length} \left(H_{(\mathcal{F}^{\text{can}})^*}^1(\mathbb{Q}, \mathbf{T}^*) \right)$$

Elliptic curves: construction of the Euler system

We apply the results to $\mathbf{T} = T_p E \otimes \chi$.

Elliptic curves: construction of the Euler system

Kolyvagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

We apply the results to $\mathbf{T} = T_p E \otimes \chi$. We assume the following:

- (E0) E is defined over \mathbb{Q} .
- (χ 1) The conductor of χ is not divisible by p or any bad prime of E .
- (χ 2) The order of χ is prime to p .

Elliptic curves: construction of the Euler system

We apply the results to $\mathbf{T} = T_p E \otimes \chi$. We assume the following:

- (E0) E is defined over \mathbb{Q} .
- (χ 1) The conductor of χ is not divisible by p or any bad prime of E .
- (χ 2) The order of χ is prime to p .

Modularity There exists a modular form $f_\chi = \sum \chi(n) a_n q^n$ such that

$$T_{f_\chi} = T_p E \otimes \chi$$

Elliptic curves: construction of the Euler system

We apply the results to $T = T_p E \otimes \chi$. We assume the following:

- (E0) E is defined over \mathbb{Q} .
- (χ 1) The conductor of χ is not divisible by p or any bad prime of E .
- (χ 2) The order of χ is prime to p .

Modularity There exists a modular form $f_\chi = \sum \chi(n) a_n q^n$ such that

$$T_{f_\chi} = T_p E \otimes \chi$$

Kato constructed an Euler system for this representation.

Elliptic curves: Bloch-Kato Selmer structure

Bloch-Kato Selmer structure

The **classical local conditions** are defined by Bloch-Kato condition

$$\begin{cases} H_{\mathcal{F}_{BK}}^1(\mathbb{Q}_\ell, \mathbf{T}) = H_f^1(\mathbb{Q}_\ell, \mathbf{T}) \quad \forall \ell \neq p \\ H_{\mathcal{F}_{BK}}^1(\mathbb{Q}_p, \mathbf{T}) = \ker \left(H^1(\mathbb{Q}_p, \mathbf{T}) \rightarrow H^1(\mathbb{Q}_p, \mathbf{T} \otimes_{Z_p} B_{\text{crys}}) \right) \end{cases}$$

Elliptic curves: Bloch-Kato Selmer structure

Bloch-Kato Selmer structure

The **classical local conditions** are defined by Bloch-Kato condition

$$\begin{cases} H_{\mathcal{F}_{BK}}^1(\mathbb{Q}_\ell, \mathbf{T}) = H_f^1(\mathbb{Q}_\ell, \mathbf{T}) \quad \forall \ell \neq p \\ H_{\mathcal{F}_{BK}}^1(\mathbb{Q}_p, \mathbf{T}) = \ker \left(H^1(\mathbb{Q}_p, \mathbf{T}) \rightarrow H^1(\mathbb{Q}_p, \mathbf{T} \otimes_{Z_p} B_{\text{crys}}) \right) \end{cases}$$

Assume the following:

- (E1) $\rho : G_{\mathbb{Q}} \rightarrow \text{Aut}(\mathbf{T})$ is surjective.

Elliptic curves: Bloch-Kato Selmer structure

Kolyagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

Bloch-Kato Selmer structure

The **classical local conditions** are defined by Bloch-Kato condition

$$\begin{cases} H_{\mathcal{F}_{BK}}^1(\mathbb{Q}_\ell, \mathbf{T}) = H_f^1(\mathbb{Q}_\ell, \mathbf{T}) \quad \forall \ell \neq p \\ H_{\mathcal{F}_{BK}}^1(\mathbb{Q}_p, \mathbf{T}) = \ker \left(H^1(\mathbb{Q}_p, \mathbf{T}) \rightarrow H^1(\mathbb{Q}_p, \mathbf{T} \otimes_{Z_p} B_{\text{crys}}) \right) \end{cases}$$

Assume the following:

- (E1) $\rho : G_{\mathbb{Q}} \rightarrow \text{Aut}(\mathbf{T})$ is surjective.

Proposition

Assuming (E1), \mathcal{F}_{BK} satisfies all the assumptions (H0), (H1) and (H2) and $\chi(\mathcal{F}_{BK}) = 0$.

Elliptic curves: construction of the Kolyvagin system

Kolyvagin derivative

The Kolyvagin derivative operator applied to Kato's Euler system, produces a Kolyvagin system for $\mathcal{F}^{\text{can}} = (\mathcal{F}_{BK})^p$.

Elliptic curves: construction of the Kolyvagin system

Kolyvagin derivative

The Kolyvagin derivative operator applied to Kato's Euler system, produces a Kolyvagin system for $\mathcal{F}^{\text{can}} = (\mathcal{F}_{BK})^p$.

Denote K_χ to the fixed field of χ . Assume:

- $(\chi^3) E((K_\chi)_p)[p] = \{O\}$ for every prime p above p .

Elliptic curves: construction of the Kolyvagin system

Kolyvagin derivative

The Kolyvagin derivative operator applied to Kato's Euler system, produces a Kolyvagin system for $\mathcal{F}^{\text{can}} = (\mathcal{F}_{BK})^p$.

Denote K_χ to the fixed field of χ . Assume:

- $(\chi 3) E((K_\chi)_p)[p] = \{O\}$ for every prime p above p .

Proposition

Assuming $(\chi 3)$, the Selmer structure $(\mathcal{F}_{BK})^p$ satisfies all the assumptions (H0), (H1) and (H2) and $\chi((\mathcal{F}_{BK})^p) = 1$.

Elliptic curves: construction of the Kolyvagin system

Kolyvagin derivative

The Kolyvagin derivative operator applied to Kato's Euler system, produces a Kolyvagin system for $\mathcal{F}^{\text{can}} = (\mathcal{F}_{BK})^p$.

Denote K_χ to the fixed field of χ . Assume:

- $(\chi 3)$ $E((K_\chi)_p)[p] = \{O\}$ for every prime p above p .

Proposition

Assuming $(\chi 3)$, the Selmer structure $(\mathcal{F}_{BK})^p$ satisfies all the assumptions (H0), (H1) and (H2) and $\chi((\mathcal{F}_{BK})^p) = 1$.

Assume further:

- $(\chi 4)$ The Tamagawa numbers of E over K_χ are prime to p .
- $(\chi 5)$ Iwasawa main conjecture (in the sense of Kato) holds for f_χ .

Proposition

The Kolyvagin derivative produced from Kato's Euler system is a primitive Kolyvagin system.

Elliptic curves: Kurihara numbers

The final goal is to compute $\delta_{n,\chi} = \text{loc}_p(\kappa_n)$. Assume

- (E2) The Manin constant is prime to p .

Elliptic curves: Kurihara numbers

The final goal is to compute $\delta_{n,\chi} = \text{loc}_p(\kappa_n)$. Assume

- (E2) The Manin constant is prime to p .

Proposition (Kurihara numbers)

Let n be a square-free product of Kolyvagin systems for \mathbf{T}/p^k .

Elliptic curves: Kurihara numbers

The final goal is to compute $\delta_{n,\chi} = \text{loc}_p(\kappa_n)$. Assume

- (E2) The Manin constant is prime to p .

Proposition (Kurihara numbers)

Let n be a square-free product of Kolyvagin systems for \mathbf{T}/p^k .

$$\delta_{n,\chi} = \sum_{a \in (\mathbb{Z}/nc)^*} \chi(a) \left(\left[\frac{a}{cn} \right]^+ + \left[\frac{a}{cn} \right]^- \right) \prod_{\ell|n} \log_{\eta_\ell}(a) \in \mathbb{Z}_p[\chi]/p^k$$

Elliptic curves: Kurihara numbers

The final goal is to compute $\delta_{n,\chi} = \text{loc}_p(\kappa_n)$. Assume

- (E2) The Manin constant is prime to p .

Proposition (Kurihara numbers)

Let n be a square-free product of Kolyvagin systems for \mathbf{T}/p^k .

$$\delta_{n,\chi} = \sum_{a \in (\mathbb{Z}/nc)^*} \chi(a) \left(\left[\frac{a}{cn} \right]^+ + \left[\frac{a}{cn} \right]^- \right) \prod_{\ell|n} \log_{\eta_\ell}(a) \in \mathbb{Z}_p[\chi]/p^k$$

where

- c is the conductor of χ .

Elliptic curves: Kurihara numbers

The final goal is to compute $\delta_{n,\chi} = \text{loc}_p(\kappa_n)$. Assume

- (E2) The Manin constant is prime to p .

Proposition (Kurihara numbers)

Let n be a square-free product of Kolyvagin systems for \mathbf{T}/p^k .

$$\delta_{n,\chi} = \sum_{a \in (\mathbb{Z}/nc)^*} \chi(a) \left(\left[\frac{a}{cn} \right]^+ + \left[\frac{a}{cn} \right]^- \right) \prod_{\ell|n} \log_{\eta_\ell}(a) \in \mathbb{Z}_p[\chi]/p^k$$

where

- c is the conductor of χ .
- $\left[\frac{a}{cn} \right]^\pm$ are the real and imaginary part of the modular symbols of E .

Elliptic curves: Kurihara numbers

The final goal is to compute $\delta_{n,\chi} = \text{loc}_p(\kappa_n)$. Assume

- (E2) The Manin constant is prime to p .

Proposition (Kurihara numbers)

Let n be a square-free product of Kolyvagin systems for \mathbf{T}/p^k .

$$\delta_{n,\chi} = \sum_{a \in (\mathbb{Z}/nc)^*} \chi(a) \left(\left[\frac{a}{cn} \right]^+ + \left[\frac{a}{cn} \right]^- \right) \prod_{\ell|n} \log_{\eta_\ell}(a) \in \mathbb{Z}_p[\chi]/p^k$$

where

- c is the conductor of χ .
- $\left[\frac{a}{cn} \right]^\pm$ are the real and imaginary part of the modular symbols of E .
- η_ℓ is a primitive root of $(\mathbb{Z}/\ell)^\times$ and $\log_{\eta_\ell}(a)$ is the image of the logarithm under the projection $(\mathbb{Z}/\ell)^\times \cong \mathbb{Z}/(\ell-1) \rightarrow \mathbb{Z}/p^k$.

Elliptic curves: Kurihara numbers

The final goal is to compute $\delta_{n,\chi} = \text{loc}_p(\kappa_n)$. Assume

- (E2) The Manin constant is prime to p .

Proposition (Kurihara numbers)

Let n be a square-free product of Kolyvagin systems for \mathbf{T}/p^k .

$$\delta_{n,\chi} = \sum_{a \in (\mathbb{Z}/nc)^*} \chi(a) \left(\left[\frac{a}{cn} \right]^+ + \left[\frac{a}{cn} \right]^- \right) \prod_{\ell|n} \log_{\eta_\ell}(a) \in \mathbb{Z}_p[\chi]/p^k$$

where

- c is the conductor of χ .
- $\left[\frac{a}{cn} \right]^\pm$ are the real and imaginary part of the modular symbols of E .
- η_ℓ is a primitive root of $(\mathbb{Z}/\ell)^\times$ and $\log_{\eta_\ell}(a)$ is the image of the logarithm under the projection $(\mathbb{Z}/\ell)^\times \cong \mathbb{Z}/(\ell-1) \rightarrow \mathbb{Z}/p^k$.

Remark

If K/\mathbb{Q} is an abelian extension such that all the characters of $\text{Gal}(K/\mathbb{Q})$ satisfy $(\chi 1) - (\chi 5)$, then the **twisted Kurihara numbers** determine $\text{Sel}(K, E[p^\infty])$ up to isomorphism of $\mathbb{Z}_p[\text{Gal}(K/\mathbb{Q})]$ -modules.

Proofs: first inequality

Proposition

Consider the exact sequence of (\mathbb{Z}/p^k) -modules

$$0 \longrightarrow C \longrightarrow M \xrightarrow{\phi} (\mathbb{Z}/p^k)^i$$

Then

$$(p)^{\text{length}(C)} \subset \text{Fitt}_i(M)$$

Proofs: first inequality

Proposition

Consider the exact sequence of (\mathbb{Z}/p^k) -modules

$$0 \longrightarrow C \longrightarrow M \xrightarrow{\phi} (\mathbb{Z}/p^k)^i$$

Then

$$(p)^{\text{length}(C)} \subset \text{Fitt}_i(M)$$

If we choose ϕ and C maximizing the image of ϕ , the equality holds.

Proofs: first inequality

Proposition

Consider the exact sequence of (\mathbb{Z}/p^k) -modules

$$0 \longrightarrow C \longrightarrow M \xrightarrow{\phi} (\mathbb{Z}/p^k)^i$$

Then

$$(p)^{\text{length}(C)} \subset \text{Fitt}_i(M)$$

If we choose ϕ and C maximizing the image of ϕ , the equality holds.

Theorem (first inequality)

$$\Theta_i \subset \text{Fitt}_i(\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee})$$

Proofs: first inequality

Proposition

Consider the exact sequence of (\mathbb{Z}/p^k) -modules

$$0 \longrightarrow C \longrightarrow M \xrightarrow{\phi} (\mathbb{Z}/p^k)^i$$

Then

$$(p)^{\text{length}(C)} \subset \text{Fitt}_i(M)$$

If we choose ϕ and C maximizing the image of ϕ , the equality holds.

Theorem (first inequality)

$$\Theta_i \subset \text{Fitt}_i(\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee})$$

Proof Apply the proposition to

$$0 \longrightarrow \text{Sel}_{(\mathcal{F}^*)_n}(\mathbb{Q}, T) \longrightarrow \text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*) \longrightarrow \prod_{\ell|n} H_f^1(\mathbb{Q}_{\ell}, T^*) \cong (\mathbb{Z}/p^k)^{\nu(n)}$$

Proofs: first inequality

Proposition

Consider the exact sequence of (\mathbb{Z}/p^k) -modules

$$0 \longrightarrow C \longrightarrow M \xrightarrow{\phi} (\mathbb{Z}/p^k)^i$$

Then

$$(p)^{\text{length}(C)} \subset \text{Fitt}_i(M)$$

If we choose ϕ and C maximizing the image of ϕ , the equality holds.

Theorem (first inequality)

$$\Theta_i \subset \text{Fitt}_i(\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee})$$

Proof Apply the proposition to

$$0 \longrightarrow \text{Sel}_{(\mathcal{F}^*)_n}(\mathbb{Q}, T) \longrightarrow \text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*) \longrightarrow \prod_{\ell|n} H_f^1(\mathbb{Q}_{\ell}, T^*) \cong (\mathbb{Z}/p^k)^{\nu(n)}$$

Then

$$(p)^{\text{length}(\text{Sel}_{(\mathcal{F}^*)_n}(\mathbb{Q}, T^*))} \subset (p)^{\text{length}(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*))} \subset \text{Fitt}_i(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T))$$

Proofs: first inequality

Proposition

Consider the exact sequence of (\mathbb{Z}/p^k) -modules

$$0 \longrightarrow C \longrightarrow M \xrightarrow{\phi} (\mathbb{Z}/p^k)^i$$

Then

$$(p)^{\text{length}(C)} \subset \text{Fitt}_i(M)$$

If we choose ϕ and C maximizing the image of ϕ , the equality holds.

Theorem (first inequality)

$$\Theta_i \subset \text{Fitt}_i(\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee})$$

Proof Apply the proposition to

$$0 \longrightarrow \text{Sel}_{(\mathcal{F}^*)_n}(\mathbb{Q}, T) \longrightarrow \text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*) \longrightarrow \prod_{\ell|n} H_f^1(\mathbb{Q}_{\ell}, T^*) \cong (\mathbb{Z}/p^k)^{\nu(n)}$$

Then

$$(p)^{\text{length}(\text{Sel}_{(\mathcal{F}^*)_n}(\mathbb{Q}, T^*))} \subset (p)^{\text{length}(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*))} \subset \text{Fitt}_i(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T))$$

The proof is completed by taking the minimum over all $n \in \mathcal{N}_i(\mathcal{P})$.

Proofs: what is needed for the equality?

- The localization map

$$\mathrm{loc}_\ell : \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*) \rightarrow H_f^1(\mathbb{Q}_\ell, T^*)$$

has the largest possible image.

Proofs: what is needed for the equality?

- The localization map

$$\mathrm{loc}_\ell : \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*) \rightarrow H_f^1(\mathbb{Q}_\ell, T^*)$$

has the largest possible image.

This can be achieved using Chebotarev density theorem.

Proofs: what is needed for the equality?

- The localization map

$$\mathrm{loc}_\ell : \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*) \rightarrow H_f^1(\mathbb{Q}_\ell, T^*)$$

has the largest possible image.

This can be achieved using Chebotarev density theorem.

- We want to choose $n \in \mathcal{N}(\mathcal{P})$ such that

$$\mathrm{Sel}_{(\mathcal{F}^*)_n}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*)$$

Proofs: what is needed for the equality?

- The localization map

$$\text{loc}_\ell : \text{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*) \rightarrow H_f^1(\mathbb{Q}_\ell, T^*)$$

has the largest possible image.

This can be achieved using Chebotarev density theorem.

- We want to choose $n \in \mathcal{N}(\mathcal{P})$ such that

$$\text{Sel}_{(\mathcal{F}^*)_n}(\mathbb{Q}, T^*) = \text{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*)$$

Proposition

Assume the following localization map is surjective

$$\text{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow H_f^1(\mathbb{Q}_\ell, T)$$

Then

$$\text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) = \text{Sel}_{(\mathcal{F}^*)_{\ell(n)}}(\mathbb{Q}, T^*)$$

Proofs: what is needed for the equality?

Proposition

Assume the following localization map is surjective

$$\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow H_f^1(\mathbb{Q}_\ell, T)$$

Then

$$\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_{\ell(n)}}(\mathbb{Q}, T^*)$$

Proof



$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathrm{Sel}_{\mathcal{F}_\ell(n)}(\mathbb{Q}, T) & \longrightarrow & \mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) & \longrightarrow & H_f^1(\mathbb{Q}_\ell, T) \\
& & & & & & \searrow \\
& & & & & & \mathrm{Sel}_{(\mathcal{F}^*)^\ell(n)}(\mathbb{Q}, T^*)^\vee & \longrightarrow & \mathrm{Sel}_{(\mathcal{F}^*)(n)}(\mathbb{Q}, T^*)^\vee & \longrightarrow & 0
\end{array}$$

Proofs: what is needed for the equality?

Proposition

Assume the following localization map is surjective

$$\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow H_f^1(\mathbb{Q}_\ell, T)$$

Then

$$\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)$$

Proof

■

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{Sel}_{\mathcal{F}_\ell(n)}(\mathbb{Q}, T) & \longrightarrow & \mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) & \longrightarrow & H_f^1(\mathbb{Q}_\ell, T) \\ & & & & & & \searrow \\ & & & & & & \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)^\vee \longrightarrow \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*)^\vee \longrightarrow 0 \end{array}$$

■ $\mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)$

Proofs: what is needed for the equality?

Proposition

Assume the following localization map is surjective

$$\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow H_f^1(\mathbb{Q}_\ell, T)$$

Then

$$\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)$$

Proof

■

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{Sel}_{\mathcal{F}_\ell(n)}(\mathbb{Q}, T) & \longrightarrow & \mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) & \longrightarrow & H_f^1(\mathbb{Q}_\ell, T) \\ & & & & & & \searrow \\ & & & & & & \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)^\vee \longrightarrow \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*)^\vee \longrightarrow 0 \end{array}$$

■ $\mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)$

■ $\mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)} = \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}} \cap \mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}$

Proofs: what is needed for the equality?

Proposition

Assume the following localization map is surjective

$$\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow H_f^1(\mathbb{Q}_\ell, T)$$

Then

$$\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)$$

Proof

■

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{Sel}_{\mathcal{F}_\ell(n)}(\mathbb{Q}, T) & \longrightarrow & \mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) & \longrightarrow & H_f^1(\mathbb{Q}_\ell, T) \\ & & & & & \searrow & \uparrow \\ & & & & & \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)^\vee & \longrightarrow & \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*)^\vee & \longrightarrow & 0 \end{array}$$

■ $\mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)$

■ $\mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)} = \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}} \cap \mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}} = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)} \cap \mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}$

Proofs: what is needed for the equality?

Proposition

Assume the following localization map is surjective

$$\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow H_f^1(\mathbb{Q}_\ell, T)$$

Then

$$\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)$$

Proof

■

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathrm{Sel}_{\mathcal{F}_\ell(n)}(\mathbb{Q}, T) & \longrightarrow & \mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) & \longrightarrow & H_f^1(\mathbb{Q}_\ell, T) \\ & & & & & \searrow & \text{hook} \\ & & & & & \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)^\vee & \longrightarrow & \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*)^\vee & \longrightarrow & 0 \end{array}$$

■ $\mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T^*) = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)}(\mathbb{Q}, T^*)$

■ $\mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)} = \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}} \cap \mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}} = \mathrm{Sel}_{(\mathcal{F}^*)_\ell(n)} \cap \mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}} = \mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}$

Proofs: equality in rank one

Proposition

$$\mathrm{Fitt}_i(\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee}) = \Theta_i := (p)^{\min\{\mathrm{ord}(\kappa_n): n \in \mathcal{N}_i(\mathcal{P})\}}$$

Proofs: equality in rank one

Proposition

$$\mathrm{Fitt}_i(\mathrm{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee}) = \Theta_i := (p)^{\min\{\mathrm{ord}(\kappa_n) : n \in \mathcal{N}_i(\mathcal{P})\}}$$

Proof Inductively, assume we have constructed some n_i such that

$$\mathrm{Fitt}_i(\mathrm{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T^*)^{\vee}) = p^{\mathrm{ord}(\kappa_{n_i})}$$

Proofs: equality in rank one

Proposition

$$\text{Fitt}_i(\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee}) = \Theta_i := (p)^{\min\{\text{ord}(\kappa_n) : n \in \mathcal{N}_i(\mathcal{P})\}}$$

Proof Inductively, assume we have constructed some n_i such that

$$\text{Fitt}_i(\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T^*)^{\vee}) = p^{\text{ord}(\kappa_{n_i})}$$

Since the core rank is one,

$$\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T) \cong (\mathbb{Z}/p^k) \oplus \text{Sel}_{\mathcal{F}^*(n_i)}(\mathbb{Q}, T^*)$$

Then there exists a surjective map $\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T) \rightarrow \mathbb{Z}/p^k$.

Proofs: equality in rank one

Proposition

$$\text{Fitt}_i(\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^\vee) = \Theta_i := (p)^{\min\{\text{ord}(\kappa_n) : n \in \mathcal{N}_i(\mathcal{P})\}}$$

Proof Inductively, assume we have constructed some n_i such that

$$\text{Fitt}_i(\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T^*)^\vee) = p^{\text{ord}(\kappa_{n_i})}$$

Since the core rank is one,

$$\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T) \cong (\mathbb{Z}/p^k) \oplus \text{Sel}_{\mathcal{F}^*(n_i)}(\mathbb{Q}, T^*)$$

Then there exists a surjective map $\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T) \rightarrow \mathbb{Z}/p^k$.

By Chebotarev density theorem, we can find a prime ℓ_{i+1} such that

- $\text{loc}_{\ell_{i+1}} : \text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T) \rightarrow H_f^1(\mathbb{Q}_\ell, T)$ is surjective.
- $\text{loc}_{\ell_{i+1}} : \text{Sel}_{\mathcal{F}^*(n_i)}(\mathbb{Q}, T^*) \rightarrow H_f^1(\mathbb{Q}_\ell, T^*)$ has maximal image.

Proofs: equality in rank one

Proposition

$$\text{Fitt}_i(\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee}) = \Theta_i := (p)^{\min\{\text{ord}(\kappa_n) : n \in \mathcal{N}_i(\mathcal{P})\}}$$

Proof Inductively, assume we have constructed some n_i such that

$$\text{Fitt}_i(\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T^*)^{\vee}) = p^{\text{ord}(\kappa_{n_i})}$$

Since the core rank is one,

$$\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T) \cong (\mathbb{Z}/p^k) \oplus \text{Sel}_{\mathcal{F}^*(n_i)}(\mathbb{Q}, T^*)$$

Then there exists a surjective map $\text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T) \rightarrow \mathbb{Z}/p^k$.

By Chebotarev density theorem, we can find a prime ℓ_{i+1} such that

- $\text{loc}_{\ell_{i+1}} : \text{Sel}_{\mathcal{F}(n_i)}(\mathbb{Q}, T) \rightarrow H_f^1(\mathbb{Q}_{\ell}, T)$ is surjective.
- $\text{loc}_{\ell_{i+1}} : \text{Sel}_{\mathcal{F}^*(n_i)}(\mathbb{Q}, T^*) \rightarrow H_f^1(\mathbb{Q}_{\ell}, T^*)$ has maximal image.

For $n_{i+1} := n_i \ell_{i+1}$, we get that

$$\text{Sel}_{(\mathcal{F}^*)(n_{i+1})}(\mathbb{Q}, T^*) = \text{Sel}_{(\mathcal{F}^*)_{\ell_{i+1}}(n_i)}(\mathbb{Q}, T^*)$$

Moreover,

$$(p)^{\text{ord}(\kappa_{n_{i+1}})} = (p)^{\text{length}(\text{Sel}_{(\mathcal{F}^*)(n_{i+1})})} = \text{Fitt}_{i+1}(\text{Sel}_{(\mathcal{F}^*)}(\mathbb{Q}, T))$$

Proofs: equality in rank zero: characteristic reduction

When $\chi(\mathcal{F}) = 0$,

$$\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \cong \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T) \quad \forall n \in \mathcal{N}(\mathcal{P})$$

It might not exist a surjective map $\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow \mathbb{Z}/p^k$.

Proofs: equality in rank zero: characteristic reduction

When $\chi(\mathcal{F}) = 0$,

$$\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \cong \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T) \quad \forall n \in \mathcal{N}(\mathcal{P})$$

It might not exist a surjective map $\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow \mathbb{Z}/p^k$.

By the structure theorem,

$$\mathrm{Sel}_{\mathcal{F}(n)} = \mathbb{Z}/p^{e_1} \times \cdots \times \mathbb{Z}/p^{e_s}$$

for some $e_1 \geq \cdots \geq e_s$.

Trick Swap T by $T_{e_1} := T/p^{e_1}$.

Proofs: equality in rank zero: characteristic reduction

When $\chi(\mathcal{F}) = 0$,

$$\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \cong \mathrm{Sel}_{(\mathcal{F}^*)_{(n)}}(\mathbb{Q}, T) \quad \forall n \in \mathcal{N}(\mathcal{P})$$

It might not exist a surjective map $\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T) \rightarrow \mathbb{Z}/p^k$.

By the structure theorem,

$$\mathrm{Sel}_{\mathcal{F}(n)} = \mathbb{Z}/p^{e_1} \times \cdots \times \mathbb{Z}/p^{e_s}$$

for some $e_1 \geq \cdots \geq e_s$.

Trick Swap T by $T_{e_1} := T/p^{e_1}$.

Similarly, we can find a prime ℓ such that the maps

- $\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, T_{e_1}) \rightarrow H_f^1(\mathbb{Q}_\ell, T_{e_1})$.
- $\mathrm{Sel}_{\mathcal{F}(n)}(\mathbb{Q}, (T_{e_1})^*) \rightarrow H_f^1(\mathbb{Q}_\ell, (T_{e_1})^*)$.

are surjective. We obtain the following for the Selmer group over T_{e_1} .

$$\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, (T_{e_1})^*) = \mathrm{Sel}_{(\mathcal{F}^*)_{\ell}(n)}(\mathbb{Q}, (T_{e_1})^*) \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}$$

Proofs: equality in rank zero: recover information

What information can we deduce from this to the Selmer group over T ?

Kolyvagin
systems
and
Fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs

What information can we deduce from this to the Selmer group over T ?

■ $\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*)[p^{e_1}] \cong \mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, (T_{e_1})^*) \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}.$

What information can we deduce from this to the Selmer group over T ?

■ $\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*)[p^{e_1}] \cong \mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, (T_{e_1})^*) \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}.$

■ If either $e_1 = k$ or $e_1 > e_2$, we can conclude that

$$\mathrm{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}$$

What information can we deduce from this to the Selmer group over T ?

- $\text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*)[p^{e_1}] \cong \text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, (T_{e_1})^*) \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}.$

- If either $e_1 = k$ or $e_1 > e_2$, we can conclude that

$$\text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}$$

- When $e_1 = e_2$, we only know that

$$\begin{aligned} \text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) &\subset \text{Sel}_{(\mathcal{F}^*)_{\ell}(n)}(\mathbb{Q}, T^*) \cong \\ \mathbb{Z}/p^k \times \text{Sel}_{\mathcal{F}_{\ell}(n)}(\mathbb{Q}, T) &\cong \mathbb{Z}/p^k \times \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s} \end{aligned}$$

What information can we deduce from this to the Selmer group over T ?

- $\text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*)[p^{e_1}] \cong \text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, (T_{e_1})^*) \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}.$

- If either $e_1 = k$ or $e_1 > e_2$, we can conclude that

$$\text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}$$

- When $e_1 = e_2$, we only know that

$$\begin{aligned} \text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) &\subset \text{Sel}_{(\mathcal{F}^*)_{\ell}(n)}(\mathbb{Q}, T^*) \cong \\ &\mathbb{Z}/p^k \times \text{Sel}_{\mathcal{F}_{\ell}(n)}(\mathbb{Q}, T) \cong \mathbb{Z}/p^k \times \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s} \end{aligned}$$

- The structure theorem implies that

$$\text{Sel}_{(\mathcal{F}^*)_{(n\ell)}}(\mathbb{Q}, T^*) \cong \mathbb{Z}/p^{f_2} \times \mathbb{Z}/p^{e_3} \times \cdots \times \mathbb{Z}/p^{e_s}$$

for some $f_2 \geq e_2$.

Proofs: equality in rank zero: inductive step

We start with $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)$ and choose a prime ℓ_1 such that the localization maps for T_{e_1} and $T_{e_1}^*$ are surjective, and minimizing f_2 .

We have two cases

Proofs: equality in rank zero: inductive step

We start with $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)$ and choose a prime ℓ_1 such that the localization maps for T_{e_1} and $T_{e_1}^*$ are surjective, and minimizing f_2 .

We have two cases

- If $f_2 = e_2$, then $\Theta_1 = \text{Fitt}_1(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)^{\vee})$.

Proofs: equality in rank zero: inductive step

We start with $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)$ and choose a prime ℓ_1 such that the localization maps for T_{e_1} and $T_{e_1}^*$ are surjective, and minimizing f_2 .

We have two cases

- If $f_2 = e_2$, then $\Theta_1 = \text{Fitt}_1(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)^{\vee})$.
- If $f_2 > e_2$, then $\Theta_1 \subsetneq \text{Fitt}_1(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)^{\vee})$.

Proofs: equality in rank zero: inductive step

We start with $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)$ and choose a prime ℓ_1 such that the localization maps for T_{e_1} and $T_{e_1}^*$ are surjective, and minimizing f_2 .

We have two cases

- If $f_2 = e_2$, then $\Theta_1 = \text{Fitt}_1(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)^{\vee})$.
- If $f_2 > e_2$, then $\Theta_1 \subsetneq \text{Fitt}_1(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)^{\vee})$.

In this case,

$$\text{Sel}_{\mathcal{F}(\ell_1)} \cong \mathbb{Z}/p^{f_2} \times \mathbb{Z}/p^{e_3} \times \cdots \times \mathbb{Z}/p^{e_s}$$

Since $f_2 > e_3$, we can choose a prime ℓ_2 in a way such that $f_3 = e_3$, so

$$\Theta_2 = \text{Fitt}_2(\text{Sel}_{\mathcal{F}^*}(\mathbb{Q}, T^*)^{\vee})$$

Non self-dual representations

- Chebotarev density theorem is stronger in this case:

For every pair of subgroups $C \subset \text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)$ and $D \subset \text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)$ such that the quotients $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)/C$ and $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)/D$ are cyclic, we can find a prime ℓ such that the kernels of the localization maps are C and D .

Non self-dual representations

- Chebotarev density theorem is stronger in this case:

For every pair of subgroups $C \subset \text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)$ and $D \subset \text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)$ such that the quotients $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)/C$ and $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)/D$ are cyclic, we can find a prime ℓ such that the kernels of the localization maps are C and D .

- If $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T) \cong \mathbb{Z}/p^{e_1} \times \cdots \times \mathbb{Z}/p^{e_s}$, a technical argument constructs a prime ℓ such that

$$\text{Sel}_{\mathcal{F}(\ell)} \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}$$

Non self-dual representations

- Chebotarev density theorem is stronger in this case:

For every pair of subgroups $C \subset \text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)$ and $D \subset \text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)$ such that the quotients $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T)/C$ and $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)/D$ are cyclic, we can find a prime ℓ such that the kernels of the localization maps are C and D .

- If $\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T) \cong \mathbb{Z}/p^{e_1} \times \cdots \times \mathbb{Z}/p^{e_s}$, a technical argument constructs a prime ℓ such that

$$\text{Sel}_{\mathcal{F}(\ell)} \cong \mathbb{Z}/p^{e_2} \times \cdots \times \mathbb{Z}/p^{e_s}$$

- Therefore, the equality $\Theta_i = \text{Fitt}_i(\text{Sel}_{\mathcal{F}}(\mathbb{Q}, T^*)^{\vee})$ holds for all i .

Thank you for your attention!



Preprint

Kolyvagin
systems
and
fitting
ideals of
Selmer
group of
rank 0

Introduction

Selmer
structures

Fitting
ideals

Duality
pairings

Kolyvagin
systems

Structure
of Selmer
groups

Euler
system

Elliptic
Curves

Proofs