Universidad Autonoma de Nuevo Leon
Facultad de Ingenieria Mecanica y Electrica
Posgrado de Ingenieria de Sistemas

RESEARCH PROJECT PROPOSAL

Name of Candidate: Jose Alberto Benavides Vazquez

Date of Admission to this programme: March 10, 2022

Full time mode

Abstract

Brief summary of the proposal and its sections.

1. Project Title

Impact of the air pollutants in the population health in the Monterrey Metropolitan Area.

2. Introduction

This study focuses on the impact that the air quality have in the population welfare primarily in the Monterrey Metropolitan Area between 2018 and 2019 obtained from the Comprehensive Environmental Monitoring System (abbreviated SIMA in Spanish from Sistema Integral de Monitoreo Ambiental) [32]. Also, it have been incorporated a database containing the hospital discharge records for Mexico over the same span of time from the open repository of the Department of Health of Mexico [12].

As the atmosphere is result of the combination of 78% of N_2, 21% of O_2 and 1% of combined helium, water vapor, oxides, methane, noble gases, sulfides, ozone, ammonia, formaldehyde, and hydroxyl radical, can be defined that the air is *polluted* when these concentrations are altered by human activities resulting in harmful environments to biological ecosystems [13].

The human activities that cause pollution have been increasing over the years. Air quality became an urban problem since 1930 when a fog in Belgium was associated with the death of 63 people and then in London during 1952 [10] when another fog caused $4,000$ deaths [28] being pneumonia the principal illness diagnosed in both cases. Its negative impact on health and general welfare concerned authorities that design regulations in order to minimize those undesirable effects. In 1981 it became an international problem when the CO concentrations were seen from space for the first time [1]. Such images show that the pollution was not only generated from expected sources, but from other unforeseen activities, and that the pollution generated in one place affected distant populations over time.

Air pollution can be caused by events that impact the atmosphere, like forest fires, increased volcanic activity, animal and vegetable decomposition, climate change [21], etc. The main difference from natural and human pollutants is that the natural ones tend to occur far from large human populations and are less often than human sources of pollution, which are generally located in or near large human populations, and their emissions are more frequent. The main human sources of air pollution are associated with transportation, fuel combustion, and industrial processes [8].

The International Classification of Diseases (ICD)[1] is a code associated to diseases by their trends and statistical values, established by the [35]. It consists, generally, in a letter and three

---

[1]The name of the diseases corresponding to the ICD code is obtained from https://icd.who.int/browse10/2019/en

numbers between 0 and 9. For example, the code for `pneumonia due to staphylococcus` is `J152`. This system comprehends from the values `A00.0` to `Z99.9`, leaving the `U` letter for yet unknown sources. This classification was proposed by Farr at the end of the XIX century [34], and its purpose is to understand the causes of morbidity and mortality in order to improve the quality of life of the human population [34].

3. Hypothesis

The contamination levels reported can explain the statistically significant variations and trends of georeferenced data from health reports. Also, it is possible to automatize these relationships in an interactive and free web service.

4. Project objectives

- Stablish a clear and scientifically sustained relationship between air pollution and the human diseases that them can cause.
- Study the impact of the air pollution in the Metropolitan Monterrey Area during 2017 and 2019 in the health population.
- Choose or propose a methodology to calculate the causation between air pollution and associated human diseases.
- Identify the areas where they are more air pollution and stablish a relationship between the people that live nearby those areas.
- Propose a rank to determine which air pollutants have the most direct effect in the human health.
- Propose pollution emission limits, scientifically justified, that can be used by organizations interested in this matter.
- Generate interactive and free tools that let people and specialist know the levels, relationships, risks, etc. related to the air pollutants studied.

5. Research questions

- Is it possible to stablish statistically significant relationships between air pollution and human diseases?
- What are the best methods to detect the causation between air pollution and human diseases?
- How to eliminate the climate and seasonal components from the relationship between air pollution and human diseases?
- Does the environmental and geographical factors can be used to predict the amount of air pollution in an area and time?
- Which air pollutants cause more human diseases?
- It is possible to stablish short (days), medium (months) and large (years) interactions between air pollution and human diseases?

6. Background and related literature

The first correspondence to discuss relationship between air quality and human health was registered during December 1930 in Belgium, when a three-day fog s was declared to be the cause of death of 63 people — during the fog, disease and death were observed to increase, whereas after its dissipation, the figures normalized again [10]. Similarly, in December 1952, a four-day fog in London was associated with four thousand deaths [28]. The majority of the fog-related deaths were related to respiratory diseases.

In 1959 Lawther [24] used a method to quantify the severity of respiratory illness based on their condition (negative numbers indicated they were recovered and positive numbers that they were getting worse). Then, he plotted a time series superposed with concentrations of two pollutants — SO_2 and smoke — and two weather variables — temperature and humidity; he discovered that the pollutants bore a similarity to the severity whereas the climate ones did not.

During the 1970s, short-term exposure to different pollutants was studied [7, 15, 16, 27] with experiments that implied some ethical and legal considerations, focusing on physiological analysis of the subjects. A compilation was published in by Ferris [9] who documented that the pollutants, in regulated concentrations, had a negative impact on health, primarily on respiratory diseases.

A set of methodologies are used to measure the relationships that are the focus of this study. They are **multiple regression analysis** [17, 18, 20, 25] with variations in the distributions such as logistic regressions [22], **multivariate analysis** [14, 23], **auto regressive models** [19], **causality models** [30] and **case-crossover approximations** [4, 29].

7. Methodology

The study will focus in the **Monterrey Metropolitan Area (MMA)** between 2018 and 2019. The air quality samples in the are taken each hour from 13 sensors over the MMA that measure concentrations of **CO**, **NO**, **NO$_2$**, **O$_3$**, **SO$_2$**, **PM$_{10}$**, **PM$_{2.5}$**, and **atmospheric pressure**, **rainfall**, **relative humidity**, **solar radiation**, **temperature**, **wind velocity and direction**. This data is provided by the *Sistema Integral de Monitoreo Ambiental de Nuevo León* (SIMA) [32]. The Mexican diseases data was obtained from the Mexican *Department of Health* [31] and contains information from all states and municipalities in Mexico such as **date of admission**, **egress date**, **age at the admission**, **gender**, **weight**, **height**, **ICD code upon arrived**, **ICD code upon diagnosis**, and **reason of egress**.

The air quality data needs to be interpolated because it contains imputed records. Different temporal interpolation techniques are used and compared [11]. Also, an spatio-temporal interpolation is performed to obtain the missing data values [26]. Both data sets are processed and converted to georeferenced time series [33] that are stationary [20] in order to establish their relationship by cross correlation [6], multiple regression analysis [3], vector autorregresive approaches, causality models [30] and geographic interactions [5]. The results are ranked by metrics like R^2, the Akaike (AIC), and the Bayesian (BIC) information criteria [2]. Finally, it will be produced a web application that allow general and specialized population to interact with the data and obtain forecasts, interactions and visualizations of the models described.

8. Expected results

9. Significance

While the Introduction mention the relevance of the research, here it need to be highlighted and demonstrated. Focus specially in the contributions that the results could bring to the state of the art and/or the solutions that can provide to the problems involved.

10.   External Collaboration

If any, include authors, published papers, works in research programs or conferences where the research have been involved.

11.   Project schedule

Add a schedule that includes the steps of the process, from investigation, possible publications and participation in conferences, to final products like thesis, libraries…A Gantt diagram is recommended.

References

[1] Hajime Akimoto. Global air quality and pollution. Science, 302(5651), 2003. doi: 10.1126/science.1092666.

[2] Jim Albert. Bayesian computation with R. Springer-Verlag GmbH, July 2007. doi: 10. 1007/978-0-387-92298-0. URL https://www.ebook.de/de/product/24984339/jim_albert_ bayesian_computation_with_r.html.

[3] Peter Brockwell. Introduction to time series and forecasting. Springer, New York, 2002. doi: 10.1007/978-3-319-29854-2.

[4] Eduardo Carracedo-Martínez, Aurelio Tobías, Marc Saez, Margarita Taracido, and Adolfo Figueiras. Fundamentos y aplicaciones del diseño de casos cruzados. Gaceta Sanitaria, 23: 161–165, April 2009. ISSN 0213-9111. URL http://scielo.isciii.es/scielo.php?script= sci_arttext&pid=S0213-91112009000200017&nrm=iso.

[5] Alexis Comber and Wen Zeng. Spatial interpolation using areal features: A review of methods and opportunities using new forms of data with coded illustrations. Geography Compass, 13 (10), August 2019. doi: 10.1111/gec3.12465.

[6] Timothy R. Derrick and Joshua M. Thomas. Innovative analyses of human movement, chapter Time Series Analysis: The Cross-Correlation Function, pages 189–205. Kinesiology Publications, 2004.

[7] Richard Ehrlich, J. C. Findlay, J. D. Fenters, and D. E. Gardner. Health effects of short-term inhalation of nitrogen dioxide and ozone mixtures. Environmental Research, 14(2):223–231, October 1977. doi: 10.1016/0013-9351(77)90034-2.

[8] Jes Fenger. Urban air quality. Atmospheric Environment, 33(29), 1999. doi: 10.1016/ s1352-2310(99)00290-3.

[9] Benjamin G. Ferris. Health effects of exposure to low levels of regulated air pollutants. Journal of the Air Pollution Control Association, 28(5):482–497, May 1978. doi: 10.1080/00022470.1978. 10470621.

[10] J. Firket. Fog along the Meuse Valley. Transactions of the Faraday Society, 32:1192, 1936. doi: 10.1039/tf9363201192.

[11] Milton Friedman. The interpolation of time series by related series. Journal of the American Statistical Association, 57(300):729–757, December 1962. doi: 10.1080/01621459.1962. 10500812.

[12] Gobierno de México, Secretaría de Salud. Dirección general de información en salud. https://www.gob.mx/salud/acciones-y-programas/ direccion-general-de-informacion-en-salud-dgis, July 2020.

[13] Thad Godish, Wayne T. Davis, and Joshua S. Fu. Air quality. Florida, The address, 5 edition, 2015. doi: 10.1201/9781003032595.

[14] Hans W. Gottinger. Air pollution health effects in the Munich metropolitan area: Preliminary results based on a statistical model. Environment International, 9(3):207–220, January 1983. doi: 10.1016/0160-4120(83)90038-7.

[15] Jack D. Hackney, William S. Linn, Ramon D. Buckley, E. Eugene Pedersen, Sarunas K. Karuza, David C. Law, and D. Armin Fischer. Experimental studies on human health effects of air pollutants. Archives of Environmental Health: An International Journal, 30(8):373–378, August 1975. doi: 10.1080/00039896.1975.10666728.

[16] Jack D. Hackney, Frederick C. Thiede, William S. Linn, E. Eugene Pedersen, Charles E. Spier, David C. Law, and D. Armin Fischer. Experimental studies on human health effects of air pollutants. IV. Short-term physiological and dlinical effects of nitrogen dioxide exposure. Archives of Environmental Health: An International Journal, 33(4):176–181, July 1978. doi: 10.1080/00039896.1978.10667330.

[17] Victor Hasselblad. Comparison of methods for the analysis of panel studies. Technical report, Environmental Protection Agency, June 1978. https://nepis.epa.gov/Exe/ZyPDF.cgi?Dockey=2000ZPAO.PDF.

[18] A. Hatzakis, K. Katsouyanni, A. Kalandidi, N. Day, and D. Trichopoulos. Short-term effects of air pollution on mortality in Athens. International Journal of Epidemiology, 15(1):73–81, March 1986. doi: 10.1093/ije/15.1.73.

[19] Steven C. Hillmer and William W. S. Wei. Time series analysis: Univariate and multivariate methods., volume 86. JSTOR, March 1991. doi: 10.2307/2289741.

[20] Rob Hyndman. Forecasting: principles and practice. OTexts, Melbourne, 2018. ISBN 9780987507112.

[21] Daniel Jacob and Darrel Winner. Effect of climate change on air quality. Atmospheric Environment, 43(1), 2009. doi: 10.1016/j.atmosenv.2008.09.051.

[22] Edward L. Korn and Alice S. Whittemore. Methods for analyzing panel studies of acute health effects of air pollution. Biometrics, 35(4):795, December 1979. doi: 10.2307/2530111.

[23] Lester B. Lave and Eugene P. Seskin. An analysis of the association between U.S. mortality and air pollution. Journal of the American Statistical Association, 68(342):284–290, June 1973. doi: 10.1080/01621459.1973.10482421.

[24] P. J. Lawther. Chronic bronchitis and air pollution. Journal (Royal Society of Health), 79(1):4–9, January 1959. doi: 10.1177/146642405907900102.

[25] Michael D. Lebowitz. A comparative analysis of the stimulus-response relationship between mortality and air pollution-weather. Environmental Research, 6(1):106–118, March 1973. doi: 10.1016/0013-9351(73)90023-6.

[26] Lixin Li and Peter Revesz. A comparison of spatio-temporal interpolation methods. In Geographic Information Science, pages 145–160. Springer Berlin Heidelberg, 2002. doi: 10.1007/3-540-45799-2_11.

[27] William S. Linn, Ramon D. Buckley, Charles E. Spier, Raymond L. Blessey, Michael P. Jones, D. Armin Fischer, and Jack D. Hackney. Health effects of ozone exposure in asthmatics. American Review Of Respiratory Disease, 117(5):835–843, May 1978. doi: 10.1164/arrd.1978.117.5.835.

[28] W. P. D. Logan. Mortality in the London fog incident, 1952. Lancet (London, England), 1:336–8, February 1953. doi: 10.1016/s0140-6736(53)91012-5.

[29] M. Maclure. The case-crossover design: A method for studying transient effects on the risk of acute events. American journal of epidemiology, 133:144–53, January 1991. doi: 10.1093/oxfordjournals.aje.a115853.

[30] F. Popescu and I. Guyon. Causality in time series. Microtome PublishingBrookline, 2013.

[31] Secretaría de Salud. Egresos Hospitalarios. Datos abiertos, August 2020. URL http://www.dgis.salud.gob.mx/contenidos/basesdedatos/da_egresoshosp_gobmx.html.

[32] Sistema Integral de Monitoreo Ambiental (SIMA), Nuevo León. aire.nl.gob.mx | Home. http://aire.nl.gob.mx/, 2020. [Accesed 14/may/2020].

[33] William Wei. Time series analysis univariate and multivariate methods. Pearson, Boston, 2019. doi: 10.2307/2289741.

[34] World Health Organization. International statistical classification of diseases and related health problems. World Health Organization, Geneva, 2011. ISBN 9789241548342.

[35] World Health Organization. International classification of diseases. https://www.who.int/classifications/icd/en/, 2020.