

REGULATORY DOSSIER SUBMISSION

Dossier ID: DOS-E1073DDEC9

Country: Burkina Faso

Submission Date: 2026-03-11

Product: Simvvvia (simvastatin) | ATC: C10AA01 | Strength: 20 mg

Applicant: United Therapeutics | Manufacturer: Mwanza Pharma Industries

Policy Label: standard\_review | Risk Score: 0.5

AMR Stewardship: AWaRe=not\_applicable | Unmet Need=not\_applicable | GLASS Trend=not\_applicable  
| Watch Similarity=not\_applicable

SECTION CONTENT

[1] m1\_application\_admin - Application Form and Administrative Information

Labels: presence=present; length=length\_ok; correctness=correct

This dossier constitutes a formal application for Marketing Authorization of Simvvvia (simvastatin) submitted on 2026-03-11 to the national regulatory authority of Burkina Faso. The applicant, United Therapeutics, formally requests approval for the commercial distribution of this medicinal product. The proposed pharmaceutical form is tablet with a strength of 20 mg, classified under ATC code C10AA01. All required administrative declarations, legal attestations, letters of authorization, and regional application forms have been completed, signed by the responsible Qualified Person, and appended to Module 1. Risk mitigation activities are documented with ownership, timelines, and follow-up checkpoints. A comprehensive table of contents and glossary of abbreviations are provided to facilitate regulatory review. All referenced documents are available in the dossier annex and traceable by section i

[1] m1\_manufacturer\_gmp - Manufacturer and GMP Evidence

Labels: presence=present; length=length\_ok; correctness=correct

The primary commercial manufacturing site for the finished pharmaceutical product is Mwanza Pharma Industries, located in Burkina Faso. The facility was subject to a comprehensive GMP inspection by a recognized stringent regulatory authority. The latest inspection concluded on 2024-09-14, resulting in a formal compliance status of 'compliant'. The active GMP certificate number is GMP-444519, which remains valid until 2026-05-12. Site master files, a summary of recent inspection observations, and evidence of completed Corrective and Preventive Actions (CAPA) are included in the annex. All referenced documents are available in the dossier annex and traceable by section identifier. Recent regulatory inspections yielded no critical observations, and all major findings have been effectively closed. Inspection history, CAPA closure records, and site responsibilities were reconciled across the manufacturing chain. Method validation records and quality controls were reviewed for internal consistency. The GMP evidence package includes certificate status, inspection scope, and manufacturing-site accountability records. Data were analyzed using a mixed-effects model for repeated measures, adjusting for ba

Site Inspection History Summary:

| Inspection Date | Authority     | Scope       | Outcome   |
|-----------------|---------------|-------------|-----------|
| 2024-09-14      | National Auth | FPP/General | compliant |
| 2022-05-12      | SRA Joint     | API/Sterile | compliant |

[1] m1\_product\_information - Product Information and Naming

Labels: presence=present; length=length\_ok; correctness=correct

The proposed proprietary name for this medicinal product is Simvvvia, containing the active pharmaceutical ingredient simvastatin. The draft Summary of Product Characteristics (SmPC), Patient Information Leaflet (PIL), and primary/secondary packaging labels are provided for review. These documents detail the approved therapeutic indications, posology, contraindications, special warnings, and precautions for use. A comprehensive risk management plan to minimize medication errors is also submitted. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. Risk mitigation activities are documented with ownership, timelines, and follow-up checkpoints. Stability testing was conducted in climatic zones III and IV, reflecting the intended distribution markets. The analytical procedure was validated in accordance with ICH Q2(R1) guidelines, demonstrating acceptable accuracy, precision, and specificity. All referenced documents are available in the dossier annex and traceable by section identifier. Where applicable, protocol deviations were assessed for impact on interpretation of outcomes. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentration-time profiles. The applicant confirms alignment with applicable technical guidance and regional submission standards. Data were analyzed using a mixed-effects model.

## [2] m2\_quality\_overall\_summary - Quality Overall Summary

Labels: presence=present; length=length\_ok; correctness=correct

This Quality Overall Summary (QOS) provides a critical evaluation of the chemistry, manufacturing, and controls (CMC) data for simvastatin. The control strategy justifies the proposed specifications for both the active substance and the finished product, emphasizing critical quality attributes (CQAs) and critical process parameters (CPPs). Impurity profiles, including degradation products and potential genotoxic impurities, have been thoroughly characterized. Batch analysis data from three commercial-scale validation batches demonstrate consistent manufacturing performance and compliance with all release criteria. The analytical procedure was validated in accordance with ICH Q2(R1) guidelines, demonstrating acceptable accuracy, precision, and specificity. The risk management plan includes routine pharmacovigilance and proposed educational materials for healthcare professionals. The applicant confirms alignment with applicable technical guidance and regional submission standards. The manufacturing site operates under a fully implemented Pharmaceutical Quality System (PQS) aligned with ICH Q10. Method validation records and quality controls were reviewed for internal consistency. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentration-time profiles. Adverse events were coded using the Medical Dictionary for Regulatory Activities (MedDRA) version 24.0. Stability testing was conducted in climatic zones III and IV, reflecting the intended distribution markets. All referenced documents are available in the dossier annex and traceable by section identifier. Statistical significance was set at an alpha level of 0.05, and all tests were two-sided. Data were analyzed using a mixed-effects model for repeated measures, adjusting for baseline covariates. Risk mitigation activities are documented with ownership, timelines, and follow-up checkpoints. Where applicable, protocol deviations were assessed for impact on interpretation of outcomes. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. The manufacturing site operates under a fully implemented Pharmaceutical Quality System (PQS) aligned with ICH Q10. Statistical significance was set at an alpha level of 0.05, and all tests were two-sided. Stability testing was conducted in climatic zones III and IV, reflectin

[2] m2\_clinical\_overview - Clinical Overview and Benefit-Risk Summary

Labels: presence=present; length=length\_ok; correctness=correct

The Clinical Overview synthesizes the efficacy and safety data supporting the use of Simvvia for the treatment of bacterial skin infection. The pivotal clinical development program comprised 2 well-controlled studies. Based on the integrated analysis, the reported clinical outcome category is 'endpoint\_met'. The overall benefit-risk profile is considered highly favorable for the target patient population. Safety findings were consistent with the known pharmacological class effects, and appropriate risk minimization measures have been integrated into the proposed prescribing information. Where applicable, protocol deviations were assessed for impact on interpretation of outcomes. Stability testing was conducted in climatic zones III and IV, reflecting the intended distribution markets. Clinical interpretation focuses on whether efficacy, safety, and benefit-risk conclusions are supported by the submitted evidence. Benefit-risk language was reviewed for consistency with the submitted efficacy and safety summaries. The manufacturing site operates under a fully implemented Pharmaceutical Quality System (PQS) aligned with ICH Q10. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. Risk mitigation activities are documented with ownership, timelines, and follow-up checkpoints. Data were analyzed using a mixed-effects model for repeated measures, adjusting for baseline covariates. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. The target population demographics are broadly representative of the intended commercial patient cohort. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentration-time profiles. The risk management plan includes routine pharmacovigilance and proposed educational materials for healthcare professionals. No new safety signals were identified during the extended ope

[3] m3\_api\_quality - API Quality and Control Strategy

Labels: presence=present; length=length\_ok; correctness=correct

The active pharmaceutical ingredient (API) section details the complete synthesis route, starting from well-defined regulatory starting materials. Robust in-process controls and intermediate specifications ensure the consistent quality of the final drug substance. The analytical procedures used for release and stability testing have been fully validated for accuracy, precision, linearity, and robustness. The impurity control strategy adequately addresses organic impurities, residual solvents, and elemental impurities in accordance with current ICH guidelines. Forced degradation studies confirm the stability-indicating nature of the assay methods. Batch analysis, validation, and impurity management records support the proposed API quality framework. The analytical procedure was validated in accordance with ICH Q2(R1) guidelines, demonstrating acceptable accuracy, precision, and specificity. All referenced documents are available in the dossier annex and traceable by section identifier. Residual solvents and elemental impurities are controlled well below ICH Q3C and Q3D permissible daily exposures. Method validation records and quality controls were reviewed for internal consistency. Risk mitigation activities are documented with ownership, timelines, and follow-up checkpoints. Stability testing was conducted in climatic zones III and IV, reflecting the intended distribution markets. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. The risk management plan includes routine pharmacovigilance and proposed educational materials for healthcare professionals. Data were analyzed using a mixed-effects model for repeated measures, adjusting for baseline covariates. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentration-time profiles. Where applicable, protocol deviations were assessed for impact on interpretation of outcomes. The applicant confirms alignment with

applicable technical guidance and regional submission standards. Statistical significance was set at an alpha level of 0.05, and all tests were two-sided. The manufacturing site operates under a fully implemented Pharmaceutical Quality System (PQS) aligned with ICH Q10. Adverse events were coded using the Medical Dictionary for Regulatory Activities (MedDRA) version 24.0. The API package links specification limits to validated analytical methods and impurity-control decisions. Critical material attributes and control strategy elements were reviewed for consistency across the quality dossier. The synthetic route involves three well-characterized steps with defined starting materials and isolated intermediates. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentration-time profiles. The risk management plan includes routine pharmacovigilance and proposed educational materials for healthcare professionals. Method validation records and quality controls were reviewed for internal consistency. P

### [3] m3\_fpp\_manufacturing - FPP Manufacturing Process and Controls

Labels: presence=present; length=length\_ok; correctness=correct

The finished pharmaceutical product (FPP) manufacturing process utilizes standard, scalable unit operations. A formal quality risk assessment (e.g., FMEA) was conducted to identify critical process parameters (CPPs) that impact critical quality attributes (CQAs). Process performance qualification (PPQ) reports for three consecutive commercial-scale batches verify that the process is maintained in a state of control. In-process controls, intermediate hold-time studies, and packaging validation data are presented to substantiate the robustness and reproducibility of the commercial manufacturing operations. Statistical significance was set at an alpha level of 0.05, and all tests were two-sided. Adverse events were coded using the Medical Dictionary for Regulatory Activities (MedDRA) version 24.0. The analytical procedure was validated in accordance with ICH Q2(R1) guidelines, demonstrating acceptable accuracy, precision, and specificity. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentration-time profiles. The applicant confirms alignment with applicable technical guidance and regional submission standards. The risk management plan includes routine pharmacovigilance and proposed educational materials for healthcare professionals. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. Stability testing was conducted in climatic zones III and IV, reflecting the intended distribution markets. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. Risk mitigation activities are documented with ownership, timelines, and follow-up checkpoints. Where applicable, protocol deviations were assessed for impact on interpretation of outcomes. Method validation records and quality controls were reviewed for internal consistency.

### [3] m3\_stability - Stability and Shelf-Life Justification

Labels: presence=present; length=length\_ok; correctness=correct

A comprehensive stability program has been executed to justify the proposed shelf-life and storage conditions. Data from both accelerated (40°C/75% RH for 6 months) and long-term storage conditions (spanning up to 36 months) are presented for multiple primary stability batches. Statistical evaluation using linear regression and poolability tests confirms that degradation trends remain well within the acceptable specification limits. Photostability and freeze-thaw cycling studies demonstrate that the product does not require specific handling precautions. A post-approval stability protocol commitment is included. The applicant confirms alignment with applicable technical guidance and regional submission standards. Method validation records and quality controls were reviewed for internal consistency. Risk mitigation activities are documented with ownership, timelines, and follow-up checkpoints. Where applicable, protocol deviations were assessed for impact on interpretation of outcomes. Stability testing was

conducted in climatic zones III and IV, reflecting the intended distribution markets. Data were analyzed using a mixed-effects model for repeated measures, adjusting for baseline covariates. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. Adverse events were coded using the Medical Dictionary for Regulatory Activities (MedDRA) version 24.0. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. All referenced documents are available in the dossier annex and traceable by section identifier. The manufacturing site operates under a fully implemented Pharmaceutical Quality System (PQS) aligned with ICH Q10. The risk management plan includes routine pharmacovigilance and proposed educational materials for healthcare professionals

[4] m4\_nonclinical\_summary - Nonclinical Study Summary

Labels: presence=present; length=length\_ok; correctness=correct

The nonclinical testing program evaluated the primary pharmacodynamics, secondary pharmacodynamics, safety pharmacology, and comprehensive toxicology profile of the compound. Repeat-dose toxicity studies in two mammalian species established clear no-observed-adverse-effect levels (NOAELs). Genotoxicity testing (Ames and in vivo micronucleus assays) yielded negative results. Reproductive and developmental toxicity studies revealed no teratogenic signals at clinically relevant exposures. The calculated safety margins support the proposed clinical dosing regimen without major safety concerns. The analytical procedure was validated in accordance with ICH Q2(R1) guidelines, demonstrating acceptable accuracy, precision, and specificity. Data were analyzed using a mixed-effects model for repeated measures, adjusting for baseline covariates. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer

[5] m5\_trial\_listing - Tabular Listing of Clinical Studies

Labels: presence=present; length=length\_ok; correctness=correct

The tabular listing of clinical studies provides a comprehensive inventory of all trials conducted in support of the proposed indication (bacterial skin infection). This includes Phase I pharmacokinetic/pharmacodynamic studies, Phase II dose-finding studies, and the pivotal Phase III efficacy and safety trials. For each study, the table details the protocol identifier, study design, randomization ratio, number of enrolled subjects, study duration, and current completion status. Statistical significance was set at an alpha level of 0.05, and all tests were two-sided. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. All referenced documents are available in the dossier annex and traceable by section identifier. Data were analyzed using a mixed-effects model for repeated measures, adjusting for baseline covariates. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. The analytical procedure was validated in accordance with ICH Q2(R1) guidelines, demonstrating acceptable accuracy, precision, and specificity. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentration-time profiles. Method validation records and quality controls were reviewed for internal consistency. The risk management plan includes routine pharmacovigilance and proposed educational materials for healthcare professionals. Stability testing was conducted in climatic zones III and IV, reflecting the intended distribution marke

Pivotal Clinical Trial Inventory:

| Protocol ID | Phase | Enrollment | Status    |
|-------------|-------|------------|-----------|
| PROT-001    | III   | 404        | completed |
| PROT-002    | III   | 655        | completed |

[5] m5\_pivotal\_trial\_reports - Pivotal Clinical Trial Reports

Labels: presence=present; length=length\_ok; correctness=correct

Detailed clinical study reports (CSRs) for the pivotal efficacy trials are provided. These randomized, double-blind, actively-controlled trials evaluated the primary and secondary endpoints in accordance with the pre-specified statistical analysis plan (SAP). The trials resulted in a formal outcome category of 'endpoint\_met', demonstrating robust treatment effects in the intent-to-treat (ITT) and per-protocol populations. Subgroup analyses across age, gender, and baseline disease severity strata were consistent with the primary findings. Serious adverse events (SAEs) and discontinuations were systematically analyzed and adjudicated. Discontinuations due to treatment-emergent adverse events (TEAEs) were low and balanced across all study arms. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. Risk mitigation activities are documented with ownership, timelines, and follow-up checkpoints. The primary efficacy endpoint was met, demonstrating a statistically significant improvement over placebo ( $p < 0.001$ ). The risk management plan includes routine pharmacovigilance and proposed educational materials for healthcare professionals. Statistical significance was set at an alpha level of 0.05, and all tests were two-sided. The analytical procedure was validated in accordance with ICH Q2(R1) guidelines, demonstrating acceptable accuracy, precision, and specificity. Method validation records and quality controls were reviewed for internal consistency. Clinical study reports document protocol adherence, endpoint handling, and safety interpretation in the target population. Where applicable, protocol deviations were assessed for impact on interpretation of outcomes. The manufacturing site operates under a fully implemented Pharmaceutical Quality System (PQS) aligned with ICH Q10. The applicant confirms alignment with applicable technical guidance and regional submission standards. Stability testing was conducted in climatic zones III and IV, reflecting the intended distribution markets. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. Adverse events were coded using the Medical Dictionary for Regulatory Activities (MedDRA) version 24.0. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentr

[5] m5\_bioequivalence - Biopharmaceutics and Bioequivalence Evidence

Labels: presence=present; length=length\_ok; correctness=correct

Biopharmaceutics data substantiate the formulation development bridging between clinical trial materials and the proposed commercial product. In vivo bioequivalence studies demonstrated that the 90% confidence intervals for the geometric mean ratios of C<sub>max</sub> and AUC parameters fell entirely within the standard acceptance criteria of 80.00% to 125.00%. Additionally, multi-point comparative dissolution profiles across physiological pH ranges (pH 1.2, 4.5, and 6.8) confirmed similarity ( $f_2 > 50$ ). The bioanalytical methods used for plasma quantification were fully validated. All referenced documents are available in the dossier annex and traceable by section identifier. Stability testing was conducted in climatic zones III and IV, reflecting the intended distribution markets. A randomized, double-blind, parallel-group study design was employed to minimize selection and observer bias. The manufacturing site operates under a fully implemented Pharmaceutical Quality System (PQS) aligned with ICH Q10. Data were analyzed using a mixed-effects model for repeated measures, adjusting for baseline covariates. The applicant confirms alignment with applicable technical guidance and regional submission standards. Where applicable, protocol deviations were assessed for impact on interpretation of outcomes. Pharmacokinetic parameters were derived using non-compartmental analysis from the concentration-time profiles. The container closure system consists of PVC/PVDC-Aluminum blisters, compliant with pharmacopoeial standards. The analytical procedure was validated in accordance with ICH Q2(R1) guidelines, demonstrating acceptable accuracy, precision, and

specificity. Adverse events were coded using the Medical Dictionary for Regula