

# HERMITIAN MODULAR FORMS FOR FIELDS OF LOW DISCRIMINANT

DIPLOMA THESIS  
in Mathematics

by  
Albert Zeyer

submitted to the  
Faculty of Mathematics, Computer Science and Natural Science of  
RWTH Aachen University

October 2012  
revised version from January 23, 2013

Supervisor: Prof. Dr. Aloys Krieg  
Second examiner: Dr. Martin Raum

written at the  
Lehrstuhl A für Mathematik  
Prof. Dr. A. Krieg

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Preliminaries</b>	<b>4</b>
2.1	Siegel modular forms . . . . .	5
2.2	Elliptic modular forms . . . . .	5
2.3	Hermitian modular forms . . . . .	6
<b>3</b>	<b>Theory</b>	<b>8</b>
<b>4</b>	<b>Implementation</b>	<b>10</b>
<b>5</b>	<b>Conclusion</b>	<b>11</b>
<b>6</b>	<b>References</b>	<b>12</b>

# Chapter 1

## Introduction

We develop an algorithm to compute Fourier expansions of Hermitian modular forms of degree 2 over  $\mathrm{Sp}_2(\mathcal{O})$  for  $\mathcal{O} \subseteq \mathbb{Q}(\sqrt{-\Delta})$ ,  $\Delta \in \{3, 4, 8\}$ .

In [PY07], spaces of Siegel modular cusp forms are calculated.

A similar algorithm is also [Rau12, Algorithm 4.3] for Jacobi forms.

We are doing the same for Hermitian modular forms.

## Chapter 2

### Preliminaries

$\mathbb{N}$  denotes the set  $\{1, 2, 3, \dots\}$ ,  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$  and  $\mathbb{Z}$  are all **integers**.  $\mathbb{Q}$  are all the **rational numbers**,  $\mathbb{R}$  are the **real numbers** and  $\mathbb{C}$  are the **complex numbers**.  $\mathbb{R}^+ := \{x \in \mathbb{R} \mid x > 0\}$ ,  $\mathbb{R}^\times$  and  $\mathbb{C}^\times$  denotes all non-zero numbers.

Let  $\text{Mat}_n(R)$  be the set of all  $n \times n$  **matrices** over some commutative ring  $R$ . Likewise,  $\text{Mat}_n^T(R)$  are the **symmetric**  $n \times n$  matrices.  $X^T$  is the **transposed** matrix of  $X \in \text{Mat}_n(R)$ .  $\bar{Z}$  is the **conjugated** matrix of  $Z \in \text{Mat}_n(\mathbb{C})$ . A matrix  $Y \in \text{Mat}_n(\mathbb{C})$  is greater 0 if and only if  $\forall x \in \mathbb{C}^n - \{0\} : Y[x] := \bar{x}^T Y x \in \mathbb{R}^+$ . Such symmetric matrices are called the **positive definitive matrices**, defined by

$$\mathcal{P}_n(R) = \{X \in \text{Mat}_n^T(R) \mid X > 0\}.$$

For  $A, X \in \text{Mat}_n(\mathbb{C})$ , we define  $A[X] := \bar{X}^T A X$ . The **denominator** of a matrix  $Z \in \text{Mat}_n(\mathbb{Q})$  is the smallest number  $x \in \mathbb{N}$  such that  $xZ \in \text{Mat}_n(\mathbb{Z})$ .

The **general linear group** is defined by

$$\text{GL}_n(R) = \{X \in \text{Mat}_n(R) \mid \det(X) \text{ is a unit in } R\}$$

and the **special linear group** by

$$\text{SL}_n(R) = \{X \in \text{Mat}_n(R) \mid \det(X) = 1\}.$$

The **orthogonal group** is defined by

$$\text{O}_n(R) = \{X \in \text{GL}_n(R) \mid X^T 1_n X = 1_n\}.$$

For  $R \subseteq \mathbb{C}$ ,  $\bar{R} \subseteq R$ , the set of **Hermitian matrices** in  $R$  is defined as

$$\text{Her}_n(R) = \left\{ Z \in \text{Mat}_n(R) \mid \bar{Z}^T = Z \right\}.$$

The **symplectic group** is defined by

$$\text{Sp}_n(R) = \left\{ X \in \text{GL}_{2n}(R) \mid \bar{X}^T J_n X = J_n \right\}$$

where  $J_n := \begin{pmatrix} 0 & 1_n \\ -1_n & 0 \end{pmatrix} \in \text{SL}_{2n}(R)$ .  $\text{Sp}_n(R)$  is also called the **unitary group**. For  $Z \in \text{Mat}_n(\mathbb{C})$ , we call

$$\Re(Z) = \frac{1}{2} \left( Z + \bar{Z}^T \right) \in \text{Mat}_n(\mathbb{C})$$

the **real** part and

$$\Im(Z) = \frac{1}{2i} (Z - \overline{Z}^T) \in \text{Mat}_n(\mathbb{C})$$

the **imaginary** part of  $Z$  and we have  $Z = \Re(Z) + i\Im(Z)$ . Note that we usually have  $\Re(Z), \Im(Z) \notin \text{Mat}_n(\mathbb{R})$  but we have  $\Re(Z), \Im(Z) \in \text{Her}_n(\mathbb{C})$ .

We say that some function  $f: \mathcal{A} \rightarrow \mathcal{B}$  with  $\mathcal{A} \subseteq \text{Mat}_n(R), \mathcal{B} \subseteq R$  is  **$k$ -invariant** under some  $\mathcal{X} \subseteq \text{Mat}_n(R)$  where  $\mathcal{A}[\mathcal{X}] \subseteq \mathcal{A}$  if and only if  $\det(U)^k f(T[U]) = f(T)$  for all  $T \in \mathcal{A}, U \in \mathcal{X}$ .

Let  $S$  be a set with  $G$ -action. Then the set of  $G$ -invariants  $S^G$  is the set of all  $s \in S$  satisfying  $gs = s$  for all  $G$ . We can equip the set of functions  $\mathcal{F} \rightarrow \mathbb{C}$  with the action  $(gf)(T) = \det(g)^k f(T[g])$  and this lead to the definition that we need.

## 2.1 Siegel modular forms

Let  $\mathcal{H}_n := \{Z \in \text{Mat}_n^T(\mathbb{C}) \mid \Im(Z) > 0\}$  be the **Siegel upper half space**. Thus,  $\mathcal{H}_1$  is the **Poincaré upper half plane**. We call  $\text{Sp}_n(\mathbb{Z})$  the **Siegel modular group**.

A **Siegel modular cusp form** of degree  $n \in \mathbb{N}$  for some  $\Gamma \subseteq \text{Sp}_n(\mathbb{Z}), \Gamma$  subgroup of  $\text{Sp}_n(\mathbb{Z})$ , is a holomorphic function

$$f: \mathcal{H}_n \rightarrow \mathbb{C}$$

with

- (1)  $f|_k y = f \quad \forall y \in \Gamma$
- (2) for  $n = 1$ :  $f(Z) = O(1)$  for  $Z \rightarrow i\infty$

where

$$\left( f|_k \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right) (Z) = f((AZ + B)(CZ + D)^{-1}) \cdot \det(CZ + D)^{-k}$$

with  $Z \in \mathcal{H}_n, \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma$ .

$M_k^{\mathcal{H}_n}(\Gamma)$  denotes the vector space of such Siegel modular forms.

## 2.2 Elliptic modular forms

We define

$$\Gamma_0(l) := \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \text{Sp}_n(\mathbb{Z}) \mid C \equiv 0 \pmod{l} \right\} \subseteq \text{Sp}_n(\mathbb{Z})$$

as a subgroup of  $\mathrm{Sp}_n(\mathbb{Z})$ .

**Elliptic modular forms** are Siegel modular cusp forms of degree 1 over  $\Gamma_0(l)$  for some  $l \in \mathbb{N}$ .

$M_k(\Gamma)$  denotes the vector space of such Elliptic modular forms.

### 2.3 Hermitian modular forms

Let  $\mathbb{H}_n := \{Z \in \mathrm{Mat}_n(\mathbb{C}) \mid \Im(Z) > 0\}$  be the **Hermitian upper half space**. Note that these matrices are not symmetric as  $\mathcal{H}_n$  but we have  $\mathcal{H}_n \subseteq \mathbb{H}_n$  and  $\mathcal{H}_1 = \mathbb{H}_1 \subseteq \mathbb{C}$ .

Let  $\Delta \in \mathbb{N}$  so that we have the imaginary quadratic number field  $\mathbb{Q}(\sqrt{-\Delta})$  where  $-\Delta$  is the fundamental discriminant. Then, let  $\mathcal{O} \subseteq \mathbb{Q}(\sqrt{-\Delta})$  be the maximum order. We call  $\mathrm{Sp}_n(\mathcal{O})$  the **Hermitian modular group**. Let  $\Gamma$  be a subgroup of  $\mathrm{Sp}_n(\mathcal{O})$ . Let  $\nu: \Gamma \rightarrow \mathbb{C}^\times$  be an abel character of  $\mathrm{Sp}_n(\mathcal{O})$ .

A **Hermitian modular form** of degree  $n \in \mathbb{N}$  with weight  $k \in \mathbb{Z}$  over  $\Gamma$  and  $\nu$  is a holomorphic function

$$f: \mathbb{H}_n \rightarrow \mathbb{C}$$

with

- (1)  $f(M \cdot Z) = \nu(M) \det(CZ + D)^k f(Z)$ ,  $M = \begin{pmatrix} * & * \\ C & D \end{pmatrix} \in \Gamma, Z \in \mathbb{H}_n$ ,
- (2) for  $n = 1$ :  $f$  is holomorphic in all cusps.

$M_k^{\mathbb{H}_n}(\Gamma, \nu)$  denotes the vector space of such Hermitian modular forms.

In this work, we will concentrate on Hermitian modular forms of degree 2. We will start with  $\Delta \in \{3, 4, 8\}$ .

Because  $-\Delta$  is fundamental, we have two possible cases:

- 1.  $\Delta \equiv 3 \pmod{4}$  and  $\Delta$  is square-free, or
- 2.  $\Delta \equiv 0 \pmod{4}$ ,  $\Delta/4 \equiv 1, 2 \pmod{4}$  and  $\Delta/4$  is square-free.

And for the maximum order  $\mathcal{O}$ , we have (compare [Der01])

$$\begin{aligned} \mathcal{O} &= \mathbb{Z} + \mathbb{Z} \frac{-\Delta + i\sqrt{\Delta}}{2}, \\ \mathcal{O}^\# &= \mathbb{Z} \frac{i}{\sqrt{\Delta}} + \mathbb{Z} \frac{1 + i\sqrt{\Delta}}{2}. \end{aligned}$$

From now on, we will always work with Hermitian modular forms of degree 2, i.e. we will always have  $n = 2$ , except if otherwise stated.

## Chapter 3

### Theory

**Lemma 3.1.** *Let  $f: \mathbb{H}_2 \rightarrow \mathbb{C}$  be a Hermitian modular form of weight  $k$ . Let  $S \in \mathcal{P}_2(\mathbb{C})$ . Then,  $f(S\tau): \mathbb{H}_1 \subseteq \mathbb{C} \rightarrow \mathbb{C}$  is an elliptic modular form of weight  $2k$  to  $\Gamma_0(l)$ , where  $l$  is the denominator of  $S^{-1}$ .*

**Lemma 3.2.** *Prop 7.3. von Poor für herm Modulformen.  $\Gamma(\mathcal{L}) \supseteq \Gamma_0(l)$  for  $l \in \mathbb{Z}^+$ ,  $ls^{-1} \in \mathcal{P}_n(\mathcal{O})$ .*

We want to calculate a generating set for the Fourier expansions of Hermitian modular forms. Now we will formulate the main algorithm of our work.

**Algorithm 3.3.** We have the Hermitian modular form degree  $n = 2$  fixed, as well as some  $\Delta$  (for now,  $\Delta \in \{3, 4, 8\}$ ). Then we select some form weight  $k \in \mathbb{Z}$  ( $k \in \{1, \dots, 20\}$  or so), some  $\mathcal{O} \subseteq \mathbb{Q}(\sqrt{-\Delta})$  and some subgroup  $\Gamma$  of  $\mathrm{Sp}_2(\mathcal{O})$ . Then we select an abel character  $\nu: \Gamma \rightarrow \mathbb{C}^\times$  of  $\mathrm{Sp}_2(\mathcal{O})$ .

We define the index set

$$\Lambda := \left\{ 0 \leq \begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} \in \mathrm{Mat}_2(\mathcal{O}^\#) \mid a, c \in \mathbb{Z} \right\}.$$

Fix  $B \in \mathbb{N}$  as a limit. Select a precision

$$\mathcal{F} := \left\{ \begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} \mid 0 \leq a, c < B, b \in \mathcal{O}^\# \right\} \subseteq \Lambda.$$

1. We start with  $l = 1$  and increase it but only use the square-free numbers.
2. Set  $\mathcal{S} = \{\}$ ,
3. Enumerate matrices  $S \in \mathrm{Mat}_2^T(\mathbb{Z})$ , and set  $\mathcal{S} \leftarrow \mathcal{S} \cup \{S\}$  and for each time you add a new matrix perform the following steps.



4. We set

$$\mathcal{M}_{k,S,\mathcal{F}}^H := \{(a[S])_{S \in \mathcal{S}} \mid a \in \mathbb{Q}^{\mathcal{F}} \text{ is } \mathrm{GL}_2(\mathcal{O}) \text{ invariant}\} \subseteq \bigoplus_{S \in \mathcal{S}} \mathbb{Q}^{\mathcal{F}(S)},$$

where

$$a[S] := \mathbb{H}_1 \rightarrow \mathbb{Q}, \tau \mapsto a(S\tau),$$

The elements  $a$  are the Fourier expansions, thus  $a(T) \in \mathbb{Q}$  for  $T \in \mathcal{F}$  are the Fourier coefficients. Recall that  $a$  being invariant under  $\mathrm{GL}_2(\mathcal{O})$  means that we have

$$\det(U)^k a(T[U]) = a(T) \quad \forall U \in \mathrm{GL}_2(\mathcal{O}).$$

Thus, to represent such invariant  $a$ , we can use `¡TODO¿`.

And we set

$$\mathcal{M}_{k,S,\mathcal{F}} := \bigoplus_{S \in \mathcal{S}} \mathcal{FE}_{\mathcal{F}(S)}(\mathrm{M}_k(\Gamma_0(l_S)))$$

where  $\mathrm{M}_k(\Gamma_0(l_S))$  is the vectorspace of Elliptic modular forms over  $\Gamma_0(l_S)$ .

5. If

$$\dim \mathcal{M}_{k,S,\mathcal{F}}^H \cap \mathcal{M}_{k,S,\mathcal{F}} = \dim \mathrm{M}_k^{\mathbb{H}^2}(\Gamma, \nu),$$

then we are ready and we can reconstruct the Fourier expansion in the following way:

...

If not, then return to step 3, and enlarge  $\mathcal{S}$ .

## **Chapter 4**

### **Implementation**

In this chapter, we are describing the implementation.

## **Chapter 5**

## **Conclusion**

Blub

## Chapter 6

### References

- [Der01] T. Dern. *Hermiteische Modulformen zweiten Grades*. Mainz, 2001.
- [PY07] C. Poor and D.S. Yuen. Computations of spaces of siegel modular cusp forms. *Journal of the Mathematical Society of Japan*, 59(1):185–222, 2007.
- [Rau12] M. Raum. Computing Jacobi Forms and Linear Equivalences of Special Divisors. *ArXiv e-prints*, December 2012.