

# HERMITIAN MODULAR FORMS FOR FIELDS OF LOW DISCRIMINANT

DIPLOMA THESIS  
in Mathematics

by  
Albert Zeyer

submitted to the  
Faculty of Mathematics, Computer Science and Natural Science of  
RWTH Aachen University

October 2012  
revised version from December 14, 2012

Supervisor: Prof. Dr. Aloys Krieg  
Second examiner: Dr. Martin Raum

written at the  
Lehrstuhl A für Mathematik  
Prof. Dr. A. Krieg

**Contents**

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Background results</b>	<b>4</b>
2.1	Preliminaries . . . . .	4
<b>3</b>	<b>Theory</b>	<b>5</b>
<b>4</b>	<b>Implementation</b>	<b>6</b>
<b>5</b>	<b>Conclusion</b>	<b>7</b>
<b>6</b>	<b>References</b>	<b>8</b>

# Chapter 1

## Introduction

In [PY07], spaces of Siegel modular cusp forms are calculated.

We are doing the same for hermitian modular forms.

## Chapter 2

### Background results

#### 2.1 Preliminaries

Let  $M_n(\mathcal{K})$  be the set of all  $n \times n$  matrices over some field  $\mathcal{K}$ . Likewise,  $M_n^T(\mathcal{K})$  are the symmetric  $n \times n$  matrices. A matrix  $Y \in M_n(\mathbb{R})$  is greater 0 iff  $\forall x \in \mathbb{R}^n - \{0\} : Y[x] := x^T Y x > 0$ . Let  $\mathbb{H}_n := \{Z = X + iY \in M_n^T(\mathbb{C}) \mid Y > 0\}$ . Thus,  $\mathbb{H}_1$  is the Poincaré upper half plane.

The general linear group is defined by  $GL_n(\mathcal{K}) = \{X \in M_n(\mathcal{K}) \mid \det(X) \neq 0\}$  and the special linear group by  $SL_n(\mathcal{K}) = \{X \in M_n(\mathcal{K}) \mid \det(X) = 1\}$ . The orthogonal group is defined by  $O_n(\mathcal{K}) = \{X \in GL_n(\mathcal{K}) \mid X^T X = 1_n\}$ . The symplectic group is defined by  $Sp_n(\mathcal{K}) = \{X \in GL_{2n}(\mathcal{K}) \mid X^T J_n X = J_n\}$  where  $J_n := \begin{pmatrix} 0 & 1_n \\ -1_n & 0 \end{pmatrix} \in SL_{2n}(\mathcal{K})$ .

A siegel modular cusp form is a holomorphic function

$$f: \mathbb{H}_n \rightarrow \mathbb{C}$$

with

- (1)  $f|_k y = f \forall y \in \Gamma \subseteq Sp_n(\mathbb{Z})$
- (2) for  $n = 1$ :  $f(Z) = O(1)$  for  $Z \rightarrow i\infty$

where

$$\left( f|_k \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right) (Z) = f((AZ + B)(CZ + D)^{-1}) \cdot \det(CZ + D)^{-k}$$

with  $Z = S\tau$ .

## Chapter 3

### Theory

**Lemma 3.1.** *Let  $f: \mathbb{M}_2(\mathbb{C}) \rightarrow \mathbb{C}$  be a hermitian modular form of weight  $k$ . Then,  $f(S\tau): \mathbb{H}_1 \subseteq \mathbb{C} \rightarrow \mathbb{C}$  is an elliptic modular form of weight  $2k$  for some matrix  $S \in \mathbb{M}_2(\mathbb{Z})$  with  $\Gamma(S) \subseteq \mathrm{SL}_2(\mathbb{Z})$ .*

**Algorithm 3.2.** 1. Select a set of matrices  $\mathcal{S} \subseteq \mathbb{M}_2^T(\mathbb{Z})$  with  $0 < S \in \mathcal{S}$ . Make  $\mathcal{S}$  big enough. Now, for some  $S \in \mathcal{S}$ :

2. Fix  $B \in \mathbb{N}$  as a limit. Or select a precision

$$\mathcal{F} = \left\{ \begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} \middle| 0 \leq ac < B \right\} \subseteq \Lambda,$$

where

$$\Lambda := \left\{ 0 \leq \begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} \in \mathbb{M}_2(o^\#) \middle| a, c \in \mathbb{Z} \right\}.$$

3.

$$\mathcal{M}_{k,S,\mathcal{F}}^H = \{fS \mid f \in \mathbb{Q}^{\mathcal{F}}\},$$

$$\mathcal{M}_{k,S} = \mathrm{FourierExpansion}_{\mathcal{F}(S)}(\mathbb{M}_k(\Gamma(S)))$$

4. If

$$\dim \mathcal{M}_{k,S,\mathcal{F}}^H \cap \bigoplus_{S \in \mathcal{S}} \mathcal{M}_{k,S} = \dim M_k^H,$$

then we are ready and we can reconstruct the Fourier expansion in the following way:

...

## **Chapter 4**

### **Implementation**

In this chapter, we are describing the implementation.

## Chapter 5

## Conclusion

Blub

## Chapter 6

### References

- [PY07] C. Poor and D.S. Yuen. Computations of spaces of siegel modular cusp forms.  
*Journal of the Mathematical Society of Japan*, 59(1):185–222, 2007.