

HERMITIAN MODULAR FORMS FOR FIELDS OF LOW DISCRIMINANT

DIPLOMA THESIS
in Mathematics

by
Albert Zeyer

submitted to the
Faculty of Mathematics, Computer Science and Natural Science of
RWTH Aachen University

October 2012
revised version from December 18, 2012

Supervisor: Prof. Dr. Aloys Krieg
Second examiner: Dr. Martin Raum

written at the
Lehrstuhl A für Mathematik
Prof. Dr. A. Krieg

Contents

1	Introduction	3
2	Background results	4
2.1	Preliminaries	4
3	Theory	6
4	Implementation	7
5	Conclusion	8
6	References	9

Chapter 1

Introduction

We develop an algorithm to compute Fourier expansions of Hermitian Modular Forms of degree 2 over $\mathrm{Sp}_2(\mathcal{O})$ for $\mathcal{O} \subseteq \mathbb{Q}(\sqrt{-\Delta})$, $\Delta \in \{3, 4, 8\}$.

In [PY07], spaces of Siegel Modular Cusp forms are calculated.

A similar algorithm is also [Rau12, Algorithm 4.3] for Jacobi forms.

We are doing the same for Hermitian Modular forms.

Chapter 2

Background results

2.1 Preliminaries

Let $M_n(\mathbb{K})$ be the set of all $n \times n$ matrices over some field \mathbb{K} . Likewise, $M_n^T(\mathbb{K})$ are the symmetric $n \times n$ matrices. A matrix $Y \in M_n(\mathbb{R})$ is greater 0 iff $\forall x \in \mathbb{R}^n - \{0\} : Y[x] := x^T Y x > 0$. Let $\mathbb{H}_n := \{Z = X + iY \in M_n^T(\mathbb{C}) \mid Y > 0\}$. Thus, \mathbb{H}_1 is the Poincaré upper half plane.

The general linear group is defined by $GL_n(\mathbb{K}) = \{X \in M_n(\mathbb{K}) \mid \det(X) \neq 0\}$ and the special linear group by $SL_n(\mathbb{K}) = \{X \in M_n(\mathbb{K}) \mid \det(X) = 1\}$. The orthogonal group is defined by $O_n(\mathbb{K}) = \{X \in GL_n(\mathbb{K}) \mid X^T 1_n X = 1_n\}$. The symplectic group is defined by $Sp_n(\mathbb{K}) = \{X \in GL_{2n}(\mathbb{K}) \mid X^T J_n X = J_n\}$ where $J_n := \begin{pmatrix} 0 & 1_n \\ -1_n & 0 \end{pmatrix} \in SL_{2n}(\mathbb{K})$. $Sp_n(\mathbb{K})$ will also be called the Hermitian modular group.

A **Siegel modular cusp form** of degree $n \in \mathbb{N}$ for some $\Gamma \subseteq Sp_n(\mathbb{Z})$ is a holomorphic function

$$f: \mathbb{H}_n \rightarrow \mathbb{C}$$

with

- (1) $f|_k y = f \quad \forall y \in \Gamma$
- (2) for $n = 1$: $f(Z) = O(1)$ for $Z \rightarrow i\infty$

where

$$\left(f|_k \begin{pmatrix} A & B \\ C & D \end{pmatrix}\right)(Z) = f((AZ + B)(CZ + D)^{-1}) \cdot \det(CZ + D)^{-k}$$

with $Z \in \mathbb{H}_n$, $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma$.

A **Hermitian modular form** of degree $n \in \mathbb{N}$ is a holomorphic function

$$f: \mathbb{H}_n \rightarrow \mathbb{C}$$

with weight $k \in \mathbb{Z}$ for some $\Gamma \subseteq Sp_n(\mathcal{O})$, $\mathcal{O} \subseteq \mathbb{Q}(\sqrt{-\Delta})$, $\Delta \in \mathbb{N}$, $\nu: \Gamma \rightarrow \mathbb{C}^\times$ is an abel character of $Sp_n(\mathcal{O})$, with

- (1) $f(M \cdot Z) = \nu(M) \det(CZ + D)^k f(Z)$, $M = \begin{pmatrix} * & * \\ C & D \end{pmatrix} \in \Gamma$, $Z \in \mathbb{H}_n$,
- (2) for $n = 1$: f is holomorphic in all peaks.

$[\Gamma, k, \nu]$ denotes the vector space of such hermitian modular forms.

In this work, we will concentrate on Hermitian Modular forms of degree 2. We will start with $\Delta \in \{3, 4, 8\}$.

Note that if Δ is fundamental, we have

$$\begin{aligned}\mathcal{O} &= \mathbb{Z} + \mathbb{Z} \frac{-\Delta + \sqrt{-\Delta}}{2}, \\ \mathcal{O}^\# &= \mathbb{Z} \frac{i}{\sqrt{-\Delta}} + \mathbb{Z} \frac{1 + \sqrt{-\Delta}}{2}.\end{aligned}$$

From now on, we will always work with Hermitian modular forms of degree 2, i.e. we will always have $n = 2$, except if otherwise stated.

Chapter 3

Theory

Lemma 3.1. *Let $f: M_2(\mathbb{C}) \rightarrow \mathbb{C}$ be a Hermitian Modular form of weight k . Then, $f(S\tau): \mathbb{H}_1 \subseteq \mathbb{C} \rightarrow \mathbb{C}$ is an elliptic modular form of weight $2k$ for some matrix $S \in M_2(\mathbb{Z})$ with $\Gamma(S) \subseteq \text{SL}_2(\mathbb{Z})$.*

Lemma 3.2. *Prop 7.3. von Poor für herm Modulformen. $\Gamma(\mathcal{L}) \supseteq \Gamma_0(l)$ for $l \in \mathbb{Z}^+, ls^{-1} \in \mathcal{P}_n(\mathcal{O})$.*

Now we will formulate the main algorithm of our work.

Algorithm 3.3. 1. Select a set of matrices $\mathcal{S} \subseteq M_2^T(\mathbb{Z})$ with $0 < S \in \mathcal{S}$. Make \mathcal{S} big enough. Now, for some $S \in \mathcal{S}$:

2. Fix $B \in \mathbb{N}$ as a limit. Or select a precision

$$\mathcal{F} := \left\{ \begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} \left| 0 \leq ac < B, b \in \mathcal{O}^\# \right. \right\} \subseteq \Lambda,$$

where

$$\Lambda := \left\{ 0 \leq \begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} \in M_2(\mathcal{O}^\#) \left| a, c \in \mathbb{Z} \right. \right\}.$$

3.

$$\mathcal{M}_{k,\mathcal{S},\mathcal{F}}^H = \{f[S] \mid f \in \mathbb{Q}^{\mathcal{F}}\},$$

$$\mathcal{M}_{k,S} = \mathcal{FE}_{\mathcal{F}(S)}(M_k(\Gamma(S)))$$

4. If

$$\dim \mathcal{M}_{k,\mathcal{S},\mathcal{F}}^H \cap \bigoplus_{S \in \mathcal{S}} \mathcal{M}_{k,S} = \dim M_k^H,$$

then we are ready and we can reconstruct the Fourier expansion in the following way:

...

Chapter 4

Implementation

In this chapter, we are describing the implementation.

Chapter 5

Conclusion

Blub

Chapter 6

References

- [PY07] C. Poor and D.S. Yuen. Computations of spaces of siegel modular cusp forms.
Journal of the Mathematical Society of Japan, 59(1):185–222, 2007.
- [Rau12] M. Raum. Computing Jacobi Forms and Linear Equivalences of Special Divisors.
ArXiv e-prints, December 2012.