

CALCULATIONS OF HERMITIAN MODULAR FORMS

BERECHNUNGEN HERMITESCHER MODULFORMEN

DIPLOMA THESIS
in Mathematics

by
Albert Zeyer

submitted to the
Faculty of Mathematics, Computer Science and Natural Science of
RWTH Aachen University

October 2012
revised version from November 29, 2012

Supervisor: Prof. Dr. Aloys Krieg
Second examiner: Dr. Martin Raum

written at the
Lehrstuhl A für Mathematik
Prof. Dr. A. Krieg

Contents

1	Introduction	3
2	Background results	4
2.1	Preliminaries	4
3	Theory	5
4	Implementation	6
5	Conclusion	7
6	References	8

Chapter 1

Introduction

In [PY07], spaces of Siegel modular cusp forms are calculated.

We are doing the same for hermitian modular forms.

Chapter 2

Background results

2.1 Preliminaries

Let $M_n(\mathcal{K})$ be the set of all $n \times n$ matrices over some field \mathcal{K} . Likewise, $M_n^T(\mathcal{K})$ are the symmetric $n \times n$ matrices. A matrix $Y \in M_n(\mathbb{R})$ is greater 0 iff $\forall x \in \mathbb{R}^n - \{0\} : Y[x] := x^T Y x > 0$. Let $\mathbb{H}_n := \{Z = X + iY \in M_n^T(\mathbb{C}) \mid Y > 0\}$. Thus, \mathbb{H}_1 is the Poincaré upper half plane.

The general linear group is defined by $GL_n(\mathcal{K}) = \{X \in M_n(\mathcal{K}) \mid \det(X) \neq 0\}$ and the special linear group by $SL_n(\mathcal{K}) = \{X \in M_n(\mathcal{K}) \mid \det(X) = 1\}$. The orthogonal group is defined by $O_n(\mathcal{K}) = \{X \in GL_n(\mathcal{K}) \mid X^T X = 1_n\}$. The symplectic group is defined by $Sp_n(\mathcal{K}) = \{X \in GL_{2n}(\mathcal{K}) \mid X^T J_n X = J_n\}$ where $J_n := \begin{pmatrix} 0 & 1_n \\ -1_n & 0 \end{pmatrix} \in SL_{2n}(\mathcal{K})$.

A siegel modular cusp form is a holomorphic function

$$f: \mathbb{H}_n \rightarrow \mathbb{C}$$

with

- (1) $f|_k y = f \forall y \in \Gamma \subseteq Sp_n(\mathbb{Z})$
- (2) for $n = 1$: $f(Z) = O(1)$ for $Z \rightarrow i\infty$

where

$$\left(f|_k \begin{pmatrix} A & B \\ C & D \end{pmatrix}\right)(Z) = f((AZ + B)(CZ + D)^{-1}) \cdot \det(CZ + D)^{-k}$$

with $Z = S\tau$.

Chapter 3

Theory

Lemma 3.1. *Let $f: \mathbb{M}_2(\mathbb{C}) \rightarrow \mathbb{C}$ be a hermitian modular form of weight k . Then, $f(S\tau): \mathbb{H}_1 \subseteq \mathbb{C} \rightarrow \mathbb{C}$ is an elliptic modular form of weight $2k$ for some matrix $S \in \mathbb{M}_2(\mathbb{Z})$ with $\Gamma(S) \subseteq \mathrm{SL}_2(\mathbb{Z})$.*

Algorithm 3.2. 1. Select a set of matrices $\mathcal{S} \subseteq \mathbb{M}_2^T(\mathbb{Z})$ with $0 < S \in \mathcal{S}$. Make \mathcal{S} big enough. Now, for some $S \in \mathcal{S}$:

2. Fix $B \in \mathbb{N}$ as a limit. Or select a precision

$$\mathcal{F} = \left\{ \begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} \middle| 0 \leq ac < B \right\} \subseteq \Lambda,$$

where

$$\Lambda := \left\{ 0 \leq \begin{pmatrix} a & b \\ \bar{b} & c \end{pmatrix} \in \mathbb{M}_2(o^\#) \middle| a, c \in \mathbb{Z} \right\}.$$

3.

$$\mathcal{M}_{k,S,\mathcal{F}}^H = \{fS \mid f \in \mathbb{Q}^{\mathcal{F}}\},$$

$$\mathcal{M}_{k,S} = \text{FourierExpansion}_{\mathcal{F}(S)}(\mathbb{M}_k(\Gamma(S)))$$

4. If

$$\dim \mathcal{M}_{k,S,\mathcal{F}}^H \cap \bigoplus_{S \in \mathcal{S}} \mathcal{M}_{k,S} = \dim M_k^H,$$

then we are ready and we can reconstruct the Fourier expansion in the following way:

...

Chapter 4

Implementation

In this chapter, we are describing the implementation.

Chapter 5

Conclusion

Blub

Chapter 6

References

- [PY07] C. Poor and D.S. Yuen. Computations of spaces of siegel modular cusp forms.
Journal of the Mathematical Society of Japan, 59(1):185–222, 2007.