

Streaming Data Management and Time Series Analysis

Alberto Filosa

28/9/2020

Indice

1 Statistical Prediction	1
1.1 Optimal Linear Predictor	1
2 Time Series Concepts	1
2.1 Moving Average Process	2
2.2 Autoregressive Process	2
2.3 Integrated Process	2
2.4 ARIMA Models	2
3 Unobserved Components Model	3
3.1 Trend	3

2. *Orthogonality of the prediction error:* $\mathbb{E}\{(Y - \hat{Y})g(x)\} = \mathbb{E}\{(Y - \mathbb{E}[Y|X])g(x)\} = 0$;
3. *Functions of conditioning variables:* $\mathbb{E}[Yg(x)|X] = \mathbb{E}[Y|X]g(x)$;
4. *Independence with the conditioning variables:* $\mathbb{E}[Y|X] = \mathbb{E}[Y]$ if $X \perp\!\!\!\perp Y$
5. *Law of iterated expectations:* $\mathbb{E}[Y] = \mathbb{E}\{\mathbb{E}[Y|X]\}$;
6. *Law of total variance:* $\text{Var}[Y] = \mathbb{E}[\text{Var}(Y|X)] + \text{Var}[\mathbb{E}[Y|X]]$.

1.1 Optimal Linear Predictor

Sometimes it can be easier to limit the search to a smaller class of functions, like linear combination. The main advantage is that the covariance structure of the random variables is all needed to compute the prediction:

1 Statistical Prediction

A *Statistical Prediction* is a guess about the value of a random variable Y based on the outcome of other random variables X_1, \dots, X_m . A *Predictor* is a function of the random variables: $\hat{Y} = p(X_1, \dots, X_m)$. An optimal prediction is the *Loss Function* which maps the prediction error to its cost; if the prediction error is zero, also the loss is 0 (exact guess = no losses). A loss function can be symmetric about 0 or asymmetric: $l(-x) = l(x)$, but the most used is generally asymmetric, like the *Quadratic loss function*: $l_2(x) = x^2$.

The predictor $\hat{Y} = \hat{p}(X_1, \dots, X_m)$ is optimal for Y with respect to the loss l if it minimizes the expected loss among the class of measurable functions:

$$\mathbb{E} l(Y - \hat{Y}) = \min_{p \in M} \mathbb{E} l[Y - p(X_1, \dots, X_m)]$$

The optimal predictor under quadratic loss is the conditional expectation $\hat{Y} = \mathbb{E}[Y|X_1, \dots, X_m]$.

The properties of the conditional expectation are:

1. *Linearity:* $\mathbb{E}[aY + bZ + c|X] = a\mathbb{E}[Y|X] + b\mathbb{E}[Z|X] + c$, with a, b, c , constants;

$$\mathbb{P}[Y|X_1, \dots, X_m] = \mu_Y + \Sigma_{YX} \Sigma_{XX}^{-1} (X - \mu_X)$$

The Properties of the optimal linear predictor are:

1. *Unbiasedness:* $\mathbb{E}[Y - \mathbb{P}[Y|X]] = 0$;
2. *Mean Square Error (MSE)* of the prediction: $MSE_{lin} = \Sigma_{YY} - \Sigma_{YX} \Sigma_{XX}^{-1} \Sigma_{XY}$
3. *Orthogonality of the prediction error:* $\mathbb{E}[(Y - \mathbb{P}[Y|X])X^t] = 0$;
4. *Linearity:* $\mathbb{P}[aY + bZ + c|X] = a\mathbb{P}[Y|X] + b\mathbb{P}[Z|X] + c$
5. *Law of iterated projections:* if $\mathbb{E}(X - \mu_X)(Z - \mu_Z)^t = 0$, then $\mathbb{P}[Y|X, Z] = \mathbb{P}[Y|X] + \mathbb{P}[Y - \mathbb{P}[Y|X]|Z]$

2 Time Series Concepts

A *Time Series* is a sequence of observations ordered to a time index t taking values in an index set S . If S contains finite numbers we speak about *Discrete* time series y_t , otherwise a *Continuous* time series $y(t)$.

The most important form of time homogeneity is *Stationarity*, defined as time-invariance of the whole probability of the data generating process (*strict stationarity*) or just of its two moments (*weak stationarity*).

The process $\{Y_t\}$ is *Strictly* stationary if $\forall k \in \mathbb{N}, h \in \mathbb{Z}$ and $(t_1, \dots, t_k) \in \mathbb{Z}^k$,

$$(Y_{t_1}, \dots, Y_{t_k}) \stackrel{d}{=} (Y_{t_1+h}, \dots, Y_{t_k+h})$$

The process $\{Y_t\}$ is *Weakly* stationary if, $\forall h, t \in \mathbb{Z}$, with $\gamma(0) < \infty$

$$\begin{aligned} \mathbb{E}(Y_t) &= \mu \\ \text{Cov}(Y_t, Y_{t-h}) &= \gamma(h) \end{aligned}$$

If a time series is strictly stationary, then it is also weakly stationary if and only if $\text{Var}(Y_t) < \infty$. If a time series a Gaussian process, then strict and weak stationarity are equivalent.

The most elementary stationarity process is the white noise. A stochastic process is *White Noise* if has $\mu = 0, \sigma^2 > 0$ and covariance function

$$\gamma(h) = \begin{cases} \sigma^2 & \text{for } h = 0 \\ 0 & \text{for } h \neq 0 \end{cases}$$

The *Autocovariance Function* is a function characterized by a weakly stationary process, while the *Autocorrelation Function* (**ACF**) is the scale independent version of the autocovariance function:

If $\{Y_t\}$ is a stationary process with autocovariance $\gamma(\cdot)$, the its ACF is $p(h) = \text{Cor}(Y_t, Y_{t-h}) = \gamma(h)/\gamma(0)$.

An alternative summary of the linear dependence of a stationary process can be obtained from the *Partial Autocorrelation Function*, that measures the correlation between Y_t and Y_{t-h} after their linear dependence on the intervening random variables has been removed

2.1 Moving Average Process

A **Moving Average Process (MA)** is a process that estimates the trend-cycle at time t obtained by averaging values of the time series within k periods of t . The average eliminates some of the randomness in the data, leaving a smooth trend-cycle component.

$$\begin{aligned} Y_t &= \varepsilon_t + \theta_1 \varepsilon_{t-1} + \dots + \theta_q \varepsilon_{t-q} \\ Y_{t-1} &= \varepsilon_{t-1} + \theta_1 \varepsilon_{t-2} + \dots + \theta_q \varepsilon_{t-q-1} \end{aligned}$$

A $MA(q)$ process has $PACF = 0$ for $h > p$ and its characteristic equation $1 + \theta_1 X + \dots + \theta_q X^q = 0$ that has only external solution of the unitary circle is a process such that $Y_t = k + \psi_1 Y_{t-1} + \dots + \psi_q Y_{t-q} + \varepsilon_t$

2.2 Autoregressive Process

An Autoregressive Process is a process that forecasts the Y variable using a linear combination of past values of the variable. The term autoregression indicates that it is a regression of the variable against itself.

$$Y_t = k + \phi_1 Y_{t-1} + \dots + \phi_p Y_{t-p} + \varepsilon_t$$

An $AR(p)$ process is stationary for the characteristic equation $1 - \phi_1 X - \dots - \phi_p X^p = 0$ and all solutions are external of the unitary circle

If Y_t is stationary, $\mathbb{E}(Y_t) = \frac{k}{1 - \phi_1 - \dots - \phi_p}$.

The $AR(1)$ process is $Y_t = k + \phi Y_{t-1} + \varepsilon_t$, so $1 - \phi x = 0 \implies x = 1/\phi$. $AR(1)$ is stationary if $|\phi| < 1$. If in $AR(1)$ $\phi = 1$ we obtain a non stationary process (integrate) called Random Walk:

$$Y_t = k + Y_{t-1} + \varepsilon$$

2.3 Integrated Process

An integrated process $\{Y_t\} \sim I(d)$ is a non stationary process, but its first difference is stationary:

$$\Delta Z_t = Z_t - Z_{t-1} \sim I(0)$$

A Z_t process is integrated of order d if it is not stationary, $\Delta^{d-1} Z_{t-1}$ non stationary, while $\Delta^d Z_t$ is stationary.

The process $\{Y_t\}$ is $ARMA(p, q)$ if it is stationary and satisfies:

$$Y_t = \phi_1 Y_{t-1} + \dots + \phi_p Y_{t-p} + \theta_1 Z_{t-1} + \dots + \theta_q Z_{t-q}$$

2.4 ARIMA Models

If we combine differencing with autoregression and a moving average model, we obtain a non-seasonal **AutoRegressive Integrated Moving Average (ARIMA)**.

$$Y'_t = c + \phi_1 Y'_{t-1} + \dots + \phi_p Y'_{t-p} + \theta_1 \varepsilon_{t-1} + \dots + \theta_q \varepsilon_{t-q} + \varepsilon_t$$


where Y'_t is the differenced series. This is an $ARIMA(p, d, q)$, where p is the order of the **AutoRegressive** part, d is the degree of first differencing

involved (**I**ntegrated) and q is the order of the **M**oving **A**verage part.

The backward shift operator $B(y_t) = y_{t-1}$ is a useful notational device when working with time series lags, so it has the effect of shifting the data back one period. The backward shift operator is convenient for describing the process of differencing. A first difference can be written as:

$$y'_t = y_t - y_{t-1} = y_t - B(y_t) = (1 - B)y_t$$

Backshift notation is particularly useful when combining differences, as the operator can be treated using ordinary algebraic rules. In particular, terms involving B can be multiplied together.



3 Unobserved Comoponents Model

A natural way humans tend to think about time series is as sum of non directly observable componen-
ts such as trends, seasonality and cycle. **U**nobserved **C**omponents **M**odels (**UCM**) select the best features of stochastic framework as ARIMA models, but also they tend to perform very well in forecasting .

The observable time series of UCM is the sum of trend (μ_t), cycle (ψ_t), seasonality (γ_t) and white noise (ε_t):

$$Y_t = \mu_t + \psi_t + \gamma_t + \varepsilon_t$$

Some of these components could be skipped and some other could be added. Also, they can be seen as stochastic version of the deterministic functions of time. In the

next sections we see how to build stochastically evolving components.

3.1 Trend

The *Trend* usually adopted in UCM is the local linear trend. Let us take a linear function defined as follows:

$$\mu_t = \mu_0 + \beta_0 t$$

where μ_0 is the intercept and β_0 the slope and write it in incremental form:

$$\mu_t = \mu_{t-1} + \beta_0$$

By adding the white noise η_n , we obtain a random walk with drift. In this case we can interpreted μ_t as a linear trend with a random walk intercept, but the slope remains unchanged.

It is possible obtain a time-varying slope in trends, easily achieved letting the slope evolves as a random walk:

$$\begin{aligned}\mu_t &= \mu_{t-1} + \beta_0 + \eta_n \\ \beta_t &= \beta_{t-1} + \xi_t\end{aligned}$$

These equations define the local linear trend interpreted as a linear trend where both intercept and slope evolve in time as random walks

The local linear trend has different special case of interest obtained by fixing the value of the variances:

- *Deterministic Linear Trend* if $\sigma_\eta^2 = \sigma_\xi^2 = 0$;
- *Random Walk with Drift* β_0 if $\sigma_\xi^2 = 0$ (slope constant);
- *Random Walk* if $\sigma_\xi^2 = \beta_0 = 0$ (slope = 0)
- *Integrated Random Walk* if $\sigma_\eta^2 = 0$, with a very smooth trend

Obs.: the local linear trend can be also seen as an ARI-MA process: if $\sigma_\xi^2 > 0$, $\mu_t \sim I(2)$ process as trend is non-stationary, while its second difference is stationary.

```
set.seed(20201006)
```

```
n <- 200 #-- Observations
```

```
beta <- numeric(n) #-- Slope
```

```
mu <- numeric(n) #-- Trend
```

```
sd_eta <- 1 #-- Standard Deviation White Noise
```

```
sd_xi <- 0.1 #-- Standard Deviation White Noise
```

```


for (t in 2:n){
  beta[t] <- beta[t-1] + rnorm(1,sd = sd_xi)          #-- Slope
  mu[t] <- mu[t-1] + beta[t-1] + rnorm(1, sd = sd_eta) #-- Intercept
}

```

```

par(mfrow = c(1,2))
plot(ts(beta))
plot(ts(mu))

```



```

par(mfrow = c(1,1))

```