#### BSPS 2017

# MACRO PATTERNS IN THE EVOLUTION OF HUMAN AGING

#### Tim Riffe & José Manuel Aburto





MAX PLANCK INSTITUTE FOR DEMOGRAPHIC RESEARCH

8<sup>th</sup> September 2017

# Background


▶ **Life expectancy** at birth  $(e_0)$  is the most used indicator.

# Background

- ▶ **Life expectancy** at birth  $(e_0)$  is the most used indicator.
- ▶ Demographers also use modal or median age at death.

# Background

- ▶ **Life expectancy** at birth  $(e_0)$  is the most used indicator.
- ▶ Demographers also use modal or median age at death.
- ► They **conceal variation of lifespans** and other aspects of the age at death distribution.



## Key formulas

Consider the conditional death distribution

$$f(y \mid a) = \frac{1}{\ell(a)} \mu(a+y) \ell(a+y)$$

# Key formulas

Consider the conditional death distribution

$$f(y \mid a) = \frac{1}{\ell(a)} \mu(a+y) \ell(a+y)$$

 $\mu(a) \longrightarrow$  force of mortality at age a  $\ell(a) \longrightarrow$  survivorship to age a

# Key formulas

Consider the conditional death distribution

$$f(y \mid a) = \frac{1}{\ell(a)} \mu(a+y) \ell(a+y)$$

 $\mu(a) \longrightarrow$  force of mortality at age a  $\ell(a) \longrightarrow$  survivorship to age a

Key point:  $f(y \mid a) \longrightarrow$  probability of surviving to and dying at age a + y given survival to age a.

Remaining life expectancy conditional on survival to age a is

$$e(a) = \frac{1}{\ell(a)} \int_0^\infty \ell(a+y) dy$$

The conditional deaths distribution can be described by its moments about e(a)

$$\eta_n(y \mid a) = \int_{y=0}^{\infty} (y - e(a))^n f(y \mid a) dy$$

$$\eta_n(y \mid a) = \int_{y=0}^{\infty} (y - e(a))^n f(y \mid a) dy$$

Variance 
$$\longrightarrow \sigma^2(y \mid a) = \eta_2(y \mid a)$$

$$\eta_n(y \mid a) = \int_{y=0}^{\infty} (y - e(a))^n f(y \mid a) dy$$

Variance 
$$\longrightarrow \sigma^2(y \mid a) = \eta_2(y \mid a)$$

Skewness 
$$\longrightarrow Skew(y \mid a) = \frac{\eta_3(y|a)}{\sigma^2(y|a)}$$


$$\eta_n(y \mid a) = \int_{y=0}^{\infty} (y - e(a))^n f(y \mid a) dy$$

Variance 
$$\longrightarrow \sigma^2(y \mid a) = \eta_2(y \mid a)$$

Skewness 
$$\longrightarrow$$
  $Skew(y \mid a) = \frac{\eta_3(y \mid a)}{\sigma^2(y \mid a)}$ 

Kurtosis 
$$\longrightarrow Kurt(y \mid a) = \frac{\eta_4(y|a)}{\sigma^3(y|a)-3}$$

:



# What if we take different ages?

What if we take different ages?


What if we take different years?


What if we take different ages?


What if we take different years?


We explore with our framework
















Introduction Formulae Illustration











## Go beyond the mean!

More information:

jmaburto@health.sdu.dk riffe@demogr.mpg.de

@jm\_aburto @timriffe1

@jmaburto @timriffe





FORSCHUNG

