

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

I declare this is my own work.

INTERNATIONAL A-LEVEL MATHEMATICS

(9660/MA03) Unit P2 Pure Mathematics

Friday 17 January 2020 07:00 GMT Time allowed: 2 hours 30 minutes

Materials

- For this paper you must have the Oxford International AQA booklet of formulae and statistical tables (enclosed).
- You may use a graphics calculator.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 120.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working: otherwise marks may be lost.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
TOTAL	



J A N 2 0 M A 0 3 0 1

Answer **all** questions in the spaces provided.

- 1 (a) (i)** Express $5 \sin \theta - 12 \cos \theta$ in the form $R \sin(\theta - \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$, giving the value of α to the nearest 0.1°

[3 marks]

Answer _____

- 1 (a) (ii)** Hence solve the equation

$$5 \sin \theta - 12 \cos \theta = -1$$

in the interval $-180^\circ < \theta < 180^\circ$, giving all solutions to the nearest 0.1°

[3 marks]

Answer _____



$$2 \cot^2 x = 10 - 5 \operatorname{cosec} x$$

[5 marks]

[illegible]

Answer



2 The function f is defined by

$$f(x) = \sin^{-1}(2x - 1)$$

2 (a) State the largest possible domain of f .


[1 mark]

Answer _____

2 (b) On the axes below, sketch the graph of $y = f(x)$.

Show the coordinates of the end-points on the graph.

[2 marks]



- 2 (c)** Describe a sequence of **two** geometrical transformations that maps the graph of $y = \sin^{-1} x$ onto the graph of $y = \sin^{-1} (2x - 1)$

[4 marks]

- 2 (d)** The root of the equation $\sin^{-1} (2x - 1) + x - 1 = 0$ is α .

Show that α lies between 0.6 and 0.7

[2 marks]

Question 2 continues on the next page

Turn over ►



2 (e) Use the iterative formula

$$x_{n+1} = \frac{1 + \sin(1 - x_n)}{2} \quad \text{with } x_1 = 0.6$$

to find the values of x_2 and x_3 , giving your answers to three decimal places.

[2 marks]

Answer _____

11



3 (a) Find $\int x \ln x \, dx$

[3 marks]

Answer _____

3 (b) Find $\int \ln x \, dx$

[3 marks]

Answer _____

Turn over ►



4 (a) The polynomial $f(x)$ is defined by

$$f(x) = 8x^3 + bx^2 + cx + 6$$

where b and c are constants.

When $f(x)$ is divided by $(2x - 1)$ the remainder is 5.25

When $f(x)$ is divided by $(2x - 3)$ the remainder is -3.75

Find the value of b and the value of c .

[4 marks]

[illegible]

$$b = \quad \quad \quad c =$$



where d , e and f are integers.

[illegible]

Turn over ►



5 (a) Express $\frac{12}{9-u^2}$ in the form $\frac{A}{3-u} + \frac{B}{3+u}$

[2 marks]

Answer _____



8

- 6 (a) (i)** Find the binomial expansion of $(1 + 2x)^{0.5}$ in ascending powers of x up to and including the term in x^2

[2 marks]

Answer _____

- 6 (a) (ii)** Find the binomial expansion of $(1 - 4x)^{-0.5}$ in ascending powers of x up to and including the term in x^2

[2 marks]

Answer _____

- 6 (b) (i)** Hence find the binomial expansion of $\sqrt{\frac{1+2x}{1-4x}}$ in ascending powers of x up to and including the term in x^2

[2 marks]

Answer _____



6 (b) (ii) State the values of x for which the expansion of $\sqrt{\frac{1+2x}{1-4x}}$ is valid.

[1 mark]

Answer _____

6 (c) Use your expansion of $\sqrt{\frac{1+2x}{1-4x}}$ to find an estimate for $\sqrt{2}$, giving your answer to three decimal places.

[3 marks]

Answer _____



7 A curve has equation $y = e^{3x} - 24x$

7 (a) Find an equation of the tangent to the curve at (0, 1)

[3 marks]

Answer _____

7 (b) Find the coordinates of the stationary point of the curve, giving your answer in an exact form.

[3 marks]

Answer _____



[3 marks]

[illegible]

Answer _____

9

Turn over ►



The rate of change of the evaporated mass, x grams, is directly proportional to the mass of the liquid which remains at time t minutes after the start of the experiment.

8 (a) Explain briefly why this information can be represented by the differential equation

$$\frac{dx}{dt} = k(80 - x)$$

Solve the differential equation in part (a) to find the value of k .

[illegible]

Answer



[2 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Answer _____

[3 marks]

[illegible]

Answer _____



[6 marks]

[illegible]

Answer

6



- 10 (a)** Use the mid-ordinate rule with six strips, to find an estimate for $\int_{1.5}^3 x^{-x} dx$, giving your answer to three decimal places.

[4 marks]

Answer _____

- 10 (b)** By taking logarithms of both sides of $y = x^{-x}$ and then using implicit differentiation, find $\frac{dy}{dx}$, giving your answer in terms of x only.

[4 marks]

Answer _____

Turn over ►

11 The point A has coordinates $(10, 2, -3)$.

The point B has coordinates $(2, -2, 5)$.

11 (a) Find the vector equation of the line AB and hence show that it can be written as

$$\mathbf{r} = \begin{bmatrix} 4 \\ -1 \\ 3 \end{bmatrix} + \lambda \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix}$$

[4 marks]

Answer _____

11 (b) The point D has coordinates $(-2, 1, 7)$.

The point C lies on the line AB .

The line CD is perpendicular to the line AB .

11 (b) (i) Find the coordinates of C .

[5 marks]



Answer _____

11 (b) (ii) Show that the distance $CD = \sqrt{q}$ where q is a constant.

[2 marks]

11 (c) The point $P(4 + 2p, -1 + p, 3 - 2p)$ lies on the line AB such that triangle DCP is isosceles.

Find the possible exact values of p , giving your answers in the form $s + t\sqrt{q}$, where s and t are constants.

[5 marks]

Answer _____



12

12 (a)

[5 marks]

[illegible]

12 (b)

[4 marks]

[illegible]

Answer

9



A curve C is defined by the parametric equations $x = at^2$, $y = 2at$, where a is a constant.

Given that p and q vary so that $p^2 + q^2 = 1$, find the Cartesian equation of the curve on which R lies, giving your answer in the form $y^2 = f(x)$

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Answer

7

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



[illegible]

[illegible]

[illegible]

There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.oxfordaqaexams.org.uk

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2020 Oxford International AQA Examinations and its licensors. All rights reserved.

