

Please write clearly ir	Please write clearly in block capitals.			
Centre number	Candidate number			
Surname				
Forename(s)				
Candidate signature	I declare this is my own work.	ノ		

INTERNATIONAL A-LEVEL **MATHEMATICS**

(9660/MA05) Unit M2 Mechanics

Thursday 19 January 2023 07:00 GMT Time allowed: 1 hour 30 minutes

Materials

- For this paper you must have the Oxford International AQA Booklet of Formulae and Statistical Tables (enclosed).
- You may use a graphical calculator.

Instructions


- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- The **final** answer to questions requiring the use of calculators should be given to two significant figures, unless stated otherwise.
- ullet Unless stated otherwise, the acceleration due to gravity, g, should be taken as 9.8 m s $^{-2}$

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.





There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



Answer **all** questions in the spaces provided.

1 A force **F** newtons acts on a body of mass 0.5 kg.

The position vector \mathbf{r} metres of the body at time t seconds is given by

$$\mathbf{r} = \begin{bmatrix} 3t^2 \\ \cos(4t) \\ e^{t^2} \end{bmatrix}$$

Find \mathbf{F} in terms of t	[5 marks]
	·

Answer

5




2		A person stands at the end of a uniform horizontal diving board AC, as shown below.
		FI.
		8
		A B
		Ц
		<
		1.25 m 2.75 m
		The diving board is attached to a vertical wall at point <i>A</i> and rests on a support at point <i>B</i>
		The mass of the person is 60 kg.
		The mass of the uniform diving board is 25 kg.
		The distance <i>AB</i> is 1.25 metres.
		The distance <i>BC</i> is 2.75 metres.
		The diving board is in equilibrium.
2	(a)	Find the magnitude of the normal reaction force acting on the diving board at <i>B</i> [3 marks]
		[o marko]
		Answer



2	(b)	Find the force acting on the diving board at A, giving its magnitude and direction. [3 marks]	Do not write outside the box
		Magnitude	6
		Turn over for the next question	



A motorless cart starts from rest at the point *X* and moves along a curved track through the point *Y* before coming to rest at the point *Z*, as shown below.



The mass of the cart is 16 kg.

The point X is 4.0 metres vertically above the level of Y

The point Z is 2.5 metres vertically above the level of Y

The distance travelled by the cart along the track between X and Y is 30 metres.

The distance travelled by the cart along the track between Y and Z is 18 metres.

Assume that throughout the motion a constant resistance force of magnitude $\,F\,$ newtons acts on the cart.


Find the value of F	[4 marks]
Answer	



3	(b)	Find the speed of the cart at Y		
			[3 marks]	
		Answer		
		Turn over for the next question		
				i



The diagram below shows the dimensions of a rectangular uniform lamina *OABC* in which a square hole has been cut.



4	(a)	[2 marks]
4	(b)	Find the distance of the centre of mass of the lamina from <i>OC</i> giving your answer to three significant figures.
		[3 marks]



Anguar	
Answer	
The lamina hangs in equilibrium, freely suspended from <i>B</i>	
Find the angle between OB and the vertical, giving your answer to	the nearest 0.1° [4 marks]
	[::::::::::
-	



5		A particle of mass 6 kg is released from rest at the point X on a long slope. The particle begins to slide down the slope.
		The slope is inclined at 25° to the horizontal as shown below.
		X
		25°
		The slope is smooth between X and the point Y where the distance $XY = 10$ metres.
		The slope is rough from $$ Y downwards. The coefficient of friction between the particle and this part of the slope is 0.5
5	(a) (i)	Find the magnitude of the particle's acceleration between $ X $ and $ Y $ [1 mark]
		Answer
5	(a) (ii)	Show that the speed of the particle at Y is 9.1 m s ⁻¹ correct to two significant figures. [2 marks]
5	(a) (iii)	State, with a reason, how the speed given in part (a)(ii) would change if the slope was inclined at an acute angle greater than 25° to the horizontal. [2 marks]



Do ou	no tsid	t w de ox	rit the	e

	Find the distance XZ [6 marks]
	Answer
5 (c)	The slope is now adjusted so that it is inclined at $lpha$ degrees to the horizontal.
5 (c)	
5 (c)	The particle is again released from rest at X and begins to slide down the slope.
5 (c)	
5 (c)	The particle is again released from rest at X and begins to slide down the slope.
5 (c)	The particle is again released from rest at X and begins to slide down the slope. This time the particle does not come to rest on the slope. Find the smallest possible value of α , giving your answer to one decimal place. [2 marks]
5 (c)	The particle is again released from rest at X and begins to slide down the slope. This time the particle does not come to rest on the slope. Find the smallest possible value of α , giving your answer to one decimal place. [2 marks]
5 (c)	The particle is again released from rest at X and begins to slide down the slope. This time the particle does not come to rest on the slope. Find the smallest possible value of α , giving your answer to one decimal place. [2 marks]
5 (c)	The particle is again released from rest at X and begins to slide down the slope. This time the particle does not come to rest on the slope. Find the smallest possible value of α , giving your answer to one decimal place. [2 marks]
5 (c)	The particle is again released from rest at X and begins to slide down the slope. This time the particle does not come to rest on the slope. Find the smallest possible value of α , giving your answer to one decimal place. [2 marks]

13



Do not write
outside the
box

6		The acceleration \mathbf{a} m s ⁻² of a particle at time t seconds is	given by
		$\mathbf{a} = e^{-2t}\mathbf{i} + \frac{1}{\left(1+t\right)^2}\mathbf{j} \qquad \text{for } t \ge 0$	0
		where the unit vectors $ {f i} $ and $ {f j} $ are perpendicular.	
6	(a)	The particle is initially at rest.	
		Find the velocity of the particle in terms of <i>t</i>	[4 marks]
		Answer	
		, ,,,,,,,,	



6 (b) The particle's initial position is 3 i + 2 j relative to a fixed origin O	Do r outs
Find the distance of the particle from O when $t = 5$	
	[5 marks]
Answer	



7 A particle is projected from a point $\,$ O $\,$ on horizontal ground with a speed of 15 m $\,$ s $^{-1}$ at an angle α above the horizontal, where $0^{\circ} < \alpha < 90^{\circ}$ The particle passes through the point *P* which has a horizontal displacement of 12 metres from O and a vertical displacement of 5 metres from O Find the maximum possible height that the particle can reach, giving your answer to three significant figures. [9 marks]




	Do not write outside the box
	Jox
Answer_	9



8 A satellite of mass m kg moves in a circular orbit of radius r metres with constant speed v m s⁻¹ around a planet of mass M kg

The initial position of the satellite is shown in the diagram.



The unit vectors $\, \mathbf{i} \,$ and $\, \mathbf{j} \,$ are directed as shown in the diagram.

The only force which acts on the satellite is always directed towards the centre of the planet. This force has magnitude F newtons such that

$$F = \frac{GMm}{r^2}$$

where $G \, \mathrm{m^3 \, kg^{-1} \, s^{-2}}$ is a constant.

The time for the satellite to make one revolution around its orbit is T seconds.

8 (a) Show that

$$T^2 = \frac{4\pi^2 r^3}{GM}$$

[4 marks]

8	(b)	It is given that $r = 4.2 \times 10^7$ and $M = 6.0 \times 10^{24}$	
8	(b) (i)	Find the angular speed of the satellite. Take the value of G to be 6.7×10^{-11}	
		Answer	
8	(b) (ii)	The initial position vector of the satellite relative to the planet is 4.2×10^7 j me	etres.
		Find the velocity of the satellite t seconds after leaving its initial position. Give your answer in the form $a\mathbf{i} + b\mathbf{j}$	
		Give your answer in the form $a\mathbf{i} + b\mathbf{j}$	[3 marks]
		Answer	

10




narks]
narks]



			Do not write outside the box
		Answer	
9	(c) (i)	Show that the work done by the force of magnitude T newtons in pulling the particle 25 metres up the inclined plane is 2800 J, correct to three significant figures. [1 mark]	
9	(c) (ii)	Find the rate at which the force T newtons does work on the particle. [2 marks]	
		Answer	12

END OF QUESTIONS







Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Do not write There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.oxfordaqaexams.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the

Copyright © 2023 Oxford International AQA Examinations and its licensors. All rights reserved.





outside the box