

Please write clearly in block capitals.	
Centre number	Candidate number
Surname	
Forename(s)	
Candidate signature	

INTERNATIONAL AS **MATHEMATICS**

(9660/MA01) - Pure Mathematics Unit 1

Thursday 24 May 2018

07.00 GMT

Time allowed: 1 hour 30 minutes

Materials

- For this paper you must have the Oxford International AQA booklet of formulae and statistical tables.
- · You may use a graphics calculator.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside
 the box around each page or on blank pages. If you require extra space, use a
 supplementary answer book.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.

For Examiner's Use				
Question	Mark			
1				
2				
3				
4				
5				
6				
7				
8				
9				
10				
TOTAL				



Answer all questions in the spaces provided.

- 1 (a) The expression $x^2 + 6x + 14$ can be written in the form $(x + a)^2 + b$
- **1** (a) (i) Find the value of a.

Circle your answer.

[1 mark]

2

3

5

6

1 (a) (ii) Find the value of *b*.

Circle your answer.

[1 mark]

2

3

5

6

1 (b) The graph of $y = x^2$ is translated onto the graph of $y = x^2 + 6x + 14$ by a vector.

Find this vector.

[2 marks]

Answer

4

(a)	Show that $5\sqrt{8} \times 3\sqrt{12}$ can be written in the form $a\sqrt{b}$ where a and b are integers.	
	Simplify your answer as far as possible. [2 r	nark
(b)	Show that $\frac{3\sqrt{7}-4\sqrt{6}}{2\sqrt{7}+\sqrt{6}}$ can be written in the form $\frac{p-\sqrt{q}}{r}$ where p,q and r are integrated as $\frac{1}{2\sqrt{7}+\sqrt{6}}$.	gers
	Fully justify your answer. [4 r	nark

Turn over for the next question



3	The line L has equation $4x - 5y = 8$	
3 (a) (i)	Find the gradient of L . [2 marks]	
	Answer	
3 (a) (ii)	Find the coordinates of the point where L intersects the x -axis. [2 marks]	
	Answer _	



3	(b)	The point A has coordinates $(4,9)$ and the point B has coordinates $(2,k)$		b
		The line that passes through ${\it A}$ and ${\it B}$ is perpendicular to the line ${\it L}$.		
3	(b) (i)	Find the value of k .	[3 marks]	
			_	
		k =		
3	(b) (ii)	Find the equation of the line that passes through A and B .		
		Give your answer in the form $ax + by + c = 0$, where a , b and c are integers.	[3 marks]	
		Answer		10
		Turn over for the next question		

4 (a)	The n th term of a sequence is u_n		
	This sequence is defined by where k is a constant.	$u_{n+1} = ku_n + 17$	
	The first two terms of this sequence	e are $u_1 = 5$ and $u_2 = 23$	
4 (a) (i)	Find the value of k .		[2 marks]
		k =	
4 (a) (ii	Find the values of u_3 and u_4		[2 marks]
	$u_3 = $	<i>u</i> ₄ =	

4 ('b)	The <i>n</i> th te	rm of a	different	sequence	is t
- 1	N)	1110 <i>11</i> 111 10	iiii Oi a	uniciciii	SEGUETICE	$10 \iota_n$

This sequence is defined by

$$t_{n+1} = (t_n)^2 - 12$$

Find, using algebra, all the values of t_1 for which all the terms of this sequence are the same.

[3 marks]

Answer ____

7

Turn over for the next question



5 (a)	The expression $(1 + 2x)^3$ can be written in the form $1 + ax + bx^2 + 8x^3$ where a and b
	are integers.

Find the values of a and b.

[3 marks]

5 ((b)	Given that $y = (1 + 2x)^3$ find $\frac{dy}{dx}$	
			[3 marks]

$$\frac{dy}{dx} =$$

10

5 (c)	The curve C has equation $y = (1 + 2x)^3$ and the line L has equation $y = 3 - 10x$
	Show that there is ${\bf no}$ point on C at which the tangent to C is parallel to L . [4 marks]


Turn over for the next question




6		The curve C with equation $y = f(x)$ passes through the point $(3, -45)$	
•	(-)	dy dy dy	
6	(a)	Given that $\frac{dy}{dx} = 12x^2 - 26x - 12$, find the equation of C .	narks]
		L	
		Answer	
6	(b)	Write the equation of C in the form $y = x(ax + b)(cx + d)$, where a, b, c and d are	
6	(b)	integers.	narks]
6	(b)	integers.	narks]
6	(b)	integers.	narks]
6	(b)	integers.	narks]
6	(b)	integers.	narks]
6	(b)	integers.	narks]
6	(b)	integers.	narks]
6	(b)	integers.	narks]
6	(b)	integers.	narks]
6	(b)	integers. [3 m	narks]
6	(b)	integers. [3 m	narks]
6	(b)	integers. [3 m	narks]



10



[3 marks]



Turn over for the next question



7	The terms of a sec	quence with first term u	are given by
		9	1

$$u_1 = 672$$

$$u_{n+1} = -\frac{3}{4}u_n$$

7	(a) (i)	Find the third term of the sequence.
	(u, (i,	i ind the tima term of the sequence.

[1 mark]

$u_2 =$			

7 (a) (ii) State whether this is an arithmetic sequence or a geometric sequence.

[1 mark]

7 **(b) (i)** State how you know that the series $u_1 + u_2 + u_3 + \dots$ converges.

[1 mark]



Do not write outside the box


7 (b) (ii)	Find the sum to infinity of this series. [2 marks	s]
		_
		_
		_ _
	Answer	_
	Turn over for the next question	



8 (a) (i)	Find $\int (x^2 - 8x + \frac{1}{x^2} + 7) dx$	[4 marks]
	American	
	Answer	
8 (a) (ii)	Evaluate $\int_{2}^{5} (x^2 - 8x + \frac{1}{x^2} + 7) dx$	
o (u) (ii)	Evaluate J_2 ($x = 6x + \frac{1}{x^2} + \frac{7}{7} ux$	[2 marks]
	Answer	



8 (b) The curve with equation $y = x^2 - 8x + \frac{1}{x^2} + 7$ for x > 0 is drawn below.



A straight line intersects the curve at the points A(2, -4.75) and B(5, -7.96)

Find the exact area of the shaded region bounded by the curve $y = x^2 - 8x + \frac{1}{x^2} + 7$ and the line AB.

[4 marks]

-		

Answer _____

10



9	A railway company bases the cost of running its trains on the speed at which they travel.
	One of the company's trains travels from New York to Chicago at a steady speed of ν kilometres per hour.
	The cost per kilometre in dollars, C , for the journey is given by
	$C = \frac{v}{25} + \frac{100}{v}$
9 (a)	Find the value of v for which C is a minimum.
	Fully justify your answer. [7 marks]
	v =



9

9 (b)	The distance by train from New York to Chicago is 1270 kilometres.	
	Calculate the minimum total cost of the journey.	
	Calculate the minimum total cost of the journey.	[2 marks]
	Answer	
	7 416 4061	
	Turn over for the next question	





10 (a)	The expressions $x^2 + bx + c$ and $x^2 - 3mx + 2n$ have a common factor of $(x - p)$, where b , c , m and n are positive.
	Show that $p = \frac{2n-c}{b+3m}$ [3 marks]



Find the range of possible values of k .	
Show clearly each step of your working.	[6 mark
Answer	
END OF QUESTIONS	



There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED Copyright information For confidentiality purposes, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the

For confidentiality purposes, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.oxfordaqaexams.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 Oxford International AQA Examinations and its licensors. All rights reserved.



Do not write outside the box