

OXFORD

INTERNATIONAL
AQA EXAMINATIONS

INTERNATIONAL AS MATHEMATICS MA01

(9660/MA01) Unit P1 Pure Mathematics

Mark scheme

June 2023

Version: 1.0 Final



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from oxfordaqaexams.org.uk

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2023 Oxford International AQA Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

M	Mark is for method
m	Mark is dependent on one or more M marks and is for method
A	Mark is dependent on M or m marks and is for accuracy
B	Mark is independent of M or m marks and is for method and accuracy
E	Mark is for explanation
✓ or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
–x EE	Deduct x marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)

Q	Answer	Marks	Comments
1(a)(i)	$16a^{\frac{11}{6}}$	B1	
		1	

Q	Answer	Marks	Comments
1(a)(ii)	$2a^{\frac{5}{12}}$	B1	
		1	

Q	Answer	Marks	Comments
1(b)(i)	$[500 \times 5^p \times] x^{2p+6}$ $[p =] -3$	M1 A1	<p>Correctly applies index rules to obtain a correct single power of x</p> <p>Could be seen embedded in a product. Ignore terms in their product that do not include x</p> <p>PI by $2p + 6$ or $2p = -6$ or correct answer seen.</p> <p>CAO</p>
		2	

Q	Answer	Marks	Comments
1(b)(ii)	$\left[\frac{500}{5^3} = \frac{500}{125} \right] = 4$	B1	CAO
		1	

	Question 1 Total	5	
--	------------------	---	--

Q	Answer	Marks	Comments
2(a)	$[QR =] \sqrt{(4-14)^2 + (9-(-3))^2}$ $\sqrt{244} \quad \text{or} \quad 2\sqrt{61}$	<p>M1</p> <p>A1</p>	<p>oe PI by 15.6[2049...] or correct final answer</p> <p>ISW Ignore decimal value if given as well.</p>
		2	

Q	Answer	Marks	Comments
2(b)	<p>[Mid-Point of QR =] (9, 3)</p> <p>[Gradient of QR =] $\frac{9-(-3)}{4-14}$</p> <p>[Gradient of l =] $\frac{5}{6}$</p> <p>$\frac{y-3}{x-9} = \frac{5}{6}$ oe</p> <p>and</p> <p>$y = \frac{5}{6}x - \frac{9}{2}$</p>	<p>B1</p> <p>M1</p> <p>A1ft</p> <p>A1</p>	<p>PI in later working.</p> <p>oe Correct method for finding the gradient of QR PI by $-\frac{6}{5}$ oe seen.</p> <p>oe Possibly seen in later working. ft their gradient of QR</p> <p>Forms a correct equation for l before the given answer. May see $y = \frac{5}{6}x + p$ and substitution of coordinates of the mid-point of QR to find p but must be a complete method.</p> <p>AG Must be convincingly shown</p>
		4	

Q	Answer	Marks	Comments
2(c)	$[k =] \frac{5}{6} \times 30 - \frac{9}{2}$	M1	Substitutes $x = 30$ into the equation of l PI by correct value of k
	$[k =] \frac{41}{2}$ or 20.5	A1	CAO
	$\left[20.5 = \frac{1}{4} \times 30 + d \right]$		
	$[d =] 13$	B1ft	ft follow through their $k = 7.5$ Substitutes their k into the equation of the line and evaluates d Condone equivalent fraction
2(c) ALT	$\left[\frac{5}{6}x - \frac{9}{2} = \frac{1}{4}x + d \text{ and } x = 30 \Rightarrow \right]$	M1	oe Equates equations of both lines and $x = 30$ substituted into a correct equation. PI by correct value of d
	$\frac{5}{6}(30) - \frac{9}{2} = \frac{1}{4}(30) + d$ or $\frac{7}{12}(30) = d + \frac{9}{2}$		
	$[d =] 13$	A1	CAO
	$[k =] \frac{41}{2}$ or 20.5	B1ft	ft follow through their $7.5 + d$ Substitutes their d into the equation of the line and evaluates k Condone equivalent fraction
		3	

	Question 2 Total	9	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
3(a)	$[S_{30} =] \frac{1}{2} \times 30 \times (2a + (30 - 1)d)$ $[= 30a + 435d]$ <p>or</p> $[S_{10} =] \frac{1}{2} \times 10 \times (2a + (10 - 1)d)$ $[= 10a + 45d]$ $\frac{1}{2} \times 30 \times (2a + (30 - 1)d)$ $- \frac{1}{2} \times 10 \times (2a + (10 - 1)d)$ $(30a + 435d) - (10a + 45d) \quad [= 522]$ $20a + 390d = 522$ <p>and</p> $10a + 195d = 261$	<p>M1</p> <p>M1</p> <p>A1</p>	<p>oe Could be embedded. Correct expression for S_{30} or S_{10} with values substituted simplified or unsimplified.</p> <p>oe Correct expression for $S_{30} - S_{10}$</p> <p>Integer multiple of final answer, before given answer</p> <p>AG Must be convincingly shown</p>
		3	

Q	Answer	Marks	Comments
4(a)	$\left[(1+6x)^7 = \right]$ $\left[(1)^7 + 7(1)^6(6x) + 21(1)^5(6x)^2 + 35(1)^4(6x)^3 \right]$ $[a =] 42$ $[b =] 756$	<p>M1</p> <p>For either $[1], 7, 21, [35]$ oe unsimplified.</p> <p>or $\begin{pmatrix} 7 \\ 1 \end{pmatrix} (1)^6 (6x)$ or $\begin{pmatrix} 7 \\ 2 \end{pmatrix} (1)^5 (6x)^2$ oe x not needed. PI</p> <p>A1</p> <p>Condone $42x$ Possibly embedded in expansion.</p> <p>A1</p> <p>Condone $756x^2$ Possibly embedded in expansion.</p>	
		3	

Q	Answer	Marks	Comments
4(b)	$\frac{1}{2} \times 7560 [x^3]$ <p>or $(-k) \times 756 [x^3]$</p> $3780 [x^3]$ $(3780 - 756k) [x^3] = 1512 [x^3]$ $[k =] 3$	<p>M1</p> <p>ft their b from part 4(a). Multiplying together two relevant pairs of terms. Condone if seen embedded in a full or partial expansion.</p> <p>M1</p> <p>oe Correct equation.</p> <p>A1</p> <p>CAO</p>	
		3	

	Question 4 Total	6	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
5(a)	$h = 0.4$	B1	PI
	$\left[\text{With } f(x) = 8^{\sqrt{x}} \right]$ $\left[I \approx \frac{h}{2} \{ \dots \} \right]$ $\left[\{ \dots \} = \right] f(1) + f(3)$ $+ 2(f(1.4) + f(1.8) + f(2.2) + f(2.6))$	M1	oe Summing the areas of the trapezia.
	$\left[\{ \dots \} = \right] 8 + 36.6604 \dots$ $+ 2 \times (11.7098 \dots + 16.2787 \dots$ $+ 21.8523 \dots + 28.5883 \dots)$	A1	oe Accept rounded or truncated to two decimal places. PI by AWRT 40.3
	$\left[I \approx 0.2 \times 201.5191 \dots = \right] 40.3$	A1	CAO Must be 40.3
		4	

Q	Answer	Marks	Comments
5(b)(i)	$\left[8^{\left(\frac{1}{3} + \sqrt{x}\right)} = \right] 8^{\frac{1}{3}} \left[\times 8^{\sqrt{x}} \right] \quad \text{or} \quad 2 \left[\times 8^{\sqrt{x}} \right]$	B1	PI by correct scale factor of stretch.
	Stretch in the y -direction.	E1	Both 'stretch' and 'direction' needed.
	[Scale] factor 2	E1	Accept 'sf'. Allow $8^{\frac{1}{3}}$ for 2
		3	

Q	Answer	Marks	Comments
5(b)(ii)	$\left[\int_1^3 8^{\left(\frac{1}{3} + \sqrt{x}\right)} dx = 2 \int_1^3 8^{\sqrt{x}} dx \approx \right] 2 \times 40.3$	M1	Their trapezium rule value multiplied by a scale factor PI by 80.6 or $2 \times$ part (a) trapezium rule value but not 80.2 (from calculator use)
	80.6	A1	CAO Second use of trapezium rule is M0 A0
		2	

	Question 5 Total	9	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
6(a)	$[f(4) = 4^3 + a \times 4^2 - 6b \times 4 + 7$ $64 + 16a - 24b + 7 = 23$ $71 + 16a - 24b = 23 \quad \text{oe}$ <p>and</p> $2a - 3b = -6$	<p>M1</p> <p>A1</p>	<p>Correctly substitutes $x = 4$ into $f(x)$</p> <p>Must use the Remainder Theorem. AG Must be convincingly shown</p> <p>Expression for $f(4)$ set equal to 23 with products and powers evaluated and AG Must be at least one extra line of working given before AG</p>
		2	

Q	Answer	Marks	Comments
6(b)	$\left[\frac{dy}{dx} = \right] 3x^2 + 2ax - 6b$ $3(-5)^2 + 2a(-5) - 6b = 21$ $[75 - 10a - 6b = 21]$ $5a + 3b = 27$	<p>M1</p> <p>m1</p> <p>A1</p>	<p>Condone one error in a term or one term omitted.</p> <p>Substitutes $x = -5$ into their derivative and sets equal to 21</p> <p>CAO oe must be in the correct form</p>
		3	

Q	Answer	Marks	Comments
6(c)	$[a =] 3$ and $[b =] 4$	B1	CAO
		1	

Q	Answer	Marks	Comments
6(d)	$[g'(x) =] 48 + 2x - x^2$ $48 + 2x - x^2 > 0$ $[x =] -6 \quad \text{and} \quad [x =] 8$ $-6 < x < 8$ $-6 < x < -4 \quad \text{or} \quad 2 < x < 8$	M1 M1 A1 A1ft M1 A1	Allow one error in a term or one term omitted. PI correct inequality or correct critical values. Condone given as equality. ft their $g'(x)$ Both correct critical values. Correct solution to $g'(x) > 0$ PI by both correct intervals in final answer. ft their two critical values. M1 : One correct interval. Ignore other incorrect intervals given. A1 : Both correct intervals and no others. Do not condone 'and' for 'or'.
		6	
	Question 6 Total	12	

Q	Answer	Marks	Comments
7(a)	$[y =] x^2 - 6x^{\frac{4}{3}} + 16$ $\left[\frac{dy}{dx} =\right] 2x - 8x^{\frac{1}{3}}$	B1 B1ft	Correct expansion. PI by correct derivative. oe Simplified or unsimplified. ft their expansion provided it contains a fractional power of x
		2	

Q	Answer	Marks	Comments
7(b)	$2x - 8x^{\frac{1}{3}} = 0$ $2x\left(1 - 4x^{-\frac{2}{3}}\right) = 0 \Rightarrow x = 0$ When $x = 0$, $y = 16$ and $(0, 16)$ $x^{\frac{2}{3}} - 4 = 0$ or $1 - 4x^{-\frac{2}{3}} = 0$ or $x^2 = 64$ $[x_Q =] 8$ $(8, -16)$	M1 A1 M1 A1 A1	oe ft their first derivative equal to zero. Statement that $x = 0$ from correct first derivative Correct coordinates of P Condone not given as coordinates but must be clearly identified. oe PI Correct x -coordinate of Q Correct coordinates of Q
		5	

Q	Answer	Marks	Comments
7(c)(i)	$\left[\frac{d^2y}{dx^2} =\right] 2 - \frac{8}{3}x^{-\frac{2}{3}}$	B1ft	oe ft their $\frac{dy}{dx}$ provided it contains a fractional power of x
		1	

Q	Answer	Marks	Comments
8	$\left[\frac{6x+5x^2}{x^2\sqrt{x}} = \frac{6}{x\sqrt{x}} + \frac{5}{\sqrt{x}} = \right] 6x^{-\frac{3}{2}} + 5x^{-\frac{1}{2}}$ $\left[\int \frac{6x+5x^2}{x^2\sqrt{x}} dx = \right]$ $-12x^{-\frac{1}{2}} + 10x^{\frac{1}{2}} [+c]$ $\left[\int_{a^2}^{25a^2} \frac{6x+5x^2}{x^2\sqrt{x}} dx = \right]$ $\left(-12(25a^2)^{-\frac{1}{2}} + 10(25a^2)^{\frac{1}{2}} \right)$ $- \left(-12(a^2)^{-\frac{1}{2}} + 10(a^2)^{\frac{1}{2}} \right) [=44]$ $-\frac{12}{5a} + 50a + \frac{12}{a} - 10a [=44]$ or $\frac{48}{5a} + 40a [=44]$ $200a^2 - 220a + 48 = 0$ or $50a^2 - 55a + 12 = 0$ $[(10a-3)(5a-4)=0 \Rightarrow]$ $a = \frac{3}{10} \quad \text{or} \quad a = \frac{4}{5}$	<p>B1</p> <p>B2,1ft</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>Correctly written as a sum of powers of x PI by correct integration</p> <p>oe ft their $6x^{-\frac{3}{2}} + 5x^{-\frac{1}{2}}$ provided each term they integrate has a fractional powers of x B2 both terms correct or B1 for one term correct. Simplified or unsimplified.</p> <p>Forms $F(25a^2) - F(a^2)$ for their integration.</p> <p>oe Simplifies the powers of a and removes the brackets.</p> <p>oe Correctly rearranges to form a quadratic equation in a. Must '=' 0' PI by correct final answer.</p> <p>CAO oe Both correct values.</p>
		7	

	Question 8 Total	7	
--	-------------------------	----------	--

[illegible]

9(a) ALT	$\left[u_1 = 27^{2p+1} \right] 3^{6p+3} \text{ or } 3^{3(2p+1)}$ $\left[r = \frac{u_2}{u_1} = \frac{3^{18p}}{27^{2p+1}} = \frac{3^{18p}}{3^{6p+3}} \right] 3^{12p-3}$ or $\left[r = \frac{u_3}{u_2} = \frac{3^{6p+1}}{3^{18p}} \right] 3^{1-12p}$ $3^{6p+1} = 3^{6p+3} \times (3^{12p-3})^2$ or $3^{6p+1} = 3^{6p+3} \times 3^{24p-6}$ or $3^{6p+1} = 3^{6p+3} \times (3^{1-12p})^2$ or $3^{6p+1} = 3^{6p+3} \times 3^{2-24p}$ $6p + 1 = 30p - 3$ or $6p + 1 = 5 - 18p$ or $24p = 4$ and $p = \frac{1}{6}$	B1	PI Writing u_1 as a power of 3
		M1	oe PI by $r^2 = \frac{1}{9}$ A correct expression for the common ratio (possibly squared) as a single power of 3
		M1	oe PI Correct equation in terms of p only for $u_3 = u_1 \times r^2$ Allow u_1 and r^2 unsimplified.
		A1	oe Correctly equates powers of 3 to form a linear equation in p before AG CSO
		4	

Q	Answer	Marks	Comments
9(b)	$[u_1 = a =] \quad 81$ $r = \frac{1}{3}$ $\left[[54 \times] \sum_{n=k+1}^{6k} u_n = [54 \times] \left(\sum_{n=1}^{6k} u_n - \sum_{n=1}^k u_n \right) \right]$ $[54 \times] \left(\frac{81 \left(1 - \left(\frac{1}{3} \right)^{6k} \right)}{1 - \frac{1}{3}} - \frac{81 \left(1 - \left(\frac{1}{3} \right)^k \right)}{1 - \frac{1}{3}} \right)$ $\left[54 \sum_{n=k+1}^{6k} u_n = \right]$ $81 \left(81 \left(\frac{1}{3} \right)^k \left(1 - \left(\frac{1}{3} \right)^{5k} \right) \right)$ <p>or</p> $6561 \times \left(\frac{1}{3} \right)^k \left(1 - \left(\frac{1}{3} \right)^{5k} \right)$ $\left[54 \sum_{n=k+1}^{6k} u_n = \right] \quad 3^{8-k} \left(1 - 3^{-5k} \right)$	<p>B1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A2,1</p>	<p>Allow $a = 3^4$</p> <p>oe Correct substitution into $\sum_{n=1}^{6k} u_n - \sum_{n=1}^k u_n$</p> <p>ft their a and r</p> <p>Allow $a = 3^4$</p> <p>oe</p> <p>Multiplication of $\sum_{n=1}^{6k} u_n - \sum_{n=1}^k u_n$ by 54, fractions cleared and $\left(\frac{1}{3} \right)^k$ taken out as a factor.</p> <p>In correct form. A1 b and c or b and d correct. A2 Fully correct answer.</p>
		6	

	Question 9 Total	10	
--	-------------------------	-----------	--