

INTERNATIONAL A-LEVEL MATHEMATICS

MA04

(9660/MA04) Unit S2 Statistics

Mark scheme

January 2024

Version: 1.0 Final



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [oxfordaqa.com](https://www.oxfordaqa.com)

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2024 OxfordAQA International Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

M	Mark is for method
m	Mark is dependent on one or more M marks and is for method
A	Mark is dependent on M or m marks and is for accuracy
B	Mark is independent of M or m marks and is for method and accuracy
E	Mark is for explanation
✓ or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
–x EE	Deduct x marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)
ISW	Ignore subsequent working

Q	Answer	Marks	Comments
1(a)	Exponential	B1	
		1	

Q	Answer	Marks	Comments
1(b)	$\left[\frac{1}{1/7} = \right] 7$	B1	
		1	

Q	Answer	Marks	Comments
1(c)	$[P(X < 7) - P(X < 2)]$ $= \left(1 - e^{-\frac{1}{7} \times 7} \right) - \left(1 - e^{-\frac{1}{7} \times 2} \right)$ $= 0.3836$	<p>M1</p> <p>A1</p>	<p>PI</p> <p>Attempts to find correct probability using cdf of exponential or integration of pdf</p> <p>AWRT 0.384</p>
		2	

Q	Answer	Marks	Comments
1(d)	$\left[P(X < a) = 0.8 \right] = \left(1 - e^{-\frac{1}{7}a} \right) = 0.8$	M1	oe, PI
	= 11.27	A1	AWRT 11.3
		2	

Q	Answer	Marks	Comments
1(e)	$\left[P(X > 8 X > 5) = P(X > 3) = \right] e^{-\frac{1}{7} \times 3}$	M1	oe, PI
	= 0.6514	A1	CAO
		2	

	Question 1 Total	8	
--	------------------	---	--


Q	Answer	Marks	Comments
2(a)	$H_0 : \mu = 300$ $H_1 : \mu > 300$ $\bar{X} \sim N\left(300, \frac{40^2}{200}\right)$ $z = \frac{306 - 300}{\frac{40}{\sqrt{200}}}$ $z = 2.1213\dots$ $z_{\text{critical}} = 2.0537$ Reject H_0 as $z_{\text{critical}} < z$ or $2.0537 < 2.1213\dots$ or $ z > 2.0537$ There is sufficient evidence to support the claim that the average time for which the app is used has increased since the new version was released [at the 2% level of significance]	B1 B1 M1 A1 B1 A1ft E1	PI by correct standardisation formulae $z = \frac{306 - 300}{\text{their } \frac{\sigma}{\sqrt{n}}}$ PI by correct z or probability AWRT 2.12 or exact value $\frac{3\sqrt{2}}{2}$ or $P(z > 2.1213\dots)$ AWFW 0.0169... to 0.0170 or comparison of probability 0.0169... to $0.0170 < 0.02$ Allow 'Accept H_1 ' Comment about H_0 and comparison 0.0169... to $0.0170 < 0.02$ Correct conclusion based upon ft their z Correct statement must be in context and must follow from fully correct solution Condone definite statement
		7	

Q	Answer	Marks	Comments
2(b)	Central limit theorem (CLT) states that [when the sample size is large enough], the sample mean will be approximately normally distributed	B1	Allow $n \geq 30$
	A sample of 200 is large enough for the CLT to apply	B1	
		2	
	Question 2 Total	9	

Q	Answer	Marks	Comments
3(a)	$\left[\int_0^1 \frac{x^2}{8} dx \right] = \left[\frac{x^3}{24} \right]_0^1$	M1	PI Correct integration and limits
	$= \frac{1}{24}$	A1	oe in exact form
		2	

Q	Answer	Marks	Comments
3(b)	$\frac{1}{24} + \int_1^6 \left(k(x-1) + \frac{1}{8} \right) dx = 1$	B1	Setting an integral and fraction summing to 1 oe PI
	$\frac{1}{24} + \left[\frac{k(x-1)^2}{2} + \frac{1}{8}x \right]_1^6 = 1$		
	$\frac{1}{24} + \left(\frac{25k}{2} + \frac{3}{4} \right) - \left(0 + \frac{1}{8} \right) = 1$	M1	For correct integration with attempt to substitute limits oe
	$\frac{25k}{2} = \frac{1}{3}$		
	$k = \frac{2}{75}$	A1	AG must be convincingly shown
		3	

Q	Answer	Marks	Comments
3(c)		B1 B1 B1	Curve from (0,0) to (1, 15/120) can be identified as coordinates or from their scale Straight line from (1, 15/120) to (6, 31/120) can be identified as coordinates or from their scale Straight line from (6,0) to (8,0)



		3	
--	--	---	--

Q	Answer	Marks	Comments
3(d)	$\frac{x^3}{24}$ $\int_1^x \left(\frac{2(x-1)}{75} + \frac{1}{8} \right) dx + \frac{1}{24}$ $= \left[\frac{(x-1)^2}{75} + \frac{1}{8}x \right]_1^x + \frac{1}{24}$ $= \frac{(x-1)^2}{75} + \frac{1}{8}x - \frac{1}{12}$ $F(x) = \begin{cases} 0 & x < 0 \\ \frac{x^3}{24} & 0 \leq x < 1 \\ \frac{(x-1)^2}{75} + \frac{1}{8}x - \frac{1}{12} & 1 \leq x \leq 6 \\ 1 & x > 6 \end{cases}$	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>Correct integration</p> <p>condone omission of $\frac{1}{24}$</p> <p>$F(x) = \frac{1}{75}x^2 + \frac{59}{600}x - \frac{7}{100}$ from</p> <p>$f(x) = \frac{2}{75}x + \frac{59}{600}$</p> <p>oe Fully correct F(x)</p>
		4	
	Question 3 total	12	

Q	Answer	Marks	Comments
4(a)	$B(100, 0.05)$ It is the only one with large n and small p	B1 B1	Condone <i>F</i> If B0 B0 , then allow SC1 for comparing means and variances
		2	

Q	Answer	Marks	Comments
4(b)(i)	$P(X = 3) = \frac{e^{-1.8} \times 1.8^3}{3!}$ $= 0.161$	M1 A1	PI AWRT 0.161
		2	

Q	Answer	Marks	Comments
4(b)(ii)	$\lambda \left[= \frac{1.8}{3} \right] = 0.6$ $P(1 < X < 4) = P(X \leq 3) - P(X \leq 1)$ $= 0.9966 - 0.8781$ $= 0.119$	B1 M1 M1 A1	PI PI At least one correct probability or finds $P(X = 2)$ and $P(X = 3)$ Correct method AWRT 0.119
		4	

Q	Answer	Marks	Comments
4(c)(i)	$\lambda [= 1.8 + 2.7] = 4.5$	B1	
		1	

Q	Answer	Marks	Comments
4(c)(ii)	$P(G < 8) = 0.9134 < 0.95$ $P(G < 9) = 0.9597 > 0.95$ Hence $a = 9$	M1 A1 B1	Sight of at least relevant probability statement Sight of both probabilities and comparison to 0.95
		3	

	Question 4 Total	12	
--	------------------	----	--

Q	Answer	Marks	Comments
5(a)	$H_0 : \mu = 13$ $H_1 : \mu \neq 13$ $\bar{x} = \frac{129.5}{10} = 12.95$ $s^2 = \frac{1}{10-1} \left(1677.05 - \frac{129.5^2}{10} \right)$ $= \frac{1}{360} = 0.002\dot{7}$ $\bar{X} \sim N \left(13, \frac{0.002\dot{7}}{10} \right)$ $t = \frac{12.95 - 13}{\sqrt{\frac{0.002\dot{7}}{10}}}$ $= -3$ $\nu = 9 \Rightarrow t_{\text{critical}} = \pm 3.250$ Do not reject H_0 as $ t < 3.250$ Insufficient evidence to suggest that Alice's mean maximum exposure time has changed at the 1% level of significance	B1 B1 M1 A1 M1 M1 A1 B1 A1ft E1	Both hypotheses Attempt at variance formula Condone one error PI by correct answer AWRT 0.00278 Accept $s = 0.0527[0462767]$ $\bar{X} \sim N \left(13, \frac{s^2}{10} \right)$ PI Calculates z with their s^2 or sight of $p=0.00747$ oe Follow through their t and t_{crit} provided signs are consistent Implied by correct conclusion in context (or comparison of p to 0.005) Must be in context, must not be definite and all the previous 9 marks must have been awarded
		10	

Q	Answer	Marks	Comments
5(b)	He uses a different significance level or He uses a one-tailed test	E1	A correct suggestion
		1	

	Question 5 Total	11	
--	-------------------------	-----------	--

Q	Answer	Marks	Comments
6(a)	$\bar{V} \sim N\left(502, \frac{2.7^2}{30}\right)$	B1	PI
	$P(\bar{V} < 501) = P\left(z < \frac{501 - 502}{\frac{2.7}{\sqrt{30}}}\right)$	M1	PI
	$P(z < -2.0286\dots)$		
	$= 1 - 0.97882$ [from tables]	M1	Calculator method gives $1 - 0.97875$ PI $1 - 'p'$ where p is from a standardised value using $n = 30$
	$= 0.0212$ (to 4 dp)	A1	AWFW 0.02118 to 0.0213 SC1 AFWF 0.355 to 0.356
		4	

Q	Answer	Marks	Comments
6(b)	$P(V > 496) = P\left(z > \frac{496 - 502}{2.7}\right)$ $= P(z > -2.22\dots)$ $= 0.98679 \quad [\text{from tables}]$ $= P(\text{all } 30) = 0.98679^{30}$ $= 0.671$	<p>M1</p> <p>A1</p> <p>m1</p> <p>A1</p>	<p>PI</p> <p>AWRT 0.987 PI</p> <p>p^{30}</p> <p>AWFW 0.671 to 0.673</p>
		4	

Q	Answer	Marks	Comments
6(c)	Let $M = V + W + W + Y$ $M \sim N(502 + 251 + 251 + 503,$ $2.7^2 + 1.5^2 + 1.5^2 + 2^2)$ $M \sim N(1507, 15.79)$ $P(M > a) = 0.95$ $z = -1.6449$ $\frac{a - 1507}{\sqrt{15.79}} = -1.6449$ $a = 1500$ [grams]	M1 A1 B1 M1 A1	Correct method for finding the mean or variance of M PI by sight of 1507 or 15.79... PI Fully correct distribution for M Allow $\pm z$ value to at least 4sf PI Standardising with compatible signs and use of their σ CAO
		5	

	Question 6 Total	13	
--	-------------------------	-----------	--

Q	Answer	Marks	Comments
7(a)	$\frac{d}{dx}F(x) = \frac{4}{x^3} \quad \text{or} \quad \frac{d}{dx}F(x) = \frac{x}{48} + \frac{1}{32}$ $E(X) = \int xf(x)dx$ $= \int_2^4 \frac{4}{x^2} dx + \int_4^8 \left(\frac{x^2}{48} + \frac{x}{32} \right) dx$ $= \left[\frac{-4}{x} \right]_2^4 + \left[\frac{x^3}{144} + \frac{x^2}{64} \right]_4^8$ $= 1 + \frac{139}{36}$ $= \frac{175}{36}$ $E(X^2) = \int x^2 f(x) dx$ $= \int_2^4 \frac{4}{x} dx + \int_4^8 \left(\frac{x^3}{48} + \frac{x^2}{32} \right) dx$ $= \left[4\ln(x) \right]_2^4 + \left[\frac{x^4}{192} + \frac{x^3}{96} \right]_4^8$ $= 4\ln(2) + \frac{74}{3}$ $E(X^2) - E(X)^2 = 4\ln(2) + \frac{74}{3} - \left(\frac{175}{36} \right)^2$ $= 4\ln(2) + \frac{1343}{1296}$	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Correct integration with correct limits of their $xf(x)$</p> <p>correct integration with correct limits of their $x^2f(x)$ oe</p> <p>correct use of their $E(X^2)$ and $E(X)^2$</p> <p>AG Must be convincingly shown</p>
		7	

Q	Answer	Marks	Comments
7(b)	$4\text{Var}(X) + 9\text{Var}(Y)$	M1	AWRT
	$= 4 \times 3.81 + 9 \times 2.5^2$	A1	
	$= 71.5$		
		2	
	Question 7 Total	9	

Q	Answer	Marks	Comments
8	$H_0 : p = 0.5$ $H_1 : p \neq 0.5$ $X \sim B(20, 0.5)$ $P(X \geq 15) = 1 - P(X \leq 14)$ $= 0.0207$ $0.0207 > 0.015$ Do not reject H_0 There is sufficient evidence at the 3% level to suggest that the student has randomly selected their answers	B1 B1 M1 A1 B1ft E1ft	Hypotheses that model the situation PI by correct probability or $X \sim B(20, p)$ AWFW 0.0206 to 0.0207 or a CR= {0,1,2,3,4, 16,17,...,20} Compares their probability with 0.015 (allow comparison to 0.03 if their alternative hypothesis is 1 tailed) Or compares 15 to a correct CR Cannot be definitive Must follow a full hypothesis test ft their conclusion consistent with their model
		6	
	Question 8 Total	6	