

OXFORD

INTERNATIONAL
AQA EXAMINATIONS

INTERNATIONAL A-LEVEL MATHEMATICS

MA03

(9660/MA03) Unit P2 – Pure Mathematics

Mark scheme

June 2022

Version: 1.0 Final



206XMA03/MS

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [oxfordaqaexams.org.uk](https://www.oxfordaqaexams.org.uk)

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2020 Oxford International AQA Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

M	Mark is for method
m	Mark is dependent on one or more M marks and is for method
A	Mark is dependent on M or m marks and is for accuracy
B	Mark is independent of M or m marks and is for method and accuracy
E	Mark is for explanation
✓ or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
–x EE	Deduct x marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)

Q	Answer	Marks	Comments										
1(a)	<table><tr><td>x</td><td>y</td></tr><tr><td>0.1</td><td>$\sin\left(e^{0.1}\right) = 0.8935409$</td></tr><tr><td>0.3</td><td>$\sin\left(e^{0.3}\right) = 0.9756924$</td></tr><tr><td>0.5</td><td>$\sin\left(e^{0.5}\right) = 0.9969654$</td></tr><tr><td>0.7</td><td>$\sin\left(e^{0.7}\right) = 0.9034885$</td></tr></table>	x	y	0.1	$\sin\left(e^{0.1}\right) = 0.8935409$	0.3	$\sin\left(e^{0.3}\right) = 0.9756924$	0.5	$\sin\left(e^{0.5}\right) = 0.9969654$	0.7	$\sin\left(e^{0.7}\right) = 0.9034885$	B1	All 4 correct x values (and no extra used) PI by 4 correct y values
	x	y											
	0.1	$\sin\left(e^{0.1}\right) = 0.8935409$											
	0.3	$\sin\left(e^{0.3}\right) = 0.9756924$											
	0.5	$\sin\left(e^{0.5}\right) = 0.9969654$											
0.7	$\sin\left(e^{0.7}\right) = 0.9034885$												
		M1	At least 3 correct y values in exact form or decimals, rounded or truncated to 2 dp or better (in table or formula) (PI by AWRT correct answer)										
	$0.2\times\left[0.89\dots+0.97\dots+0.99\dots+0.90\dots\right]$	m1	Correct sub into formula with $h = 0.2$ OE and at least 3 correct y values either listed, with + signs, or totalled. (PI by AWRT correct answer)										
	$= 0.754$	A1	CAO , must see this value exactly and no error seen										
		4											
1(b)(i)	$f\left(x\right)=\sin\left(e^x\right)-3x+2$ $f\left(0.8\right)=0.39\dots$ $f\left(0.9\right)=-0.069\dots$ Change of sign, $0.8<\alpha<0.9$	M1	Or reverse Both values rounded or truncated to at least 1sf										
		A1	Must have both statement and interval in words or symbols or comparing 2 sides: at 0.8, $\sin\left(e^{0.8}\right)>3\times0.8-2$; at 0.9, $\sin\left(e^{0.9}\right)<3\times0.9-2$ (M1) Conclusion as before (A1)										
		2											
1(b)(ii)	$x_2 = 0.931$ $x_3 = 0.856$	B1 B1											
		2											
	Total	8											

Q	Answer	Marks	Comments
2(a)	$\left[\frac{dy}{dx} = \frac{(2x+5) \times (-3) - (1-3x) \times 2}{(2x+5)^2} \right]$ $= \frac{-17}{(2x+5)^2}$	M1 A1	or use of product rule PI by correct answer
		2	
2(b)	$\left[\frac{dy}{dx} = \frac{-17}{(2x+5)^2} \times \frac{(2x+5)}{(1-3x)} \right]$ $= \frac{-17}{(2x+5)(1-3x)}$	M1 A1	their (a) $\times \frac{2x+5}{1-3x}$ oe such as $\frac{A}{1-3x} - \frac{B}{5+2x}$ with $A < 0$ and $B > 0$ ft their (a) ACF such as $\frac{-3}{1-3x} - \frac{2}{2x+5}$
		2	
	Total	4	

Q	Answer	Marks	Comments
3(a)	$[16 \sin \theta + 30 \cos \theta =]$ $R \sin \theta \cos \alpha + R \cos \theta \sin \alpha$ $\alpha = 1.08$ $R = 34$ $[34 \sin(\theta + 1.08)]$	M1 A1 B1	Implied by $16 = R \cos \alpha$ and $30 = R \sin \alpha$
		3	
3(b)(i)	[Min value =] -34	B1ft	
		1	
3(b)(ii)	$[\theta =] 3.63$	B1	oe such as -2.65 Accept 3.6 Accept $1.5\pi - 1.08$ $[\pm 2n\pi]$
		1	
	Total	5	

Q	Answer	Marks	Comments
4(a)(i)	$18(-0.5)^3 + b(-0.5)^2 + c(-0.5) - 4 = 0$ $18\left(\frac{1}{3}\right)^3 + b\left(\frac{1}{3}\right)^2 + c\left(\frac{1}{3}\right) - 4 = -5$ $b - 2c = 25$ $b + 3c = -15$ $b = 9, \quad c = -8$	<p>M1</p> <p>A1</p> <p>m1</p> <p>A1</p>	<p>At least one correct substitution or M1 use of long division $9x^2 + \left(\frac{b-9}{2}\right)x$</p> <p>Both substitutions correct or A1 for $9x^2 + \left(\frac{b-9}{2}\right)x + \frac{1}{2}\left(c - \frac{b-9}{2}\right)$</p> <p>Attempt to solve their simultaneous equations</p> <p>Both values correct</p>
		4	
4(a)(ii)	$f(x) = (2x+1)(9x^2-4)$ $= (2x+1)(3x+2)(3x-2)$	<p>M1</p> <p>A1</p>	<p>PI</p> <p>Condone $p = 3$ and $q = 2$</p>
		2	
4(b)	$\frac{f(x)}{(3x+2)(x^2-2)} = \frac{(2x+1)(3x-2)}{(x^2-2)}$ $= \frac{6x^2 - x - 2}{x^2 - 2}$ $= \frac{6x^2 - 12 - x + 10}{x^2 - 2}$ $= 6 + \frac{10-x}{x^2-2}$	<p>M1</p> <p>A1</p>	<p>Substitutes their $f(x)$ and correctly cancels the factor of $3x + 2$ in numerator and denominator</p> <p>Be convinced</p>
		2	
	Total	8	

Q	Answer	Marks	Comments
6(a)	$= 1 + \left(-\frac{1}{3}\right) \times (-x) + \frac{\left(-\frac{1}{3}\right) \times \left(-\frac{4}{3}\right) \times (-x)^2}{2}$ $+ \frac{\left(-\frac{1}{3}\right) \times \left(-\frac{4}{3}\right) \times \left(-\frac{7}{3}\right) \times (-x)^3}{6}$ $= 1 + \frac{1}{3}x + \frac{2}{9}x^2 + \frac{14}{81}x^3$	<p>M1 A1</p> <p>A1</p>	<p>M1: At least 3 terms correct (unsimplified)</p> <p>A1: All terms correct (unsimplified)</p>
		3	
6(b)(i)	$\sqrt[3]{\frac{1}{1-2x}} = (1-2x)^{-\frac{1}{3}}$ $= 1 + \frac{1}{3} \times 2x + \frac{2}{9} \times (2x)^2 + \frac{14}{81} \times (2x)^3$ $= 1 + \frac{2}{3}x + \frac{8}{9}x^2 + \frac{112}{81}x^3$	<p>M1</p> <p>A1</p>	Substitutes 2x in to their (a)
		2	
6(b)(ii)	$-0.5 < x < 0.5$	B2	<p>oe such as $x < 0.5$</p> <p>B1 for $-0.5 \leq x \leq 0.5$</p>
		2	
6(c)	$[x = 0.1] \quad 1 + \frac{2}{3} \times 0.1 + \frac{8}{9} \times 0.1^2 + \frac{112}{81} \times 0.1^3$ $[= 1.0769...]$ $[x = 0.1] \quad \frac{1}{\sqrt[3]{1-0.2}} = \frac{1}{\sqrt[3]{0.8}} = \sqrt[3]{\frac{10}{8}}$ $\sqrt[3]{10} \quad [= 2 \times 1.0769...] = 2.154$	<p>B1ft</p> <p>M1</p> <p>A1</p>	<p>Substitutes $x = 0.1$ into their (b)(i)</p> <p>oe</p> <p>AWRT 2.154 from correct use of binomial expansion Value calculated using this binomial expansion is 2.153876543</p>
		3	
	Total	10	

Q	Answer	Marks	Comments
7(a)	Translation $\begin{bmatrix} 0 \\ k \end{bmatrix}$ or Stretch + either I or II $k = 1$ I: Parallel to y -axis II: SF $\frac{1}{3}$ [Followed by] Stretch + either I or II [Followed by] Translation $\begin{bmatrix} 0 \\ k \end{bmatrix}$ I: Parallel to y -axis II: SF $\frac{1}{3}$ $k = \frac{1}{3}$	M1 A1 M1 A1	
		4	
7(b)	$x = \frac{1 + \cos y}{3}$ $3x - 1 = \cos y$ $[f^{-1}(x) =] \cos^{-1}(3x - 1)$	M1 A1	Interchanging x and y
		2	
	Total	6	

Q	Answer	Marks	Comments
8(a)(i)	$6 = a(x^2 + 1) + x(bx)$ $a = 6, b = -6$ $\frac{6}{x^3 + x} = \frac{6}{x} - \frac{6x}{x^2 + 1}$	M1 A1	
		2	
8(a)(ii)	$\int \frac{6}{x^3 + x} [dx] = \int \frac{6}{x} - \frac{6x}{x^2 + 1} [dx]$ $= 6 \ln x - 3 \ln(x^2 + 1)$ $\left[\int_1^2 \frac{6}{x^3 + x} dx \right]$ $= (6 \ln 2 - 3 \ln 5) - (0 - 3 \ln 2)$ $= \ln \left(\frac{512}{125} \right)$	M1 A1 M1 A1	ft their a and b from (a)(i) Substitutes $x = 1$ and $x = 2$ into their integration, provided it is in the form $= a \ln x + b \ln(x^2 + 1)$ ACF such as $\ln \left(\frac{2^9}{5^3} \right)$
		4	

8(b)(i)	$\frac{d}{dy}(\cos y)^{-1} = -1(\cos y)^{-2} \times (-\sin y)$ $= \frac{\sin y}{\cos^2 y}$ $= \sec y \tan y$	M1 A1	Need to see an intermediate line of working
		2	
8(b)(ii)	$\left[\frac{du}{dx} = \right] \cos x$ $\left[\int \frac{u}{(1-u^2)^{1.5}} du = \right] \int \frac{\sin x \cos x}{(1-\sin^2 x)^{1.5}} dx$ $\left[= \int \frac{\sin x \cos x}{\cos^3 x} dx \right]$ $= \int \sec x \tan x dx$ $[= \sec x]$ $\int_0^{0.5} du = \int_0^{\frac{\pi}{6}} dx$ $\left[\int_0^{0.5} \frac{u}{(1-u^2)^{1.5}} du = \right] \frac{2}{\sqrt{3}} - 1$	B1 M1 A1 B1 A1	<p>All in terms of x condone omission of dx</p> <p>Change of limits, maybe seen earlier (may change back to u and not change limits)</p> <p>oe such as $\frac{1}{3}(2\sqrt{3}-3)$</p>
		5	
	Total	13	

Q	Answer	Marks	Comments
9(a)	$\frac{1}{(30-x)(10-x)} = \frac{A}{30-x} + \frac{B}{10-x}$ $1 = A(10-x) + B(30-x)$ $A = -\frac{1}{20}, \quad B = \frac{1}{20}$ $\frac{1}{(30-x)(10-x)} = -\frac{1}{20(30-x)} + \frac{1}{20(10-x)}$	<p>M1</p> <p>A1</p>	
		2	

9(b)	$\frac{dx}{dt} = k(30-x)(10-x)$ $\int \frac{1}{(30-x)(10-x)} dx = \int \frac{-1}{20(30-x)} + \frac{1}{20(10-x)} dx$ $= \frac{1}{20} \ln(30-x) - \frac{1}{20} \ln(10-x)$ $\frac{1}{20} \ln\left(\frac{30-x}{10-x}\right) = kt + c$ $[t=0, x=0 \Rightarrow] \quad c = \frac{1}{20} \ln 3$ $\frac{1}{20} \ln\left(\frac{30-x}{10-x}\right) = kt + \frac{1}{20} \ln 3$ $[t=2, x=6 \Rightarrow] \quad \frac{1}{20} \ln\left(\frac{24}{4}\right) = 2k + \frac{1}{20} \ln 3$ $k = \frac{1}{40} \ln 2$ $\frac{1}{20} \ln\left(\frac{30-x}{10-x}\right) = \left(\frac{1}{40} \ln 2\right)t + \frac{1}{20} \ln 3$ $\ln\left(\frac{30-x}{10-x}\right) = \frac{t}{2} \ln 2 + \ln 3$ $\left(\frac{30-x}{10-x}\right) = 3 \times 2^{0.5t}$ $x = \frac{30 \times (2^{0.5t} - 1)}{3 \times 2^{0.5t} - 1}$	B1 M1 m1 A1ft m1 M1 A1 m1 A1	 Uses their partial fractions to separate variables Attempt to integrate ft their A and B from (a) Attempt to find c Attempt to find k Both c and k correct Attempt to solve ACF
		9	
	Total	11	

Q	Answer	Marks	Comments
10(a)	$\frac{dx}{dt} = -3\cos^2 t \sin t$	M1	At least one derivative correct
	$\frac{dy}{dt} = -2\cos t \sin^2 t + (2 + \cos^2 t)\cos t$ $\left[= -2\cos t(1 - \cos^2 t) + 2\cos t + \cos^3 t = 3\cos^3 t \right]$	A1	Both derivatives correct
	$\frac{dy}{dx} = \frac{-3\cos^3 t}{3\cos^2 t \sin t}$	M1	Correct use of trig identity
	$\frac{dy}{dx} = -\frac{\cos t}{\sin t} = -\cot t$	A1	AG Be convinced
		4	
10(b)	gradient at $t = p$ is $-\frac{1}{-\cot p} \left[= \frac{\sin p}{\cos p} = \tan p \right]$	M1	Must be in terms of p
	$y - (2 + \cos^2 p)\sin p = \frac{\sin p}{\cos p}(x - \cos^3 p)$	A1	ACF $y = (\tan p)x + 2\sin p$
		2	
10(c)	$[x = 0 \Rightarrow] y - (2 + \cos^2 p)\sin p = \frac{\sin p}{\cos p}(-\cos^3 p)$	M1	Maybe seen in (b)
	$y = 2\sin p + \sin p \cos^2 p - \sin p \cos^2 p = 2\sin p$	A1	
	$[y = 0 \Rightarrow] -(2 + \cos^2 p)\sin p = \frac{\sin p}{\cos p}(x - \cos^3 p)$	M1	Maybe seen in (b)
	$x = -2\cos p - \cos^3 p + \cos^3 p = -2\cos p$	A1	
	$[AB =] \sqrt{4\sin^2 p + 4\cos^2 p}$	M1	
	$[AB =] 2$	A1	CAO
		6	
	Total	12	

Q	Answer	Marks	Comments
11(a)	$A(-2.5, 0), B(0, 5)$	B1	
		1	
11(b)(i)	$\left[\frac{dy}{dx} = \right] 2e^{-x} - (5 + 2x)e^{-x}$ $\left[\frac{dy}{dx} = -3e^{-x} - 2xe^{-x} \right]$	M1 A1	M1: $ae^{-x} + bxe^{-x}$ A1: $-3e^{-x} - 2xe^{-x}$ ACF
		2	
11(b)(ii)	$-3e^{-x} - 2xe^{-x} = 0$ $-e^{-x}(3 + 2x) = 0$ $[e^{-x} \neq 0 \Rightarrow] 3 + 2x = 0$ $(-1.5, 2e^{1.5})$	M1 A1	ft their [simplified] derivative oe
		2	
11(b)(iii)	$\left[\frac{d^2y}{dx^2} = 3e^{-x} + 2xe^{-x} - 2e^{-x} \right]$ $x = -1.5, \frac{d^2y}{dx^2} = -2e^{1.5} [= -8.96...] < 0$ <p>Hence [local] maximum</p>	B1 E1	or considers first derivative either side of $x = -1.5$, e.g. $x < -1.5 \Rightarrow \frac{dy}{dx} > 0$ $x > -1.5 \Rightarrow \frac{dy}{dx} < 0$ Must have been awarded B1
		2	

11(c)	$\int (5+2x)e^{-x} dx = -(5+2x)e^{-x} + \int 2e^{-x} dx$ $= -5e^{-x} - 2xe^{-x} - 2e^{-x}$ $= -7e^{-x} - 2xe^{-x}$ $\int_{-2.5}^0 (5+2x)e^{-x} dx = [-7e^{-x} - 2xe^{-x}]_{-2.5}^0$ $= (-7) - (-7e^{2.5} + 5e^{2.5})$ $= -7 + 2e^{2.5}$ <p>[Area of Triangle =] $0.5 \times 5 \times 2.5 \left[= \frac{25}{4} \right]$</p> <p>[Area =] $-7 + 2e^{2.5} - 0.5 \times 5 \times 2.5$</p> <p>[Area =] $-13.25 + 2e^{2.5}$</p>	M1 oe Use of integration by parts Condone omission of dx throughout m1 Complete use of integration by parts A1 M1 Their integral of the form $ae^{-x} + bxe^{-x}$ evaluated between their -2.5 and 0 B1 May be seen at any point in their solution A1 oe such as $2e^{2.5} - \frac{53}{4}$	
		6	
	Total	13	

Q	Answer	Marks	Comments
12	$y = x \ln(x + y)$ $\frac{dy}{dx} = \ln(x + y) + \frac{x}{x + y} \left(1 + \frac{dy}{dx} \right)$ $(x + y) \frac{dy}{dx} = (x + y) \ln(x + y) + x \left(1 + \frac{dy}{dx} \right)$ $x \frac{dy}{dx} + y \frac{dy}{dx} = (x + y) \frac{y}{x} + x + x \frac{dy}{dx}$ $y \frac{dy}{dx} = (x + y) \frac{y}{x} + x$ $\frac{dy}{dx} = (x + y) \frac{1}{x} + \frac{x}{y}$ $\frac{dy}{dx} = 1 + \frac{y}{x} + \frac{x}{y}$	<p>M1 A1</p> <p>m1</p> <p>m1</p> <p>A1</p> <p>A1</p>	<p>M1: Attempt at implicit differentiation A1: All correct</p> <p>Eliminates the fraction</p> <p>Expands & eliminates logarithm or correctly isolates $\frac{dy}{dx}$ term</p> <p>Expands & eliminates logarithm and correctly isolates $\frac{dy}{dx}$ term</p> <p>AG Be convinced</p>
	Total	6	

12 ALT 1	$\left[\frac{y}{x} = \ln(x+y) \Rightarrow \right] e^{\frac{y}{x}} = x+y$ $\frac{d}{dx} \left(e^{\frac{y}{x}} \right) = \frac{d}{dx} (x+y)$ $e^{\frac{y}{x}} \left(x \frac{dy}{dx} - y \right) = 1 + \frac{dy}{dx}$ $(x+y) \left(x \frac{dy}{dx} - y \right) = x^2 \left(1 + \frac{dy}{dx} \right)$ $x^2 \frac{dy}{dx} - xy + yx \frac{dy}{dx} - y^2 = x^2 + x^2 \frac{dy}{dx}$ $yx \frac{dy}{dx} = y^2 + x^2 + xy$ $\frac{dy}{dx} = \frac{y^2 + x^2 + xy}{yx}$ $\frac{dy}{dx} = \frac{y}{x} + \frac{x}{y} + 1$	M1 m1 A1 m1 A1 A1	PI m1 : Attempt at implicit differentiation A1 : All correct Eliminates exponential term or correctly isolates $\frac{dy}{dx}$ term Eliminates exponential term and correctly isolates $\frac{dy}{dx}$ term AG Be convinced
	Total	6	
12 ALT 2	$\frac{y}{x} = \ln(x+y)$ $\frac{d}{dx} \left(\frac{y}{x} \right) = \frac{d}{dx} (\ln(x+y))$ $\frac{x \frac{dy}{dx} - y}{x^2} = \frac{1}{x+y} \left(1 + \frac{dy}{dx} \right)$ $(x+y) \left(x \frac{dy}{dx} - y \right) = x^2 \left(1 + \frac{dy}{dx} \right)$ $\frac{dy}{dx} (x(x+y) - x^2) = xy + y^2 + x^2$ $\frac{dy}{dx} = \frac{xy + y^2 + x^2}{xy}$ $\frac{dy}{dx} = 1 + \frac{y}{x} + \frac{x}{y}$	M1 m1 A1 m1 A1 A1	PI m1 : Attempt at implicit differentiation A1 : All correct Eliminates fractions or correctly isolates $\frac{dy}{dx}$ term Eliminates fractions and correctly isolates $\frac{dy}{dx}$ term AG Be convinced
	Total	6	

Q	Answer	Marks	Comments
13(a)	$[AB =]\sqrt{(16-2)^2 + (-1-(-3))^2 + (-1-7)^2}$ $[AB =]\sqrt{14^2 + 2^2 + (-8)^2}$ $AB = \sqrt{264}$	M1 A1	oe oe such as $2\sqrt{66}$ Condone 16.2[48...]
		2	
13(b)(i)	$2 + 14\lambda = 9 + 5\mu$ $-3 + 2\lambda = -2 - 4\mu$ $\lambda = 0.5, \quad \mu = 0$ $7 - 8\lambda = q + 5\mu$ $q = 3$	M1 A1 A1	May use B instead Equating x and y
		3	
13(b)(ii)	$\begin{bmatrix} 14 \\ 2 \\ -8 \end{bmatrix} \cdot \begin{bmatrix} 5 \\ -4 \\ 5 \end{bmatrix} = 22$ $\cos \theta = \frac{\pm 22}{\sqrt{66}\sqrt{264}} \left[= \pm \frac{1}{6} \right]$ $[\theta =] 80.4^{[^\circ]}$	M1 A1 M1 A1	M1: Use of scalar product with direction vectors of l and AB A1: Correctly finds 22 ft their 22 from the scalar product between the two correct vectors
		4	

<p>13(c)</p> $\mathbf{C}(9+5c, -2-4c, 3+5c) \text{ or } \overrightarrow{OC} = \begin{bmatrix} 9+5c \\ -2-4c \\ 3+5c \end{bmatrix}$ $\overrightarrow{CD} = \begin{bmatrix} 10+5c \\ -4-4c \\ 5c \end{bmatrix}$ $\begin{bmatrix} 10+5c \\ -4-4c \\ 5c \end{bmatrix} \cdot \begin{bmatrix} 5 \\ -4 \\ 5 \end{bmatrix} = 0$ $50 + 25c + 16 + 16c + 25c = 0$ $\mathbf{C}(4, 2, -2) \text{ or } \overrightarrow{OC} = \begin{bmatrix} 4 \\ 2 \\ -2 \end{bmatrix}$ $BC^2 = (4-16)^2 + (2-(-1))^2 + (-2-(-1))^2 = \sqrt{154}$ $AC^2 = (4-2)^2 + (2-(-3))^2 + (-2-7)^2 = \sqrt{110}$ $AB^2 = AC^2 + BC^2 \text{ [so right-angled triangle]}$	<p>B1</p> <p>M1</p> <p>m1</p> <p>A1</p> <p>m1</p> <p>A1</p>	<p>oe</p> <p>$66c = -66, \quad c = -1$</p> <p>or finding $\overrightarrow{BC} \cdot \overrightarrow{AC} = -24 + 15 + 9$</p> <p>Note $\overrightarrow{AC} = \begin{bmatrix} 2 \\ 5 \\ -9 \end{bmatrix}$ and $\overrightarrow{BC} = \begin{bmatrix} -12 \\ 3 \\ -1 \end{bmatrix}$</p> <p>$\overrightarrow{BC} \cdot \overrightarrow{AC} = 0$ [so right-angled triangle]</p>
	<p>6</p>	
	<p>Total</p>	<p>15</p>