

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

I declare this is my own work.

INTERNATIONAL A-LEVEL MATHEMATICS

(9660/MA05) Unit M2 Mechanics

Tuesday 26 January 2021 07:00 GMT Time allowed: 1 hour 30 minutes

Materials

- For this paper you must have the Oxford International AQA Booklet of Formulae and Statistical Tables (enclosed).
- You may use a graphical calculator.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- The **final** answer to questions requiring the use of calculators should be given to two significant figures, unless stated otherwise.
- Unless stated otherwise, the acceleration due to gravity, g , should be taken as 9.8 m s^{-2}

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.


For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
TOTAL	



J A N 2 1 M A 0 5 0 1

Answer **all** questions in the spaces provided.

- 1** The diagram below shows a stationary motorcycle of mass 170 kg in equilibrium on horizontal ground.



The wheels are in contact with the ground at the points A and B where $AB = 1.40$ metres

The motorcycle's centre of mass is vertically above the point C where $BC = 0.80$ metres

- 1 (a)** Find the magnitude of the reaction force which acts on the motorcycle's rear wheel at A due to its contact with the ground.

[3 marks]

Answer _____



- 1 (b)** Find the magnitude of the reaction force which acts on the motorcycle's front wheel at B due to its contact with the ground.

[2 marks]

Answer

5

Turn over for the next question

Turn over ►



- 2** A body of mass 10 kg moves so that its velocity $\mathbf{v} \text{ m s}^{-1}$ at time t seconds is given by

$$\mathbf{v} = 2\sin 2t \mathbf{i} + e^{-t} \mathbf{j} + (3t^2 - \cos t) \mathbf{k}$$

where \mathbf{i} , \mathbf{j} and \mathbf{k} are unit vectors.

- 2 (a) (i)** Find the acceleration of the body when $t = \frac{\pi}{3}$

[3 marks]

Answer _____

- 2 (a) (ii)** Find the magnitude of the resultant force acting on the body when $t = \frac{\pi}{3}$

[2 marks]

Answer _____



Find the position of the body at time t

[4 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


Answer

C



- 3** A child of mass 30 kg moves down a slide.

The diagram below shows the shape of the slide.



The child starts at position *A* which is at a height of 3.0 metres above the level of position *B*

The speed of the child at *A* is 1.2 m s^{-1}

The speed of the child at *B* is 4.0 m s^{-1}

The curved length of the slide between *A* and *B* is 12 metres

- 3 (a)** Show that the child loses 882 J of gravitational potential energy moving from *A* to *B* **[1 mark]**

- 3 (b)** It is assumed that the child experiences a resistive force of constant magnitude R newtons at all times whilst moving from *A* to *B*

Find the value of R

[4 marks]



Answer _____

- 3 (c)** A student wants to determine the maximum speed the child could have at *B* if no resistive forces act on the child.

The student uses the following method:

$$v^2 = u^2 + 2as$$

$$v^2 = 1.2^2 + 2 \times 9.8 \times 12$$

$$v^2 = 236.64$$

$$v = 15 \text{ m s}^{-1} \text{ (to 2 significant figures)}$$

Give **two reasons** why the student's method for determining the maximum speed of the child is incorrect.


[2 marks]

Reason 1 _____

Reason 2 _____



- 4** A design is made from five rectangular uniform laminas, as shown below.
The laminas do not overlap and are made from the same material.



- 4 (a) (i)** Find the distance of the centre of mass of the design from OA, giving your answer in exact form.

[4 marks]

Answer _____



- 4 (a) (ii)** Find the distance of the centre of mass of the design from OB , giving your answer in exact form.

[3 marks]

Answer _____

- 4 (a) (iii)** Explain how you have used the fact that each rectangular lamina is uniform.

[1 mark]

- 4 (b)** The design is suspended from the point A and is in equilibrium.

Find, to the nearest degree, the angle between OA and the vertical.

[3 marks]

Answer _____

Turn over ►



5 A particle of mass 5 kg is acted upon by a constant driving force **F** newtons and accelerates at $\begin{bmatrix} 2 \\ 3 \end{bmatrix} \text{ m s}^{-2}$

The initial velocity of the particle is $\begin{bmatrix} 0 \\ 4 \end{bmatrix} \text{ m s}^{-1}$

At time t seconds after the force **F** newtons begins to act the velocity of the particle is **v** m s⁻¹

5 (a) Find **F**

[1 mark]

Answer _____

5 (b) (i) Find **v** in terms of t

[2 marks]

Answer _____



5 (b) (ii) Find the kinetic energy of the particle when $t = 6$

[3 marks]

Answer _____

5 (c) The work done on the particle each second is P joules, where P is the scalar product of the driving force \mathbf{F} newtons and the velocity \mathbf{v} m s⁻¹

Find the range of values of t for which P exceeds 580

[3 marks]

Answer _____



6


A particle of mass 2.4 kg is attached to two light inextensible strings.

The first string is attached to a fixed point B and has length 1.5 metres

The second string is attached to a fixed point C and has length 0.9 metres

The point C is 1.2 metres vertically below B

The particle is set into motion from the point A , as shown in the diagram below, and moves around a horizontal circle with centre C at a constant speed of 6 m s^{-1} so that both strings are taut, as shown in the diagram.



6 (a) (i) Both strings are described as inextensible.

Explain what is meant by inextensible.

[1 mark]

6 (a) (ii) State, with a reason, whether or not the particle is accelerating.

[2 marks]



6 (b) (i) Find the tension in the string attached to B

[3 marks]

Answer _____

6 (b) (ii) Find the tension in the string attached to C , giving your answer to three significant figures.

[4 marks]

Answer _____

10

Turn over ►



- 7** A sphere of weight 500 N is initially at rest in a liquid.
- A constant force of magnitude 2000 N acts vertically upwards on the sphere.
- The sphere begins moving vertically upwards through the liquid.
- The sphere also experiences a resistive force. When the speed of the sphere is $v \text{ m s}^{-1}$ the resistive force is kv^2 newtons vertically downwards, where k is a constant.

- 7 (a)** Show that the magnitude of the acceleration of the sphere is 29.4 m s^{-2} when the sphere is initially at rest.

[2 marks]

- 7 (b)** The acceleration of the sphere is 9.8 m s^{-2} upwards when the sphere is moving upwards at 5 m s^{-1}

Find the value of k

[3 marks]

Answer _____



[2 marks]

[illegible]

Answer _____

7

Turn over for the next question

Turn over ►



- 8** An object is projected with speed $u \text{ m s}^{-1}$ at an angle θ degrees above the horizontal from a point O on horizontal ground, where $0 < \theta < 90$

- 8 (a) (i)** Show that the time taken for the object to reach its maximum height above the ground is

$$\frac{u \sin \theta}{g}$$

[2 marks]

- 8 (a) (ii)** State an assumption you have made in **part (a)(i)**.

[1 mark]




[6 marks]

[illegible]

Answer



- 9** A block of mass 12 kg starts from rest on a rough slope which is inclined at θ to the horizontal, as shown in the diagram.



The coefficient of friction between the block and the slope is 0.4

- 9 (a)** Draw a diagram to show all the forces acting on the block, writing down the names of the forces on your diagram.

[1 mark]

- 9 (b)** The block accelerates uniformly down the slope at 3.2 m s^{-2}

Find the value of θ using $A \sin \theta - B \cos \theta = R \sin(\theta - \alpha)$

Give your answer to the nearest degree.

[8 marks]



9 (c) Without further calculation, state, with a reason, how your answer to **part (b)** may change if

13



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



[illegible]

There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.oxfordaqaexams.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2021 Oxford International AQA Examinations and its licensors. All rights reserved.



2 4



2 1 1 X M A 0 5

IB/G/Jan21/MA05