

# INTERNATIONAL A-LEVEL FURTHER MATHEMATICS FM03

---

Mark scheme

June 2019

---

Version: 1.0 Final

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [aqa.org.uk](http://aqa.org.uk)

### Key to mark scheme abbreviations

<b>M</b>	Mark is for method
<b>m</b>	Mark is dependent on one or more M marks and is for method
<b>A</b>	Mark is dependent on M or m marks and is for accuracy
<b>B</b>	Mark is independent of M or m marks and is for method and accuracy
<b>E</b>	Mark is for explanation
<b>✓ or ft</b>	Follow through from previous incorrect result
<b>CAO</b>	Correct answer only
<b>CSO</b>	Correct solution only
<b>AWFW</b>	Anything which falls within
<b>AWRT</b>	Anything which rounds to
<b>ACF</b>	Any correct form
<b>AG</b>	Answer given
<b>SC</b>	Special case
<b>oe</b>	Or equivalent
<b>A2, 1</b>	2 or 1 (or 0) accuracy marks
<b>–x EE</b>	Deduct x marks for each error
<b>NMS</b>	No method shown
<b>PI</b>	Possibly implied
<b>SCA</b>	Substantially correct approach
<b>sf</b>	Significant figure(s)
<b>dp</b>	Decimal place(s)

Q	Answer	Marks	Comments
1(a)	$\left(\frac{1}{2r+1} - \frac{1}{2r+3}\right) = \frac{2r+3 - (2r+1)}{(2r+1)(2r+3)}$ $= \frac{2}{(2r+1)(2r+3)}$	M1  A1	Condone omission of brackets for the M1 mark  CSO.
1(b)	Attempt to use method of differences  $(A) \left\{ \frac{1}{3} - \frac{1}{2n+3} \right\}$ $(A) \left\{ \frac{2n+3-3}{3(2n+3)} \right\}$ $\sum_{r=1}^n \frac{1}{(2r+1)(2r+3)} = \frac{1}{2} \left\{ \frac{2n}{3(2n+3)} \right\}$ $= \frac{n}{3(2n+3)}$	M1  A1  A1  A1	Must include four terms including two which cancel eg $\left(\frac{1}{3} - \frac{1}{5}\right) + \left(\frac{1}{5} - \frac{1}{7}\right)$ with or without factor of $\frac{1}{c's k}$  $\frac{1}{3} - \frac{1}{2n+3}$ after cancellations. Ignore any non-zero multiplier  Writing $\frac{1}{3} - \frac{1}{2n+3}$ with a common denominator  CSO.
	<b>Total</b>	<b>6</b>	

Q	Answer	Marks	Comments
2(a)	$2 \cosh^2 x - 1 = 2 \left( \frac{e^x + e^{-x}}{2} \right)^2 - 1$ $= 2 \left( \frac{e^{2x} + 2 + e^{-2x}}{4} \right) - 1$ $= \left( \frac{e^{2x} + 2 + e^{-2x}}{2} \right) - 1 = \frac{e^{2x} + e^{-2x}}{2}$ $= \cosh 2x$	M1	Correct exponential form for $\cosh x$ and attempt to expand
		A1	Correct expansion
		A1	CSO, AG
2(b)	$\frac{dy}{dx} = 6 \cosh 2x - 5 \cosh x + 4$ $0 = 6(2 \cosh^2 x - 1) - 5 \cosh x + 4$ $0 = 12 \cosh^2 x - 5 \cosh x - 2$ $0 = (4 \cosh x + 1)(3 \cosh x - 2)$ $\cosh x = -\frac{1}{4}, \quad \cosh x = \frac{2}{3}$ <p>But <math>\cosh x \geq 1</math>, so no (real) solutions</p> <p>(Since) <math>\frac{dy}{dx} \neq 0</math>, (the curve has) no stationary points</p>	M1	Differentiates, at least two of three terms correct
		m1	Puts $\frac{dy}{dx} = 0$ and forms a quadratic in $\cosh x$
		A1	Correct quadratic equation in a suitable form for solving
		A1	Both values for $\cosh x$ oe
		E1	Valid reason(s) for discounting <b>both</b> the candidate's two roots.
		A1	CSO Previous 5 marks scored and conclusion stated. If ' $\frac{dy}{dx} \neq 0$ ' is missing here, accept statement ' $\frac{dy}{dx} = 0$ for stationary points' oe at any stage
Total		9	

Q	Answer	Marks	Comments
3(a)(i)	$\alpha\beta + \beta\gamma + \gamma\alpha = 3$	B1	Condone 9/3
3(a)(ii)	$\alpha^2 + \beta^2 + \gamma^2$ $= (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha)$ $\alpha + \beta + \gamma = -\frac{b}{a} = 0$ $\alpha^2 + \beta^2 + \gamma^2 = 0 - 2(3) = -6$	M1  B1 A1	Correct formula  Seen or used CSO, AG.
3(a)(iii)	$\alpha^2 + \beta^2 + \gamma^2 < 0$ so roots not all real. Coefficients (of cubic eqn) are all real so non-real roots occur in conjugate pairs ie 2 non-real and 1 real	E1  E1	oe  oe  0/2 if 'Hence' not used.
3(b)(i)	(Complex conjugate) $1 - \sqrt{6}i$ is a root  $3^{\text{rd}}$ root: $0 - (1 - \sqrt{6}i) - (1 + \sqrt{6}i) = -2$ $\alpha\beta\gamma = -2(1 + 6) = -14$	M1  A1 A1	Or equating real and imaginary parts after substituting $1 + \sqrt{6}i$ for $z$ in the given cubic equation.
3(b)(ii)	$r = -3 \times \alpha\beta\gamma = 42$	B1ft	Ft on $-3 \times$ candidate's (b)(i) answer
Total		10	

Q	Answer	Marks	Comments
4(a)	$\begin{vmatrix} 1 & 3 & c \\ 1 & 2 & 3 \\ 1 & 1 & 1 \end{vmatrix} = 0$ $1(2-3) - 3(1-3) + c(1-2) = 0$ $\begin{vmatrix} 1 & 3 & c \\ 1 & 2 & 3 \\ 1 & 1 & 1 \end{vmatrix} = 5 - c = 0; \quad c = 5$	<p>M1</p> <p>A1</p>	<p>Equates <math>\begin{vmatrix} 1 &amp; 3 &amp; c \\ 1 &amp; 2 &amp; 3 \\ 1 &amp; 1 &amp; 1 \end{vmatrix}</math> to 0 and attempts to solve for <math>c</math>.</p> <p>CSO, AG be convinced</p>
4(b)	$x + 3y + 5z = 9$ (1); $x + 2y + 3z = 6$ (2) $x + y + z = d$ (3) Eg (1) – (2): $y + 2z = 3$  $(2) - (3): y + 2z = 6 - d$  $6 - d = 3$  $d = 3$	<p>M1</p> <p>A1</p> <p>A1</p> <p>m1</p> <p>A1</p>	<p>Eliminating a variable from two eqns, no more than one indep. error</p> <p>Using a different pair of eqns to get another value for <math>k</math> (previous <math>y + 2z</math>)</p> <p>Forming equation in <math>d</math> only, for consistent equations</p> <p>[If <math>y</math>-eliminated: <math>4d - 12 = 3d - 9</math> oe ]</p> <p>[If <math>z</math>-eliminated: <math>6d - 12 = 5d - 9</math> oe ]</p> <p><math>d = 3</math></p>
	Award equivalent marks for other approaches to solving the equations.		
	<b>Total</b>	<b>7</b>	

Q	Answer	Marks	Comments
5(a)	$f(k+1) - 2f(k)$ $= 2^{k+3} + 3^{2k+3} - 2(2^{k+2} + 3^{2k+1})$ $= 2^{k+3} + 3^{2k+3} - 2^{k+3} - 2 \times 3^{2k+1}$ $= 9 \times 3^{2k+1} - 2 \times 3^{2k+1} = 7 \times 3^{2k+1}$	M1 A1 A1	Seen or used Correct expansion of brackets and $2 \times 2^{k+2} = 2^{k+3}$ used Convincingly shown
5(b)	<p>Let <math>f(n) = 2^{n+2} + 3^{2n+1}</math>                      Assume result true for <math>n = k</math>,                      ie assume <math>f(k)</math> is a multiple of 7 (*)                      ie <math>f(k) = 7 \times M</math>, <math>M</math> an integer                      From (a), <math>f(k+1) = 2f(k) + 7 \times 3^{2k+1}</math></p> <p><math>f(k+1) = 7 \times (2M + 3^{2k+1})</math>                      Now <math>2k+1</math> is a positive integer so  <math>f(k+1) = 7 \times (2M + N) = 7 \times W</math>,                      integers <math>N</math> and <math>W</math>  <math>\therefore</math> if <math>f(k)</math> is a multiple of 7 then  <math>f(k+1)</math> is a multiple of 7 (**)</p> <p><math>f(1) = 8 + 27 (= 35) = 7 \times 5 \Rightarrow f(1)</math> is a multiple of 7</p> <p>Since <math>f(1)</math> is a multiple of 7, <math>f(2)</math>, <math>f(3)</math>,  <math>\dots</math> are multiples of 7 by induction,  <math>2^{n+2} + 3^{2n+1}</math> is a multiple of 7 for all  <b>integers</b> <math>n \geq 1</math></p>	M1  A1  B1  E1	Attempt at $f(k+1) = \dots$ , ft c's integer $a$  Showing that if $f(k)$ is a multiple of 7 then $f(k+1)$ is a multiple of 7  Must explicitly show that $8 + 27$ is a multiple of 7  Precise conclusion also dep. on previous 3 marks scored and (*) and (**) present. E0 if statement is not precise, eg 'a multiple of 7 for all $n \geq 1$ '
	<b>Total</b>	<b>7</b>	



Q	Answer	Marks	Comments
6(a)	$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = 20e^{2x} + 18$ <p>PI: <math>y_{PI} = p + qxe^{2x}</math></p> $y'_{PI} = qe^{2x} + 2qxe^{2x}$ $y''_{PI} = 4qe^{2x} + 4qxe^{2x}$ $4qe^{2x} + 4qxe^{2x} + qe^{2x} + 2qxe^{2x} - 6p - 6qxe^{2x} = (20e^{2x} + 18)$ $-6p = 18 \text{ and } 5q = 20$ $p = -3; \quad q = 4$ $[y_{PI} = -3 + 4xe^{2x}]$	<p>M1</p> <p>M1</p> <p>m1</p> <p>A2,1,0</p>	<p><math>\pm ae^{2x} \pm bxe^{2x}</math></p> <p>Substitution into LHS of DE attempted</p> <p>Dep on 2<sup>nd</sup> M1 only... equating coefficients to form two equations at least one correct. PI by correct values for both <math>p</math> and <math>q</math></p> <p>A1 if both values correct but <math>xe^{2x}</math> terms in 2<sup>nd</sup> M1 line do not cancel</p>
6(b)	<p>Aux. eqn. <math>m^2 + m - 6 = 0</math>  <math>(m + 3)(m - 2) = 0</math></p> <p><math>(y_{CF}) = Ae^{-3x} + Be^{2x}</math></p> <p><math>(y_{GS}) = Ae^{-3x} + Be^{2x} - 3 + 4xe^{2x}</math></p> <p><math>x = 0, y = 5 \Rightarrow 5 = A + B - 3</math></p> <p><math>\frac{dy}{dx} = -3Ae^{-3x} + 2Be^{2x} + 4e^{2x} + 8xe^{2x}</math></p> <p>As <math>x \rightarrow -\infty, (e^{2x} \rightarrow 0)</math> and <math>xe^{2x} \rightarrow 0</math></p> <p>As <math>x \rightarrow -\infty, \frac{dy}{dx} \rightarrow 0</math> so <math>A = 0</math></p> <p>When <math>A = 0, 5 = 0 + B - 3 \Rightarrow B = 8</math></p> <p><math>y = 8e^{2x} - 3 + 4xe^{2x}</math></p>	<p>M1</p> <p>A1</p> <p>B1ft</p> <p>B1ft</p> <p>E1</p> <p>B1</p> <p>A1</p>	<p>Factorising or using quadratic formula oe on correct aux eqn. PI by correct two values of 'm' seen/used Correct CF</p> <p>c's CF + c's PI but must have exactly two arbitrary constants</p> <p>Only ft if previous B1ft scored and GS contains exponentials</p> <p>Must treat <math>xe^{2x} \rightarrow 0</math> separately</p> <p>Coefficient of <math>e^{-3x}</math> is 0</p> <p><math>y = 8e^{2x} - 3 + 4xe^{2x}</math></p>
Total		12	

Q	Answer	Marks	Comments
7(a)	$\begin{bmatrix} 1 & -1 & 2 \\ 0 & 5 & 7 \\ k & 1 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 7 \\ 1 \end{bmatrix} = \lambda_1 \begin{bmatrix} -1 \\ 7 \\ 1 \end{bmatrix}$ $\begin{bmatrix} -6 \\ 42 \\ -k+8 \end{bmatrix} = \lambda_1 \begin{bmatrix} -1 \\ 7 \\ 1 \end{bmatrix}, \quad \lambda_1 = 6$ $-k+8 = \lambda_1$ $k = 2$	M1   B1  m1 A1	$\mathbf{M}\mathbf{v}_1 = \lambda_1 \mathbf{v}_1$ with $\mathbf{M}\mathbf{v}_1$ attempted oe  $\lambda_1 = 6$  Equating c's components to form an equation involving $k$ $k = 2$
7(b)	$\det(\mathbf{M} - \lambda \mathbf{I}) = \begin{vmatrix} 1-\lambda & -1 & 2 \\ 0 & 5-\lambda & 7 \\ k & 1 & 1-\lambda \end{vmatrix}$ $= (1-\lambda) \begin{vmatrix} 5-\lambda & 7 \\ 1 & 1-\lambda \end{vmatrix} + k \begin{vmatrix} -1 & 2 \\ 5-\lambda & 7 \end{vmatrix}$ $= (1-\lambda)[(5-\lambda)(1-\lambda) - 7] + k[-17 + 2\lambda]$ <p>Characteristic eqn:  <math>-\lambda^3 + 7\lambda^2 - 36 = 0</math></p> <p>Eigenvalues are 3, (6) and <math>-2</math> so <math>-2</math> is the least eigenvalue</p>	M1  A1ft  A1  A1	oe  Correct (unsimplified) expression for expansion of $\det(\mathbf{M} - \lambda \mathbf{I})$ ; if value for $k$ used, ft on c's non-zero value ACF of the characteristic eqn correctly found eg $(\lambda - 3)(6 - \lambda)(2 + \lambda) = 0$ Eigenvalues 3 and $-2$ and final conclusion; must see a correct equation
7(c)	$\begin{bmatrix} 1 & -1 & 2 \\ 0 & 5 & 7 \\ k & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = -2 \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ $3x - y + 2z = 0; \quad \text{so } 3x + 3z = 0$ $7y + 7z = 0 \Rightarrow y = -z$ $kx + y + 3z = 0 \quad \text{so } 2x + 2z = 0$ $x = y = -z \text{ so an eigenvector is}$ $\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$	M1  A1ft  A1	$\mathbf{M}\mathbf{v} = -2\mathbf{v}$ oe and attempt to get a system of equations  Three correct ft equations with a later substitution resulting in two equations in just two variables  An eigenvector in form $\beta \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$ ,  $\beta \neq 0$
7(d)	$-x = \frac{y}{7} = z; \text{ or } x = y = -z$	B1	Either one oe. [since Q is 'write down', the case $\lambda = 3$ is not relevant]
Total		12	

Q	Answer	Mark	Comments
8(a)	$\frac{dy}{dx} + \frac{1}{x(x+1)}y = 2x+3$ $\text{I.F. is } \exp\left(\int \frac{1}{x(x+1)}(dx)\right)$ $\frac{1}{x(x+1)} = \frac{A}{x} + \frac{B}{x+1}$ $\exp\left(\int\left(\frac{1}{x} - \frac{1}{x+1}\right)(dx)\right)$ $(\text{I.F.}) = e^{\ln x - \ln(x+1)}$ $= \frac{x}{x+1}$	<p>M1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>Identified and integration attempted</p> <p>Partial fractions used as far as finding a value for <math>A</math> and a value for <math>B</math>. PI by next line</p> <p>AG be convinced</p>
8(b)	$\left(\frac{x}{x+1}\right)\frac{dy}{dx} + \frac{1}{(x+1)^2}y = \frac{x(2x+3)}{x+1}$ $\frac{d}{dx}\left[y \frac{x}{x+1}\right] = \frac{x(2x+3)}{x+1}$ $\frac{yx}{x+1} = \int \frac{x(2x+3)}{x+1}(dx)$ $\frac{yx}{x+1} = \int \frac{2x^2+3x}{x+1}(dx)$ $= \int 2x+1 - \frac{1}{x+1}(dx)$ $= x^2 + x - \ln(x+1) (+A)$ $\frac{yx}{x+1} = x^2 + x - \ln(x+1) + A$ $y = \left(\frac{x+1}{x}\right)[x^2 + x - \ln(x+1) + A]$	<p>M1</p> <p>A1</p> <p>m1</p> <p>A1</p> <p>A1</p>	<p>Multiplies both sides of the DE by <math>\frac{x}{x+1}</math> and then identifies the LHS as the derivative of <math>y \times \text{I.F.}</math> PI by next line.</p> <p>oe</p> <p>Division to reach eg <math>2x + p + \frac{q}{x+1}</math>, where <math>p</math> and <math>q</math> are non-zero integers. PI by next line</p> <p>Correct integration of <math>\frac{x(2x+3)}{x+1}</math>, condone absence of '+constant' here</p> <p>ACF of the GS</p>
Total		9	

Q	Answer	Mark	Comments
9(a)	$\mathbf{r} \cdot \begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix} = 3 \quad \mathbf{r} \cdot \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} = 4$ $\begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} = 4$ $\cos \theta = \frac{4}{\sqrt{9}\sqrt{9}} = \frac{4}{9}$ $\text{Acute angle} = \cos^{-1}\left(\frac{4}{9}\right) = 63.6^\circ$	<p>M1</p> <p>A1</p> <p>B1</p> <p>A1</p>	<p>Use of the scalar product on the two normal vectors <math>\begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix}</math> and <math>\begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}</math></p> <p>Correct evaluation of scalar product</p> <p>[Equivalent marks for the use of the modulus of the relevant vector product]</p> <p>Denominator = 9; correct product of moduli</p> <p>CAO must be 63.6</p>
9(b)	<p>eg <math>(0, a, b)</math>, <math>-a + 2b = 3</math>, <math>2a + b = 4</math> Solving gives a common pt <math>(0, 1, 2)</math></p> $\begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix} \times \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -5 \\ 2 \\ 6 \end{bmatrix}$ $\mathbf{r} = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} + t \begin{bmatrix} -5 \\ 2 \\ 6 \end{bmatrix}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>Method to find a common point: Any correct common pt. [other likely ones are <math>(2.5, 0, -1)</math>; <math>(5/3, 1/3, 0)</math>]</p> <p>Finding direction vector of the line: eg <math>\mathbf{n}_1 \times \mathbf{n}_2</math> or <math>\mathbf{n}_2 \times \mathbf{n}_1</math> attempted or by applying a correct method to obtain and use two common points</p> <p>A correct direction vector</p> <p>oe ACF with correct notation eg <math>(\mathbf{r} - (\mathbf{j} + 2\mathbf{k})) \times (-5\mathbf{i} + 2\mathbf{j} + 6\mathbf{k}) = \mathbf{0}</math></p>
Total		9	

Q	Answer	Mark	Comments
10(a)	$\frac{dx}{dt} = 1 - \cos t;$	B1	oe Correct expression for $\frac{dx}{dt}$
	$\frac{dy}{dt} = 2 \left( 2 \sin \frac{t}{2} \right) \left( \frac{1}{2} \cos \frac{t}{2} \right)$	B1	oe Correct expression for $\frac{dy}{dt}$
	$\frac{dx}{dt} = 1 - \cos t = 2 \sin^2 \frac{t}{2}$	M1	$1 - \cos t = 2 \sin^2 \frac{t}{2}$ used at any stage or better
	$\dot{x}^2 + \dot{y}^2 = 4 \sin^4 \frac{t}{2} + 4 \sin^2 \frac{t}{2} \cos^2 \frac{t}{2} =$ $4 \sin^2 \frac{t}{2} (\sin^2 \frac{t}{2} + \cos^2 \frac{t}{2}) = 4 \sin^2 \frac{t}{2}$ $\left( \frac{dx}{dt} \right)^2 + \left( \frac{dy}{dt} \right)^2 = 4 \sin^2 \frac{t}{2}$	A1	CSO, AG
(b)	$(SA=) 2\pi \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} 2 \sin^2 \frac{t}{2} \left( 2 \sin \frac{t}{2} \right) dt$	B1	Must include the $2\pi$ and $dt$ with a correct integrand. Correct limits seen here or used correctly later.
	$= 8\pi \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} (1 - \cos^2 \frac{t}{2}) \left( \sin \frac{t}{2} \right) dt$		
	Subst: Let $u = \cos \frac{t}{2}; \frac{du}{dt} = -\frac{1}{2} \sin \frac{t}{2}$	M1	Valid method to integrate $\sin^3 \frac{t}{2}$ eg substitution/inspection [PI by answer
	$= (8\pi) \int_{\frac{\sqrt{3}}{2}}^{\frac{1}{2}} (1 - u^2)(-2) du$	A1	$\left( a \cos \frac{t}{2} + \frac{b}{3} \cos^3 \frac{t}{2} \right)$ ] and by parts Subst....correct integrand and limits By parts....Must reach a stage where only integrals remaining are multiples of $\sin^3 \frac{t}{2}$ or better . By inspection...as M1 above with $b=2$ or better.
	$= (8\pi) \left[ (-2) \left( u - \frac{u^3}{3} \right) \right]_{\left( \frac{\sqrt{3}}{2} \right)}^{\left( \frac{1}{2} \right)}$	A1	Integration of $\sin^3 \frac{t}{2}$ complete and correct. By parts...the most likely form for integral of $\sin^3 \frac{t}{2}$ will be
	$= -16\pi \left[ \left( \frac{1}{\sqrt{2}} - \frac{1}{6\sqrt{2}} \right) - \left( \frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{8} \right) \right]$	m1	$-\frac{2}{3} \sin^2 \frac{t}{2} \cos \frac{t}{2} - \frac{4}{3} \cos \frac{t}{2}$ oe
	$= \frac{2\pi}{3} (9\sqrt{3} - 10\sqrt{2})$	A1	Correct use of correct limits. Award even if limits reversed on integral sign. Be convinced as the form of the answer is given
Total		10	



Q	Answer	Marks	Comments
<b>11(a)(i)</b>	$z^n = \cos n\theta + i \sin n\theta$ $z^{-n} = \cos(-n\theta) + i \sin(-n\theta)$ $= \cos n\theta - i \sin n\theta$ $z^n + \frac{1}{z^n} = 2 \cos n\theta$	M1 E1 A1	Must be shown either as in soln or using $\frac{1}{\cos n\theta + i \sin n\theta} \times \frac{\cos n\theta - i \sin n\theta}{\cos n\theta - i \sin n\theta} =$ AG Note: M1E0 A1 is possible eg for those who just quote $z^{-n} = \cos n\theta - i \sin n\theta$ .
<b>11(a)(ii)</b>	$z - \frac{1}{z} = 2i \sin \theta$	B1	
<b>11(b)</b>	$(z - z^{-1})^6 = z^6 - 6z^4 + 15z^2 - 20 + 15z^{-2} - 6z^{-4} + z^{-6}$ $= z^6 + z^{-6} - 6(z^4 + z^{-4}) + 15(z^2 + z^{-2}) - 20$ $((2i \sin \theta)^6 =)$ $= 2 \cos 6\theta - 12 \cos 4\theta + 30 \cos 2\theta - 20$ $(2i \sin \theta)^6 = 64i^6 \sin^6 \theta = -64 \sin^6 \theta$ $64 \sin^6 \theta = 20 - 30 \cos 2\theta + 12 \cos 4\theta - 2 \cos 6\theta$	M1 A1 m1 A1 A1	Attempts to find expansion of $(z - z^{-1})^6$ Expansion correct Groups terms so as to use result in (a) PI Showing RHS CSO
<b>11(c)</b>	$(\text{Area of leaf}) = \frac{1}{2} \int_0^\pi 4 \sin^6 \theta \, d\theta =$ $\frac{1}{32} \int_0^\pi (20 - 30 \cos 2\theta + 12 \cos 4\theta - 2 \cos 6\theta) \, d\theta$ $(\int_0^\pi \cos n\theta \, d\theta = [\frac{1}{n} \sin n\theta]_0^\pi = 0)$ $(\text{Area leaf}) = \frac{1}{32} [20\theta]_0^\pi = \frac{20\pi}{32}$ $(= \frac{5\pi}{8})$ $(\text{Area of disc}) = \pi \times 1^2 = \pi$ $\% \text{ area of disc not covered} =$ $\frac{3}{8} \times 100 = 37.5\%$	M1 M1 A1ft B1 A1	Use of $\frac{1}{2} \int r^2 \, (d\theta)$ or $\int_0^\pi r^2 \, (d\theta)$ oe Uses <b>11(b)</b> with c's values for $a, b, c$ ; then integrates correctly or explains/clearly indicates that cos terms when integrated are 0 at each of the limits A correct area for the leaf; ft on candidate's non-zero values for $a, b, c$ . 37.5 OE with no errors seen
<b>Total</b>		<b>14</b>	

Q	Answer	Mark	Comments
12(a)	$\sin 2x = 2x - \frac{4}{3}x^3 + \frac{4}{15}x^5 \dots$	B1	Do not accept series in powers of $2x$
12(bi)	$\frac{1}{y} \frac{dy}{dx} = \frac{1}{1+x^2}$ $\frac{1}{y} \frac{d^2y}{dx^2} - \frac{1}{y^2} \left( \frac{dy}{dx} \right)^2 = -\frac{2x}{(1+x^2)^2}$ $y \frac{d^2y}{dx^2} - \left( \frac{dy}{dx} \right)^2 = \frac{-2x}{(1+x^2)^2} = -2x \left( \frac{dy}{dx} \right)^2$ $(2x-1) \left( \frac{dy}{dx} \right)^2 + y \frac{d^2y}{dx^2} = 0 \quad (*)$	B1;B1 M1 A1  A1	$\frac{1}{y} \frac{dy}{dx}$ oe (B1); $\frac{1}{1+x^2}$ (B1) Pr. rule/Quotient rule used appropriately Correct differentiation of prev line oe  CSO, AG
12(bii)	<p><math>\ln y = \tan^{-1} x \Rightarrow y = e^{\tan^{-1} x}</math>. From McL.                      when <math>x = 0</math>, <math>y' = 1</math>, <math>y'' = 1</math>, <math>y''' = 3!p</math>,  <math>y^{(iv)} = 4!q</math>                      Differentiating (*) wrt <math>x</math>:  <math>2(y')^2 + (2x-1)2y'y'' + y'y''' + yy'''' = 0</math>                      Sub <math>x = 0</math> gives  <math>2(1)^2 + (-1)2(1)(1) + (1)(1) + (1)y''''(0) = 0</math>  <math>\Rightarrow y''''(0) = -1 \Rightarrow p = \frac{y''''(0)}{3!} = -\frac{1}{6}</math>                      Differentiating wrt <math>x</math>:  <math>4y'y'' + 4y'y'' + (4x-2)[y'y''' + (y'')^2] +</math>  <math>+ y'y'''' + (y'')^2 + y'y'''' + yy^{(iv)} = 0</math>                      Sub <math>x = 0</math> gives  <math>4(1)(1) + 4(1)(1) - 2[(1)(-1) + 1^2] +</math>  <math>(1)(-1) + 1^2 + (1)(-1) + (1)y^{(iv)}(0) = 0</math>  <math>\Rightarrow y^{(iv)}(0) = -7</math>                      From McL series, <math>q = \frac{y^{(iv)}(0)}{4!} = -\frac{7}{24}</math></p>	B2,1,0  B1  B1  M1  A1	seen or used as appropriate. B1 if two are correct,  A correct equation involving $y'''$ oe  AG Be convinced  Product rule/quotient rule used appropriately to obtain an equation involving $y^{(iv)}$ oe  CSO



Q	Answer	Mark	Comments
	<p><u>Alternative for final 4 marks</u></p> <p>From (b)(i), <math>(1+x^2)\frac{d^2y}{dx^2} = (1-2x)\left(\frac{dy}{dx}\right)</math></p> <p><math>2x\frac{d^2y}{dx^2} + (1+x^2)\frac{d^3y}{dx^3} = -2\frac{dy}{dx} + (1-2x)\frac{d^2y}{dx^2}</math></p> <p>Sub <math>x=0</math>, <math>3!p = -2+1 \Rightarrow p = -\frac{1}{6}</math></p> <p><math>2\frac{d^2y}{dx^2} + 4x\frac{d^3y}{dx^3} + (1+x^2)\frac{d^4y}{dx^4}</math></p> <p><math>= -4\frac{d^2y}{dx^2} + (1-2x)\frac{d^3y}{dx^3}</math></p> <p>Sub <math>x=0</math>, <math>2+4!q = -4-1 \Rightarrow q = -\frac{7}{24}</math></p>	<p>(B1)</p> <p>(B1)</p> <p>(M1)</p> <p>(A1)</p>	<p>CSO</p>
12(c)	<p><math>\lim_{x \rightarrow 0} \left[ \frac{e^{\tan^{-1}x} - e^x}{2x - \sin 2x} \right] =</math></p> <p><math>= \lim_{x \rightarrow 0} \frac{-\frac{x^3}{6} + qx^4 - \frac{x^3}{6} - \frac{x^4}{24} \dots}{2x - 2x + \frac{4x^3}{3} + O(x^5)}</math></p> <p><math>= \lim_{x \rightarrow 0} \frac{-\frac{2}{6} + O(x)}{\frac{4}{3} + O(x^2)}, \quad (\text{so limit exists})</math></p> <p><math>= -0.25</math></p>	<p>M1</p> <p>m1</p> <p>A1</p>	<p>Substitution of series</p> <p>Dividing numerator and denominator by <math>x^3</math> to get <math>\lim_{x \rightarrow 0} \frac{a + O(x)}{b + O(x^2)}</math>, so limit exists</p> <p><math>= a/b</math>. In place of <math>O(\quad)</math> may have equivalent term(s)</p> <p>Correct value for the limit.</p> <p>(A0 if previous 2 marks not scored)</p>
<b>Total</b>		<b>15</b>	