

**OXFORD**

INTERNATIONAL  
AQA EXAMINATIONS

---

# INTERNATIONAL AS FURTHER MATHEMATICS FM02

(9665/FM02) Unit P2 – Unit FPSM1 – Pure, Statistics and Mechanics

---

Mark scheme

January 2020

---

Version: 1.0 Final

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [oxfordaqaexams.org.uk](http://oxfordaqaexams.org.uk)

#### **Copyright information**

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2020 Oxford International AQA Examinations and its licensors. All rights reserved.

**Key to mark scheme abbreviations**

<b>M</b>	Mark is for method
<b>m</b>	Mark is dependent on one or more M marks and is for method
<b>A</b>	Mark is dependent on M or m marks and is for accuracy
<b>B</b>	Mark is independent of M or m marks and is for method and accuracy
<b>E</b>	Mark is for explanation
✓ <b>or ft</b>	Follow through from previous incorrect result
<b>CAO</b>	Correct answer only
<b>CSO</b>	Correct solution only
<b>AWFW</b>	Anything which falls within
<b>AWRT</b>	Anything which rounds to
<b>ACF</b>	Any correct form
<b>AG</b>	Answer given
<b>SC</b>	Special case
<b>oe</b>	Or equivalent
<b>A2, 1</b>	2 or 1 (or 0) accuracy marks
<b>–x EE</b>	Deduct x marks for each error
<b>NMS</b>	No method shown
<b>PI</b>	Possibly implied
<b>SCA</b>	Substantially correct approach
<b>sf</b>	Significant figure(s)
<b>dp</b>	Decimal place(s)

Q	Answer	Marks	Comments
1	$hf(x, y) = 0.25 \frac{\sqrt{7} + 4.3}{4.3\sqrt{4.3}}$ $= 0.194741$ $y_2 = 4.3 + 0.194741 = 4.494741$ $y_3 = 4.494741 + 0.25 \frac{\sqrt{7.25} + 4.494741}{4.494741\sqrt{4.494741}}$ $= 4.68330$ 4.683	<b>M1</b>  <b>A1</b>  <b>m1</b>  <b>A1ft</b>  <b>A1</b>	PI   ft their $y_2$ to at least 3 dp  CAO
	<b>Total</b>	<b>5</b>	

Q	Answer	Marks	Comments
2(a)	$\begin{vmatrix} 1 & k \\ 3 & 2 \end{vmatrix} = 0$ $2 - 3k = 0$ $k = \frac{2}{3}$	<b>M1</b>  <b>A1</b>	  CAO
2(b)	$\mathbf{M} = \mathbf{B}^{-1}\mathbf{B}\mathbf{M} = \mathbf{B}^{-1} \begin{bmatrix} 1 & 2 & -1 \\ 0 & 1 & 4 \end{bmatrix}$ $\mathbf{B}^{-1} = \left( \frac{1}{2-3k} \right) \begin{bmatrix} 2 & -k \\ -3 & 1 \end{bmatrix}$ $\mathbf{M} = \left( \frac{1}{2-3k} \right) \begin{bmatrix} 2 & -k \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & -1 \\ 0 & 1 & 4 \end{bmatrix}$ $\mathbf{M} = \left( \frac{1}{2-3k} \right) \begin{bmatrix} 2 & 4-k & -2-4k \\ -3 & -5 & 7 \end{bmatrix}$	<b>B1</b>  <b>M1</b> <b>A1ft</b>  <b>M1</b>  <b>A1</b>	Alternative method: <b>B1</b> for 6 sim. equations (at least 5 correct) (PI) <b>M1</b> for attempting to solve a pair of their equations for their unknowns <b>A1</b> for correctly solving a pair of the equations. Then last two marks as for main method. <b>M1</b> for $\begin{bmatrix} 2 & -k \\ -3 & 1 \end{bmatrix}$ <b>A1ft</b> division by their $ \mathbf{B} $ For correctly calculating the 2 <sup>nd</sup> or 3 <sup>rd</sup> column of $\begin{bmatrix} 2 & 4-k & -2-4k \\ -3 & -5 & 7 \end{bmatrix}$  oe CAO

	<b>Total</b>	<b>7</b>	
--	--------------	----------	--

<b>Q</b>	<b>Answer</b>	<b>Marks</b>	<b>Comments</b>
----------	---------------	--------------	-----------------

<b>3(a)</b>	531 405 195 -74	<b>B1</b>	
<b>3(b)</b>	All points correct Line of best fit drawn	<b>B1</b> <b>B1</b>	
<b>3(c)</b>	$X$ -intercept = 91 $x = \sqrt{91}$ (= 9.5)	<b>M1</b> <b>A1</b>	In the range [86, 94] Square root of their $X$ -intercept in the range [9.3, 9.7]
<b>3(d)</b>	$b = 650$  $a = \text{gradient}$ $= -7.1$	<b>B1</b>  <b>M1</b> <b>A1</b>	their $Y$ -intercept in the range [630, 670] No marks if not consistent with graph PI In [-7.5, -6.7]
	<b>Total</b>	<b>8</b>	

<b>Q</b>	<b>Answer</b>	<b>Marks</b>	<b>Comments</b>
----------	---------------	--------------	-----------------

<b>4(a)</b>	$f(-1) = 1$ and $f(-1.5) = -19/8$ Change of sign and $f$ is continuous on the interval so $\alpha$ is in the interval $(-1.5, -1)$ .	<b>M1</b> <b>A1</b>	
<b>4(b)</b>	$f'(x) = 3x^2 - 4x - 3$ $x_2 = -1.5 - \frac{-19/8}{39/4}$ -1.256	<b>B1</b> <b>B1</b> <b>B1</b> <b>B1</b>	PI <b>B1</b> for numerator in correct form <b>B1</b> for denominator in correct form
<b>4(c)(i)</b>	Tangent drawn at $x = -1.5$ to meet $x$ -axis	<b>B1</b>	
<b>4(c)(ii)</b>	P correct Q correct	<b>B1</b> <b>B1</b>	
	<b>Total</b>	<b>9</b>	

Q	Answer	Marks	Comments
5(a)	(3,1), (11,3) and (8,2) Drawn correctly, labelled and joined up	B1 B1	
5(b)	$\begin{vmatrix} 3 & -4 \\ 1 & -1 \end{vmatrix} = 1$ Area of $OABC = 2$  Therefore, area of $OA'B'C'$ = $1 \times$ area of $OABC$ = $1 \times 2$ = 2 square units	M1  A1	With explanation  Alt. method: $\begin{vmatrix} 3 & 11 \\ 1 & 3 \end{vmatrix} = -2$ M1 so area = 2 A1 Or use of determinant made of any pair of position vectors of vertices
5(c)	$\begin{pmatrix} 3 & -4 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ mx \end{pmatrix} = \begin{pmatrix} x \\ mx \end{pmatrix}$ $3 - 4m = 1$ so $m = \frac{1}{2}$ or $1 - m = m$ so $m = \frac{1}{2}$	M1  E1	Must have only one unknown
5(d)	$\begin{pmatrix} 3 & -4 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ kx \end{pmatrix} = \begin{pmatrix} X \\ 3X \end{pmatrix}$ $3x - 4kx = X$ $x - kx = 3X$ $x - kx = 3(3x - 4kx)$ $1 - k = 9 - 12k$ $k = \frac{8}{11}$	M1  M1  A1  M1  A1	Must have two different unknowns  For two equations in 2 unknowns  For eliminating 1 unknown
	Total	11	

Q	Answer	Marks	Comments
<b>6</b>	$G_X(t) = \sum_{k=0}^n \binom{n}{k} p^k (1-p)^{n-k} t^k$	<b>M1</b>	Applies $G_X(t)$ formula
	$= \sum_{k=0}^n \binom{n}{k} (pt)^k (1-p)^{n-k}$	<b>M1</b>	Simplifies expression
	$= (1-p + pt)^n \quad (\text{AG})$	<b>A1</b>	Complete proof with no errors seen
	<b>Total</b>	<b>3</b>	


Q	Answer	Marks	Comments
<b>7(a)</b>	$E(X^2) = \sum_{x=1}^n \frac{x^2}{n} = \frac{1^2 + 2^2 + \dots + n^2}{n}$	<b>M1</b>	Applies formula for $E(X^2)$
	$= \frac{\frac{1}{6} n(n+1)(2n+1)}{n}$	<b>A1</b>	Applies formula for $\sum n^2$
	$= \frac{(n+1)(2n+1)}{6}$		
	$\text{Var}(X) = \frac{(n+1)(2n+1)}{6} - \left(\frac{n+1}{2}\right)^2$	<b>M1</b>	Applies $\text{Var}(X) = E(X^2) - (E(X))^2$
	$= \frac{n^2 - 1}{12}$	<b>A1</b>	Complete proof with no errors seen Needs at least one intermediate line
<b>7(b)(i)</b>	$\frac{n^2 - 1}{12} = 33.25$	<b>M1</b>	Sets up equation using $\text{Var}(X) = 33.25$
	$n = 20$	<b>A1</b>	
<b>7(b)(ii)</b>	$P(D > 18) = 0.1$	<b>B1ft</b>	Accept 1/10 oe Follow through their (their $n - 18$ ) $\times 1/20$
<b>7(b)(iii)</b>	$(1 - 0.1)^4 \times 0.1$	<b>M1</b>	$(1 - \text{their } P(D > 18))^4 \times \text{their } P(D > 18)$
	0.06561	<b>A1</b>	CAO
	<b>Total</b>	<b>9</b>	

Q	Answer	Marks	Comments
8(a)	$\text{Var}(aX) = 0.25a^2$ $\text{Var}((1-a)Y) = 0.16(1-a)^2$ or $0.16 - 0.32a + 0.16a^2$ $2\rho\sqrt{\text{Var}(aX)\text{Var}((1-a)Y)} =$ $2 \times 0.25 \times 0.5a \times 0.4(1-a)$ or $0.1a - 0.1a^2$ $\text{Var}(aX + (1-a)Y)$ $= 0.25a^2 + 0.16 - 0.32a + 0.16a^2 + 0.1a -$ $0.1a^2$ $= 0.31a^2 - 0.22a + 0.16$	<b>B1</b>  <b>B1</b>  <b>M1</b>  <b>A1</b>	Uses $\text{Var}(aX) = a^2 \text{Var}(X)$ Uses $\text{Var}((1-a)Y) =$ $(1-a)^2 \text{Var}(Y)$ Finds $2\rho\sqrt{\text{Var}(aX)\text{Var}((1-a)Y)}$ Or $2a(1-a)\text{Cov}(X,Y)$ Applies $\text{Var}(aX + (1-a)Y)$ $= \text{Var}(aX) + \text{Var}((1-a)Y) +$ $2\rho\sqrt{\text{Var}(aX)\text{Var}((1-a)Y)}$ (or uses $2a(1-a)\text{Cov}(X,Y)$ here) and simplifies to given answer
8(b)(i)	$\frac{d(\text{Var})}{da} = 0.62a - 0.22$ $0.62a - 0.22 = 0$ $a = \frac{11}{31}$	<b>B1</b>  <b>M1</b>  <b>A1</b>	Correct differentiation Sets $\frac{d(\text{Var})}{da} = 0$ CAO 0.35483..... is <b>B1M1A0</b>
8(b)(ii)	$\frac{226}{31}$	<b>B1ft</b>	Follow through their $a \times 6 + (1 - \text{their } a) \times 8$ Accept AWRT 7.3
	<b>Total</b>	<b>8</b>	



Q	Answer	Marks	Comments
9	$[a] = [r][\omega]^2$ $LT^{-2} = L[\omega]^2$ $[\omega]^2 = T^{-2}$ $[\omega] = T^{-1}$	<b>M1</b> <b>A1ft</b>  <b>A1</b>	ft their dimensions of acceleration (at least two terms)
	<b>Total</b>	<b>3</b>	

Q	Answer	Marks	Comments
10(a)	$e = \frac{5}{8} = 0.625$	<b>B1</b>	
10(b)	$I = 0.2 \times 8 - 0.2 \times (-5)$ $= 2.6 \text{ Ns}$	<b>M1</b> <b>A1</b>	-2.6 is <b>M1A0</b>
10(c)	$2.6 = 0.25F$ $F = \frac{2.6}{0.25} = 10.4 \text{ N}$	<b>M1</b> <b>A1</b>	
	<b>Total</b>	<b>5</b>	

Q	Answer	Marks	Comments
11(a)	 $\frac{\sin \alpha}{4} = \frac{\sin 135^\circ}{9}$ $\sin \alpha = 0.31427$ $\alpha = 018$	<p><b>M1</b></p> <p><b>M1A1</b></p> <p><b>A1</b></p>	<p>Showing all details</p> <p>Condone 18°</p> <p>Alt. method  <math>\mathbf{s} = \begin{pmatrix} 2\sqrt{2}t \\ 5000 + 2\sqrt{2}t \end{pmatrix}</math> <b>M1</b>  <math>\mathbf{p} = \begin{pmatrix} 9(\sin \alpha^\circ)t \\ 9(\cos \alpha^\circ)t \end{pmatrix}</math> <b>M1</b>                      (condone switched)  <math>\sin \alpha^\circ = \frac{2\sqrt{2}}{9}</math> <b>A1</b>  <math>\alpha = 018</math> <b>A1</b></p>
11(b)	$\text{Time} = \frac{5000}{9 \cos \alpha - 4 \sin 45^\circ}$ $\text{Time} = 875 \text{ seconds}$	<p><b>M1A1</b></p> <p><b>A1</b></p>	<p>Use of 5 instead of 5000 is <b>M1A0</b>  <math>\frac{v}{\sin 26.7^\circ} = \frac{9}{\sin 135^\circ}</math> and <math>t = \frac{5000}{v}</math> is <b>M1A1</b>                      oe                      Accept [871, 879]</p>
	<b>Total</b>	<b>7</b>	

Q	Answer	Marks	Comments
12	<p>For A:</p> $-9 = 2 \times v_A - 2 \times 5$ $v_A = 0.5$ <p>For B:</p> $9 = 3 \times v_B - 3 \times (-2)$ $v_B = 1$ $0.5 - 1 = -e(5 - (-2))$ $e = \frac{-0.5}{-7} = \frac{1}{14}$	<p><b>M1</b> <b>A1</b></p> <p><b>M1</b> <b>A1</b></p> <p><b>A1</b></p>	Or for a correct conservation of momentum equation with one unknown
	<b>Total</b>	<b>5</b>	