



---

# INTERNATIONAL A-LEVEL MATHEMATICS

## MA04

(9660/MA04) Unit S2 Statistics

---

Mark scheme

June 2022

---

Version 1.0 Final



2 2 6 X M A 0 4 / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [oxfordaqaexams.org.uk](https://www.oxfordaqaexams.org.uk)

#### **Copyright information**

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2022 Oxford International AQA Examinations and its licensors. All rights reserved.

**Key to mark scheme abbreviations**

<b>M</b>	Mark is for method
<b>m</b>	Mark is dependent on one or more M marks and is for method
<b>A</b>	Mark is dependent on M or m marks and is for accuracy
<b>B</b>	Mark is independent of M or m marks and is for method and accuracy
<b>E</b>	Mark is for explanation
<b>✓ or ft</b>	Follow through from previous incorrect result
<b>CAO</b>	Correct answer only
<b>CSO</b>	Correct solution only
<b>AWFW</b>	Anything which falls within
<b>AWRT</b>	Anything which rounds to
<b>ACF</b>	Any correct form
<b>AG</b>	Answer given
<b>SC</b>	Special case
<b>oe</b>	Or equivalent
<b>A2, 1</b>	2 or 1 (or 0) accuracy marks
<b>–x EE</b>	Deduct x marks for each error
<b>NMS</b>	No method shown
<b>PI</b>	Possibly implied
<b>SCA</b>	Substantially correct approach
<b>sf</b>	Significant figure(s)
<b>dp</b>	Decimal place(s)

Q	Answer	Marks	Comments
1(a)	A random variable	E1	
	that is a function of known observations from a population	E1	oe
		2	
1(b)	Range of values [of the test statistic]	E1	oe
	that leads us to determine whether or not the null hypothesis is to be rejected or not	E1	oe
		2	
	Total	4	

Q	Answer	Marks	Comments
2(a)	$\lambda = 0.1$	B1	
	$P(T < 5) = 1 - e^{-0.1 \times 5}$	M1	ft their value for $\lambda$
	$= 0.3935$	A1	AWRT 0.3935
		3	
2(b)	$P(8 < T < 14)$		
	$= (1 - e^{-0.1 \times 14}) - (1 - e^{-0.1 \times 8})$	M1	Difference between two probabilities with at least one correct probability
	$[= 0.75340... - 0.55067...]$		
	$= 0.2027$	A1	AWRT 0.2027
		2	
	Total	5	

Q	Answer	Marks	Comments
3(a)	$\int_0^a \frac{4}{(2x+1)^2} dx = 1$ $\left[ \frac{4}{2 \times -1} (2x+1)^{-1} \right]_0^a = 1$ $\frac{-2}{2a+1} - (-2) = 1$ $a = 0.5$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>Correct definite integral set equal to 1 Condone missing dx</p> <p>Correct integration <b>oe</b></p> <p><b>ft</b> from their integration with correct limits substituted</p> <p><b>AG</b> Be convinced</p>
		<b>4</b>	
3(b)	$P(0.2 < X < 0.4) = \int_{0.2}^{0.4} \frac{4}{(2x+1)^2} dx$ $= \left[ \frac{4}{2 \times -1} (2x+1)^{-1} \right]_{0.2}^{0.4}$ $= (-2) \times (2 \times 0.4 + 1)^{-1} - (-2) \times (2 \times 0.2 + 1)^{-1}$ $= \frac{20}{63}$	<p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>Correct definite integral Condone missing dx <b>PI</b></p> <p>Applies limits to their integration from <b>part (a)</b> <b>PI</b></p> <p><b>oe, AWRT</b> 0.317</p>
		<b>3</b>	
	<b>Total</b>	<b>7</b>	

Q	Answer	Marks	Comments
4	$H_0: \mu = 16$ $H_1: \mu < 16$  $\bar{x} = 15.5$  $s^2 = \frac{1}{200-1} \left( 50300 - \frac{3100^2}{200} \right)$  $= 11.3[0653266]$  $\bar{X} \sim N \left( 16, \frac{11.3}{200} \right)$  $z = \frac{15.5-16}{\sqrt{\frac{11.3}{200}}}$  $= -2.1[0351581]$  $z_{\text{critical}} = -2.3263$  As $z > z_{\text{critical}}$ we fail to reject $H_0$  Evidence to suggest that the laptop battery time between charges has not decreased	<b>B1</b>  <b>B1</b>  <b>M1</b>  <b>A1</b>  <b>M1</b>  <b>M1</b>  <b>A1</b>  <b>B1</b>  <b>A1ft</b>  <b>E1</b>	Both hypotheses  <b>PI</b> by correct calculation for $z$  Attempt at variance formula Allow one slip <b>PI</b> by correct answer  <b>AWRT</b> $\left( \frac{2250}{199} \right)$ Accept $s = 3.36[2518297]$  $\bar{X} \sim N \left( 16, \frac{s^2}{200} \right)$ <b>PI</b>  Calculates $z$ with their $s^2$  <b>AWRT</b> $-2.1$  <b>AWRT</b> $-2.3$ or $P(z < -2.1) = 0.0177$ Accept $2.3$  Follow through their $z$ and $z_{\text{critical}}$ provided signs are consistent or comparing $0.0177$ to $1\%$  Must be consistent with their conclusion on whether to fail to reject $H_0$ or not or their $z$ and $z_{\text{critical}}$ if not explicitly stated
	<b>Total</b>	<b>10</b>	

Q	Answer	Marks	Comments
5(a)	Symmetrical  Mode = Mean = Median  95% of the data lies within 2 sd of the mean	<b>B1</b>  <b>B1</b>  <b>B1</b>	oe such as 'bell-shaped curve' or 'no skew'  Accept » instead of =  Accept similar accurate comment eg 99% P 3 sd 68% P 1 sd
		<b>3</b>	
5(b)(i)	$P(X > 17) = P\left(Z > \frac{17-14}{1.2}\right)$ $= P(Z < -2.5)$ $= 1 - 0.99379[03346]$ $= 0.0062$	<b>M1</b>  <b>M1</b>  <b>A1</b>	<b>PI</b> Standardises  <b>PI</b> by sight or use of 0.99379...  <b>AWRT</b> 0.0062
		<b>3</b>	
5(b)(ii)	$\text{Var}(\bar{X}) = \frac{1.2^2}{50}$ or $\text{sd} = \frac{1.2}{\sqrt{50}}$  $P(\bar{X} < 13.8) = P\left(Z < \frac{13.8-14}{\sqrt{0.0288}}\right)$ $= P(Z < -1.1785)$ $= 1 - 0.8807[035853]$ $= 0.119$	<b>B1</b>  <b>M1</b>    <b>m1</b>  <b>A1</b>	<b>PI</b> Accept $\text{Var}(\bar{X}) = 0.0288$ or <b>AWRT</b> $\text{sd} = 0.17$  <b>PI</b> Standardises using their $\text{Var}(\bar{X})$ but not $1.2^2$  <b>PI</b> by correct answer  <b>AWRT</b> 0.119
		<b>4</b>	
5(b)(iii)	$X + Y \sim N(44, 1.2^2 + 4^2)$  $P(X + Y < 35) = P\left(Z < \frac{35-44}{\sqrt{17.44}}\right)$ $= P(Z < -2.1551)$ $= 1 - 0.9844[233527]$ $= 0.0156$	<b>B1</b>  <b>M1</b>    <b>m1</b>  <b>A1</b>	<b>PI</b>  <b>PI</b> Standardises with their sum of means and their sum of variances    <b>PI</b> by correct answer  <b>AWFW</b> 0.01539 to 0.01578
		<b>4</b>	
	<b>Total</b>	<b>14</b>	

Q	Answer	Marks	Comments
6(a)	$P(X=3) = \frac{e^{-4} \times 4^3}{3!}$	M1	PI May use tables: 0.4335 – 0.2381
	= 0.195	A1	AWRT 0.195
		2	
6(b)	$\lambda = 2 \times (4 + 2.5)$	M1	PI
	$\lambda = 13$	A1	
	$H_0: \lambda = 13$ $H_1: \lambda < 13$	B1	Both hypotheses Allow $H_0: \lambda = 6.5$ $H_1: \lambda < 6.5$
	$P(X \leq 7)$	M1	Attempts $P(X \leq 7)$ or $P(X < 7)$
	$P(X \leq 7) = 0.054$	A1	AWRT 0.054
	$0.054 > 0.05$	M1	Compares their probability with 0.05
	Do not reject $H_0$	A1ft	ft their probability compared with 0.05 Implied by correct conclusion in context
	Evidence to suggest that there has not been a reduction in the total number of breakdowns of boats and buses	E1	Must be consistent with their conclusion on whether or not to reject $H_0$ or on their probability if not explicitly stated
		8	
	Total	10	



Q	Answer	Marks	Comments
7(a)	$X \sim B(100, 0.03)$ $100 \times 0.03 = 3$ $Y \sim Po(3)$ $P(Y \leq 3)$ $= 0.647$	<b>B1</b>  <b>B1</b>  <b>M1</b>  <b>A1</b>	<b>PI</b> by use of $\lambda = 3$ with Poisson distribution Identifies correct approximate distribution Identifies correct probability <b>AWRT</b> 0.647
		<b>4</b>	
7(b)(i)	$H_0: p = 0.03$ $H_1: p \neq 0.03$ $X \sim B(20, 0.03)$ $P(X \geq 3) = 0.021$ $0.021 < 0.025$ Reject $H_0$ Evidence to suggest that the proportion of viewers watching the local news programme has changed	<b>B1</b>  <b>M1</b>  <b>A1</b>  <b>M1</b>  <b>A1ft</b>  <b>E1</b>	Both hypotheses <b>PI</b> by a binomial probability calculation <b>AWRT</b> 0.021 Compares their probability with 0.025 Follow through their probability Implied by correct conclusion in context Must be in context, must not be definite and all the previous 5 marks must have been awarded.
		<b>6</b>	
7(b)(ii)	A Type I error means to reject that the proportion of viewers watching the local news programme is 3% when it is 3%	<b>E2</b>	<b>oe</b> <b>E1</b> for describing Type I error without context
		<b>2</b>	
7(b)(iii)	$P(X \geq 3) = 0.021$ [ $< 0.025$ ] $P(X \geq 2) = 0.1198$ [ $> 0.025$ ] $P(X \leq x) < 0.025$ $P(X = 0) = 0.5438$ [ $> 0.025$ ] $P(\text{Type I error}) = 0.021$	<b>M1</b>  <b>M1</b>  <b>A1</b>	Considers both probabilities <b>PI</b> by calculation $P(X = 0)$ or $P(X \leq 0)$ <b>oe</b>
		<b>3</b>	
	<b>Total</b>	<b>15</b>	

Q	Answer	Marks	Comments
8(a)	$\frac{0.4}{2}x$ or $0.2x$ $y - 0.4 = \frac{0.6}{4}(x - 2)$ or $y - 1 = \frac{0.6}{4}(x - 6)$ $0.15x + 0.1$ $F(x) = \begin{cases} 0 & x < 0 \\ 0.2x & 0 \leq x < 2 \\ 0.15x + 0.1 & 2 \leq x < 6 \\ 1 & x \geq 6 \end{cases}$	<b>B1</b>  <b>M1</b>  <b>A1</b>  <b>B1</b>  <b>A1</b>	Seen anywhere  Use of straight line methods to find second line Allow $y = mx + c$ methods only if a value is found for $c$  Seen anywhere  0 when $x < 0$ and 1 when $x \geq 6$ Allow either strict or non-strict inequalities  Completely defined function in <b>ACF</b> Allow different but consistent placement of strict inequalities
		<b>5</b>	
8(b)	$f(x) = \begin{cases} 0.2 & 0 \leq x < 2 \\ 0.15 & 2 \leq x < 6 \\ 0 & \text{otherwise} \end{cases}$	<b>B1</b>  <b>M1</b>  <b>A1ft</b>	0 and otherwise <b>oe</b> seen  0.2 <b>oe</b> or 0.15 <b>oe</b> seen anywhere <b>ft</b> their equations of lines from <b>part (a)</b>  Completely defined function in <b>ACF</b> <b>ft</b> their equations of lines from <b>part (a)</b>
		<b>3</b>	

Q	Answer	Marks	Comments
<b>8(c)</b>	$E(X^3) = \int_0^2 0.2x^3 dx + \int_2^6 0.15x^3 dx$	<b>M1</b>	Identifies correct integral for their f(x)
	$= \left[ \frac{0.2x^4}{4} \right]_0^2 + \left[ \frac{0.15x^4}{4} \right]_2^6$	<b>A1</b>	Correct integration
	$E(X^6) = \int_0^2 0.2x^6 dx + \int_2^6 0.15x^6 dx$	<b>M1</b>	Identifies correct integral for their f(x)
	$= \left[ \frac{0.2x^7}{7} \right]_0^2 + \left[ \frac{0.15x^7}{7} \right]_2^6$	<b>A1</b>	Correct integration
	$E(X^3) = 48.8 \text{ or } E(X^6) = \frac{209984}{35}$	<b>A1</b>	<b>PI, AWRT</b> 6000 for $E(X^6)$
	$\text{Var}(X^3) = E(X^6) - E(X^3)^2$		
	$\text{Var}(X^3) = \frac{209984}{35} - 48.8^2$	<b>m1</b>	Uses variance formula with their expectation values for $E(X^6)$ and $E(X^3)$
	$\text{Var}(X^3) = 3620$	<b>A1</b>	<b>AWRT</b> 3620
		<b>7</b>	
	<b>Total</b>	<b>15</b>	