

OXFORD

INTERNATIONAL
AQA EXAMINATIONS

INTERNATIONAL AS MATHEMATICS MA02

(9660/MA02) Unit PSM1 Pure Mathematics, Statistics and Mechanics

Mark scheme

June 2023

Version: 1.0 Final



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from oxfordaqaexams.org.uk


Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2023 Oxford International AQA Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

M	Mark is for method
m	Mark is dependent on one or more M marks and is for method
A	Mark is dependent on M or m marks and is for accuracy
B	Mark is independent of M or m marks and is for method and accuracy
E	Mark is for explanation
✓ or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
–x EE	Deduct x marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)

Q	Answer	Marks	Comments
1(a)	See artwork below	B1 B1	Decreasing exponential curve of the correct form in the first and second quadrants asymptotic to the positive x -axis. Correct value of y -intercept indicated. Allow correct coordinates instead of value.
			
		2	

Q	Answer	Marks	Comments
1(b)	$\left[y = \frac{1}{9^{(2 \log_9 a - 0.5)}} = \frac{1}{9^{2 \log_9 a} \times 9^{-0.5}} \right]$ $\left[9^{2 \log_9 a} = \right] a^2 \text{ or } \left[9^{-2 \log_9 a} = \right] a^{-2}$ $\frac{3}{a^2}$	M1 A1ft	PI Expressing $9^{2 \log_9 a}$ or $9^{-2 \log_9 a}$ as the correct power of a Correct y -coordinate of P in the correct form. ft their value for the y -intercept from part (a) for '3' provided it is positive. Allow $3a^{-2}$ for $\frac{3}{a^2}$
		2	

	Question 1 Total	4	
--	------------------	---	--

Q	Answer	Marks	Comments
2(a)	<p>[Arc Length =] 10θ</p> <p>or</p> <p>$AB = 6$ [cm]</p> <p>$10\theta + 10 + 10 = 26$</p> <p>or</p> <p>$10\theta + 20 = 26$</p> <p>and</p> <p>$\theta = 0.6$</p>	<p>M1</p> <p>A1</p>	<p>Correct length of arc AB in terms of θ or identifies that arc AB is 6 [cm] Possibly embedded in later working.</p> <p>Sets expression for perimeter equal to 26. PI by $10\theta = 6$ and AG Must be convincingly shown</p>
		2	

Q	Answer	Marks	Comments
2(b)	$\frac{\sin(\angle BOC)}{14} = \frac{\sin(0.7)}{10}$ $\left[\angle BOC = \sin^{-1}\left(\frac{14\sin(0.7)}{10}\right) = \right]$ $1.12[415...] \text{ [radians]}$ $[\angle OBC =] \pi - 0.7 - 1.12[415...]$ $[\angle OBC =]$ 1.32 [radians]	<p>M1</p> <p>A1</p> <p>m1</p> <p>A1</p>	<p>PI oe Use of sine rule with correct values substituted. Allow AWRT 40.1° for 0.7 radians.</p> <p>Correct $\angle BOC$ AWRT 1.12 Allow AWRT 64.4°</p> <p>oe ft their $\angle BOC$ provided M1 scored. Allow angles in degrees. PI by final answer of 1.317[43...] rounded or truncated to at least 3dp or AWRT 75.5°</p> <p>AWRT 1.32 Must be in radians.</p>
2(b) ALT	$[\text{Length of } OC = x]$ $10^2 = 14^2 + x^2 - 2 \times 14 \times x \times \cos(0.7)$ <p>or</p> $\cos(0.7) = \frac{14^2 + x^2 - 10^2}{2 \times 14 \times x}$ $[x^2 - 21.4[1558...]x + 96 = 0 \Rightarrow]$ $[x =] 15.0[2714...]$ $\frac{\sin(\angle OBC)}{15.0[2714...]} = \frac{\sin(0.7)}{10}$ <p>or</p> $\cos(\angle OBC) = \frac{10^2 + 14^2 - (15.0[2714...])^2}{2 \times 10 \times 14}$ $[\angle OBC =]$ 1.32 [radians]	<p>M1</p> <p>A1</p> <p>m1</p> <p>A1</p>	<p>oe Correct use of cosine rule with values substituted. Allow AWRT 40.1° for 0.7 radians.</p> <p>Correct length of OC AWRT 15 Condone 6.38844... seen as well.</p> <p>oe ft their length of OC provided M1 scored. Correct use of sine rule or cosine rule with values substituted. Allow AWRT 40.1° for 0.7 radians. PI by correct final answer or anything that truncates to 75°</p> <p>AWRT 1.32 Must be in radians. Accept AWRT 1.31</p>
		4	

Q	Answer	Marks	Comments
2(c)	$\left[\frac{1}{2} \times 10 \times 10 \times 0.6 = \right] 30 \text{ [cm}^2\text{]}$ $\frac{1}{2} \times 10 \times 14 \times \sin(1.31[743\dots])$ $67.76[524\dots] \text{ [cm}^2\text{]}$ $97.8 \text{ [cm}^2\text{]}$	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>PI Correct area of sector <i>OAB</i></p> <p>oe Correct method for calculating the area of triangle <i>OBC</i> with values substituted. ft their $\angle OBC$</p> <p>Correct area of triangle <i>OBC</i> PI by correct final answer. AWFW 67.55 to 67.85</p> <p>CAO AWFW 97.55 to 97.85</p>
		4	
	Question 2 Total	10	

[illegible]

Q	Answer	Marks	Comments
3(b)(i)	$x - 5y = 22$ and $y = 8x - 59$ $(7, -3)$	M1 A1	M1 : Correct x -coordinate or y -coordinate. A1 : Correct coordinates.
		2	

Q	Answer	Marks	Comments
3(b)(ii)	$(8-7)^2 + (5-(-3))^2$ or $\sqrt{(8-7)^2 + (5-(-3))^2}$ or $(11-7)^2 + ((-10)-(-3))^2$ or $\sqrt{(11-7)^2 + ((-10)-(-3))^2}$ $[r =] \sqrt{65}$ or $[r^2 =] 65$ $(x-7)^2 + (y+3)^2 = 65$	<p>M1</p> <p>A1ft</p> <p>A1ft</p>	<p>Method to find the radius or the square of the radius of <i>C</i> using either the coordinates of <i>P</i> or <i>Q</i> ft their centre of <i>C</i></p> <p>ft their centre of <i>C</i></p> <p>Correct equation in the correct form. ft their centre and r^2 provided all values are integers and M1 scored.</p>
		3	

Q	Answer	Marks	Comments
3(c)	$\left[d^2 = (7-2)^2 + (-3-(-9))^2 \right]$ $[d^2 =] 61 \text{ or } [d =] \sqrt{61}$ <p>Since $\sqrt{61} < \sqrt{65}$ (or $61 < 65$) then R lies inside the circle.</p>	<p>B1ft</p> <p>E1ft</p>	<p>Correct distance or square of distance from centre of C to R ft their (b)(i) or (b)(ii) Allow 7.8[1024...] for $\sqrt{61}$</p> <p>Compares their $\sqrt{61}$ with their $\sqrt{65}$ oe and gives a correct conclusion. ft their distance or square of distance from centre of C to R, and their r or r^2 provide both coordinates of the centre of C are integers.</p> <p>Allow 7.8[1024...] for $\sqrt{61}$ and 8[.0622...] for $\sqrt{65}$</p>
		2	

	Question 3 Total	10	
--	-------------------------	-----------	--

[illegible]

	Question 4 Total	9	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
5(a)	$\frac{\sin \theta}{1 + \cos \theta} + \frac{1}{\frac{\sin \theta}{\cos \theta}}$ or $\frac{\sin \theta}{1 + \cos \theta} + \frac{\cos \theta}{\sin \theta}$ $\frac{\sin^2 \theta + (1 + \cos \theta) \cos \theta}{(1 + \cos \theta) \sin \theta}$ or $\frac{\sin^2 \theta + \cos^2 \theta + \cos \theta}{(1 + \cos \theta) \sin \theta}$ or $\frac{\sin \theta (1 - \cos \theta)}{\sin^2 \theta} + \frac{\sin \theta \cos \theta}{\sin^2 \theta}$ $\frac{1 + \cos \theta}{(1 + \cos \theta) \sin \theta} \text{ or } \frac{\sin \theta}{\sin^2 \theta} \text{ or } \frac{1 - \cos \theta}{\sin \theta} + \frac{\cos \theta}{\sin \theta}$ and $\frac{1}{\sin \theta}$	<p>M1</p> <p>M1</p> <p>A1</p>	<p>Use of $\tan \theta = \frac{\sin \theta}{\cos \theta}$</p> <p>oe Rearrangement to give a correct expression in terms of $\sin \theta$ and $\cos \theta$ with a common denominator. Allow $1 - \cos^2 \theta$ for $\sin^2 \theta$</p> <p>Uses $\sin^2 \theta + \cos^2 \theta = 1$ AG Must be convincingly shown.</p>
		3	

Q	Answer	Marks	Comments
5(b)	$\frac{2}{\sin 2x} = 4 \sin 2x$ $2x = \sin^{-1} \left(\frac{1}{\sqrt{2}} \right)$ $[x =] 22.5^\circ, 67.5^\circ$	<p>B1</p> <p>M1</p> <p>A2,1</p>	<p>oe, condone θ for $2x$ throughout.</p> <p>PI by 45° or 135° or one correct final answer. Ignore $2x = \sin^{-1} \left(-\frac{1}{\sqrt{2}} \right)$</p> <p>A1: At least one correct answer. A2: Both correct answers with no others seen.</p>
		4	

	Question 5 Total	7	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
6(a)	$[\text{Variance} = 40 \times 0.4(1 - 0.4)] = 9.6$	B1	oe
		1	

Q	Answer	Marks	Comments
6(b)	$P(L = 19) = \binom{40}{19} \times 0.4^{19} \times (1 - 0.4)^{40-19}$ or $0.8702 - 0.7911$ $= 0.079$	M1 A1	oe, PI AWRT 0.079 Uses correct formula for $P(L = 19)$ or uses $P(L \leq 19) - P(L \leq 18)$ CAO
		2	

Q	Answer	Marks	Comments
6(c)	$[P(L > 13) = 1 - P(L \leq 13)]$ $= 1 - 0.2112$ $= 0.789$	M1 A1	PI Uses formula. AWRT 0.789
		2	

	Question 6 Total	5	
--	------------------	---	--

Q	Answer	Marks	Comments
7(a)(i)	$[a =] 1$	B1	
	$[b =] 4$	B1	
		2	

Q	Answer	Marks	Comments
7(a)(ii)	$0.4 \times 1 + 0.3 \times 4 + 0.3c = 3.4$	M1	Applies expectation formula for their a and b and sets equal to 3.4
	$[c =] 6$	A1	
		2	

Q	Answer	Marks	Comments
7(b)	$[E(X^2) =]$ $0.4 \times 1^2 + 0.3 \times 4^2 + 0.3 \times 6^2 [= 16]$	M1	Applies formula for $E(X^2)$ for their values of a , b and c PI by correct variance for their values of a , b and c
	$[Var(X) = 16 - 3.4^2 =]$ $4.44 \text{ or } \frac{111}{25}$	A1ft	Correctly finds variance for their values of a , b and c Must use $E(X) = 3.4$
	$Var(X + Y) = 17.44 \text{ or } \frac{436}{25}$	A1ft	ft their $Var(X) + 13$ Dependent on at least M1 awarded
		3	

	Question 7 Total	7	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
8(a)	0.24×0.74	M1	
	$= 0.1776$ or $\frac{111}{625}$	A1	
		2	

Q	Answer	Marks	Comments
8(b)	$0.24 + 0.61 - 0.1776$	M1	Applies the Addition rule with their part (a) [0.1776]
	$= 0.6724$ or $\frac{1681}{2500}$	A1ft	ft their part (a) [0.1776] provided final answer is between 0 and 1
		2	

Q	Answer	Marks	Comments
8(c)	$0.24 - 0.1776$ or $0.6724 - 0.61$ or 0.24×0.26	M1	ft 0.24 – their part (a) or their part (b) – 0.61
	$= 0.0624$ or $\frac{39}{625}$	A1	CAO
		2	

Q	Answer	Marks	Comments
8(d)	$\frac{0.0624}{1 - 0.61}$	M1	oe, ft their part (c)
	$= 0.16$ or $\frac{4}{25}$	A1	CAO
		2	

	Question 8 Total	8	
--	------------------	---	--

Q	Answer	Marks	Comments
9(a)	$2 - 2 \times 0.6 - k = 0$	M1	oe PI by correct answer
	or $[k =] 2 - 2 \times 0.6$		
	$[k =] 0.8$	A1	
		2	

Q	Answer	Marks	Comments
9(b)	0.6 N	B1	Condone omission of units
		1	

	Question 9 Total	3	
--	------------------	---	--

Q	Answer	Marks	Comments
10(a)	$\left[v = \int (0.72 - 0.18t) dt \right]$		
	$[v =] \ 0.72t - 0.09t^2$	B1	oe Correct expression for v Allow '+c'
	$0 = 0.72 - 0.18t$	M1	Uses $a = 0$ oe, PI
	$[t =] \ 4 \text{ [seconds]}$	A1	
	$[v =] \ 0.72t - 0.09t^2$		
	$[v_{\max} =] \ 0.72 \times 4 - 0.09 \times 4^2$	m1	oe Substitutes their 4 into their integrated expression for v
	$[v_{\max} =] \ 1.44 \text{ ms}^{-1}$	A1	CAO Condone omission of units
		5	

Q	Answer	Marks	Comments
10(b)(i)	$[v^2 = u^2 + 2as]$ $v^2 = 0^2 + 2 \times 9.8(7.68 - 2)$ $[v =] 10.55$ ms^{-1}	<p>M1</p> <p>A1</p> <p>B1</p>	<p>oe If more than one constant equation formula used it must be a complete method. PI by AWRT 111.3 Condone one sign error AWRT 11</p> <p>Exact answer is $\frac{14\sqrt{355}}{25}$</p> <p>Correct units.</p>
		3	

Q	Answer	Marks	Comments
10(b)(ii)	$\left[s = \frac{1}{2}(u + v)t\right]$ $[s =] \frac{1}{2} \times (0 + 10.55) \times 0.3$ $[s =] 1.58 \text{ [metres]}$ $[\text{Height} = 2 - 1.58 =] 0.42 \text{ metres}$	<p>M1</p> <p>A1</p> <p>B1ft</p>	<p>oe PI ft their answer to part (b)(i) If more than one constant equation formula used it must be a complete method.</p> <p>AWRT 1.58, allow 1.59 PI by correct final answer</p> <p>AWRT 0.42, allow 0.41 ft $0 < \text{their } 1.58 < 2$ Condone omission of units</p>
		3	

	Question 10 Total	11	
--	--------------------------	-----------	--

Q	Answer	Marks	Comments
11(a)	Conservation of Momentum		
	$4 \times 4.8 = 4v + 3m$	M1	oe Correct unsimplified equation. Allow sign error.
	$[4v = 19.2 - 3m]$		
	$v = 4.8 - 0.75m$	A1	AG Must be convincingly shown
		2	

Q	Answer	Marks	Comments
11(b)	$0 < 4.8 - 0.75m \quad [\Rightarrow m < 6.4]$	M1	oe Considers inequality or equality for $v = 0$ PI by 6.4 Condone equality or weak inequality
	$3 \geq 4.8 - 0.75m \quad [\Rightarrow m \geq 2.4]$	M1	oe Considers inequality or equality for $v = 3$ PI by 2.4 Condone equality or strict inequality
	$2.4 \leq m$ or $m < 6.4$	A1	oe At least one inequality correct For one of $2.4 \leq m$ or $m < 6.4$ Accept $2.4 < m$ for $2.4 \leq m$ but not $m \leq 6.4$ for $m < 6.4$
	$2.4 \leq m < 6.4$	A1	oe Both inequalities correct Accept $2.4 < m < 6.4$ but not $2.4 \leq m \leq 6.4$
		4	

	Question 11 Total	6	
--	-------------------	---	--