

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

--

Forename(s)

--

Candidate signature

--

I declare this is my own work.

INTERNATIONAL A-LEVEL FURTHER MATHEMATICS

(9665/FM03) Unit FP2 Pure Mathematics

Wednesday 24 May 2023 07:00 GMT Time allowed: 2 hours 30 minutes

Materials

- For this paper you must have the Oxford International AQA Booklet of Formulae and Statistical Tables (enclosed).
- You may use a graphical calculator.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 120.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.

For Examiner's Use

Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
TOTAL	



J U N 2 3 F M 0 3 0 1

IB/G/Jun23/E8

FM03

Answer **all** questions in the spaces provided.

1 The 3×3 matrix **N** represents a reflection in the plane $y = 0$

The 3×3 matrix **M** represents an enlargement, scale factor 2 with the origin as the centre of enlargement.

Find the matrix **NM**

[2 marks]

NM =

2



2 The cubic equation

$$z^3 - 4z^2 + 3z + c = 0$$

where c is a non-zero constant, has roots α , β and γ

2 (a) Show that

$$\alpha^2 + \beta^2 + \gamma^2 = 10$$

[3 marks]

2 (b) Explain why

$$\beta^3 - 4\beta^2 + 3\beta + c = 0$$

[1 mark]

2 (c) Show that

$$\alpha^3 + \beta^3 + \gamma^3 = 28 - 3c$$

[2 marks]



- 3** Two 3×3 matrices **A** and **B** are such that

$$\det(\mathbf{AB}) = 10 \quad \text{and} \quad \det(\mathbf{A}^{-1}) = 5$$

A three-dimensional shape S_1 with volume 6 cm^3 is mapped onto the shape S_2 by the transformation represented by matrix **B**

Find the volume of S_2

[4 marks]

Answer _____

4

- 4** A curve has Cartesian equation

$$y = x\sqrt{x} - \frac{1}{3}\sqrt{x}$$

- 4 (a)** Show that

$$1 + \left(\frac{dy}{dx}\right)^2 = \left(px^n + qx^{-n}\right)^2$$

where p , q and n are rational numbers.

[2 marks]



[illegible]

[4 marks]

[illegible]

5

$$\int_0^e \left(9x^2 \ln x + \frac{4}{1+4x^2} \right) dx$$

[7 marks]

[illegible]

7



$$\frac{dy}{dx} + \frac{8x}{x^2+2}y = 2x^3 + \frac{1}{(x^2+2)^{\frac{9}{2}}}$$

[7 marks]

[illegible]

7

7

$$u_1 = 3 \qquad u_{r+1} = 3u_r + 4$$

7 (a)

$$\sum_{r=1}^n u_r = \frac{1}{2}u_{n+1} - 2n - \frac{3}{2}$$

[3 marks]

[illegible]

[4 marks]

[illegible]

[1 mark]

Turn over ►



- 8 (a)** Find the general solution of the differential equation

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} = 0$$

[2 marks]

Answer _____



8 (b)

and when $x=0$ it is given that both $y=0$ and $\frac{d^2y}{dx^2}=4$

[7 marks]

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.
$$y =$$


9

$$\mathbf{A} = \begin{bmatrix} 3-k & 1-k & 3 \\ 5 & 7 & 4 \\ 3 & 5 & 3 \end{bmatrix}$$

where k is a **positive** constant.

9 (a)

[6 marks]

This image shows a single sheet of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

$$\mathbf{A}^{-1} =$$



9 (b) (i) Use your answer to **part (a)** to solve the equations

$$(3-k)x + (1-k)y + 3z = 1$$

$$5x + 7y + 4z = 1$$

$$3x + 5y + 3z = 1$$

Give your solution in terms of k

[3 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

$x =$ _____ $y =$ _____ $z =$ _____

9 (b) (ii) Hence, state the range of possible values of $x + y + z$

[1 mark]

Answer

10

Turn over ►



10 (a) (i) Write down $e^{\frac{i\theta}{2}} + e^{-\frac{i\theta}{2}}$ in terms of $\cos\left(\frac{\theta}{2}\right)$

[1 mark]

$$e^{\frac{i\theta}{2}} + e^{-\frac{i\theta}{2}} =$$

10 (a) (ii) Hence, given that $e^{i\theta} \neq -1$ show that

$$\frac{1}{e^{i\theta} + 1} = \frac{1}{2} - \frac{i}{2} \tan\left(\frac{\theta}{2}\right)$$

[2 marks]

10 (b) Hence, by replacing θ by $\pi - \theta$ in the equation in **part (a)(ii)**, show that for $e^{i\theta} \neq 1$

$$\frac{1}{e^{i\theta} - 1} = -\frac{1}{2} - \frac{i}{2} \cot\left(\frac{\theta}{2}\right)$$

[3 marks]



10 (c) Deduce that, for $e^{2i\theta} \neq 1$

$$\frac{1}{\cos 2\theta - 1 + i \sin 2\theta} = a + i b \left(\tan\left(\frac{\theta}{2}\right) - \cot\left(\frac{\theta}{2}\right) \right)$$

where a and b are rational numbers.

[2 marks]



11 The line L has equation $\left(\mathbf{r} - \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \right) \times \begin{bmatrix} 2 \\ 3 \\ -6 \end{bmatrix} = \mathbf{0}$

The plane Π has equation $\mathbf{r} = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix} + \lambda \begin{bmatrix} -2 \\ 1 \\ 2 \end{bmatrix} + \mu \begin{bmatrix} 1 \\ -3 \\ 4 \end{bmatrix}$

11 (a) Find $\begin{bmatrix} -2 \\ 1 \\ 2 \end{bmatrix} \times \begin{bmatrix} 1 \\ -3 \\ 4 \end{bmatrix}$

[1 mark]

Answer _____

11 (b) Use a scalar triple product to determine whether or not $\begin{bmatrix} -4 \\ 3 \\ -2 \end{bmatrix}$, $\begin{bmatrix} -2 \\ 1 \\ 2 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ -3 \\ 4 \end{bmatrix}$ are coplanar vectors.

[2 marks]

Answer _____



[4 marks]

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Answer

[4 marks]

[illegible]

Question 11 continues on the next page

Turn over ►



Answer _____

- 11 (e)** Hence, or otherwise, find the **exact** value for the shortest distance from the point Q to the plane Π

[3 marks]

Answer _____



Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Turn over ►



12

$$y = e^{\frac{7}{25}x} \operatorname{sech} x$$

The curve has exactly one stationary point P

12 (a)

Find the x -coordinate of P giving your answer in the form $\ln k$ where k is a constant.

[6 marks]

[illegible]

Answer



Give your answer in the form $a(b)^c$ where a , b and c are rational numbers.

[illegible]

12 (c) Hence, determine whether or not the line L intersects the curve $y = \tanh x$. Justify your answer.

[illegible]

- 13 (a)** Write down the Maclaurin series expansion of $\ln(1+4x)$ in ascending powers of x up to and including the term in x^3 and state the range of values of x for which this expansion is valid.

[2 marks]

$\ln(1+4x) =$ _____ valid for _____

- 13 (b)** It is given that $y = \ln(\cos x - \sin x)$

- 13 (b) (i)** Show that

$$\frac{d^2y}{dx^2} = \frac{-2}{1 - \sin 2x}$$

[3 marks]



Maclaurin series expansion of $\ln(\cos x - \sin x)$ are $-x - x^2 - \frac{2}{3}x^3$

$$\lim_{x \rightarrow 0} \left[\frac{\ln((1 - \sin 2x) \sqrt{1 + 4x})}{5x^2 + 6x^3} \right]$$


[5 marks]

[illegible]

13



- 14** The diagram shows a sketch of a curve C , the pole O and the initial line.



The end points A and B of the curve C are shown on the diagram above.

The curve C has polar equation

$$r = \frac{3}{2} \operatorname{cosec}^2\left(\frac{\theta}{2}\right) \quad \text{for} \quad \frac{\pi}{4} \leq \theta \leq \frac{7\pi}{4}$$

- 14 (a)** The end point A has polar coordinates $\left(6 + 3\sqrt{2}, \frac{\pi}{4}\right)$

Show that the area of triangle AOB is $27 + 18\sqrt{2}$

[2 marks]



- 14 (b)** Find the Cartesian equation of C giving your answer in the form $y^2 = f(x)$

[4 marks]

Answer _____

- 14 (c)** The straight line with polar equation $\tan \theta = \sqrt{3}$ intersects the curve C at the points P and Q

- 14 (c) (i)** Find the polar coordinates of P and Q

[3 marks]

Answer _____

Question 14 continues on the next page

Turn over ►



[3 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


Answer _____



Do not write
outside the
box

14 (c) (iii) Hence, find the area of the region bounded by the curve C and the line segment PQ giving your answer in the form $k\sqrt{n}$ where k is a rational number and n is a prime number.

[3 marks]

[illegible]

Answer

END OF QUESTIONS

15



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



[illegible]

*Do not write
outside the
box*

[illegible]

*Do not write
outside the
box*

[illegible]

There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.oxfordaqaexams.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2023 Oxford International AQA Examinations and its licensors. All rights reserved.

