

OXFORD

INTERNATIONAL
AQA EXAMINATIONS

INTERNATIONAL AS FURTHER MATHEMATICS FM01

(9665/FM01) Unit FP1 Pure Mathematics

Mark scheme

January 2022

Version: 1.0 Final



2 2 1 X F M 0 1 / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from oxfordaqaexams.org.uk

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2022 Oxford International AQA Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

| | |
|----------------|--|
| M | Mark is for method |
| m | Mark is dependent on one or more M marks and is for method |
| A | Mark is dependent on M or m marks and is for accuracy |
| B | Mark is independent of M or m marks and is for method and accuracy |
| E | Mark is for explanation |
| ✓ or ft | Follow through from previous incorrect result |
| CAO | Correct answer only |
| CSO | Correct solution only |
| AWFW | Anything which falls within |
| AWRT | Anything which rounds to |
| ACF | Any correct form |
| AG | Answer given |
| SC | Special case |
| oe | Or equivalent |
| A2, 1 | 2 or 1 (or 0) accuracy marks |
| –x EE | Deduct x marks for each error |
| NMS | No method shown |
| PI | Possibly implied |
| SCA | Substantially correct approach |
| sf | Significant figure(s) |
| dp | Decimal place(s) |

| Q | Answer | Marks | Comments |
|---|--|--|--|
| 1 | $\left[\sum_{r=n+1}^{2n} r^3 = \right] \sum_{r=1}^{2n} r^3 - \sum_{r=1}^n r^3$ $= \frac{1}{4}(2n)^2(2n+1)^2 - \frac{1}{4}n^2(n+1)^2$ $= \frac{1}{4}n^2 \{4(2n+1)^2 - (n+1)^2\}$ $= \frac{1}{4}n^2 \{16n^2 + 16n + 4 - (n^2 + 2n + 1)\}$ $= \frac{1}{4}n^2 \{15n^2 + 14n + 3\}$ $= \frac{1}{4}n^2(5n+3)(3n+1)$ | <p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> | <p>If M0 awarded, allow SC1 for sight of $\frac{1}{4}(2n)^2(2n+1)^2$ and $\frac{1}{4}n^2(n+1)^2$</p> <p>Factorising at least n^2 using consistent working</p> <p>Expands the two squared brackets or uses difference of two squares Allow one slip</p> |
| | Question 1 Total | 5 | |

| Q | Answer | Marks | Comments |
|------|---|--|--|
| 2(a) | $\frac{7-3\mathrm{i}}{k-5\mathrm{i}} \times \frac{k+5\mathrm{i}}{k+5\mathrm{i}}$ Real part = $\frac{7k+15}{k^2+25}$ Imaginary part = $\frac{35-3k}{k^2+25}\mathrm{i}$ | <p>M1</p> <p>A1</p> <p>A1</p> | <p>or $z = x + \mathrm{i}y$ $(x + \mathrm{i}y)(k - 5\mathrm{i}) = 7 - 3\mathrm{i}$</p> <p>Then multiplies out and equates real and imaginary parts</p> <p>Seen anywhere</p> <p>Condone $\left(\frac{35-3k}{k^2+25}\right)\mathrm{i}$</p> |
| | | 3 | |

| Q | Answer | Marks | Comments |
|------|---|--|--|
| 2(b) | <p>substituting $k = 2$</p> <p>or</p> $\frac{35 - 3k}{7k + 15} \text{ seen}$ $\left[\frac{7 - 3i}{2 - 5i} = \frac{29}{29} + i \left(\frac{29}{29} \right) = \right] 1 + i$ <p>or</p> $\frac{35 - 3 \times 2}{7 \times 2 + 15}$ $\arg \left(\frac{7 - 3i}{2 - 5i} \right) = \arg(1 + i) = \left[\tan^{-1} \left(\frac{1}{1} \right) = \right] \frac{\pi}{4}$ | <p>M1</p> <p>A1</p> <p>A1</p> | <p>AG</p> <p>Condone $\tan \theta = 1 \Rightarrow \theta = \frac{\pi}{4}$</p> <p>where $\theta = \arg \left(\frac{7 - 3i}{2 - 5i} \right)$</p> |
| | | 3 | |

| | | | |
|--|-------------------------|----------|--|
| | Question 2 Total | 6 | |
|--|-------------------------|----------|--|

| Q | Answer | Marks | Comments |
|---|---|--|--|
| 3 | $A = \pi r^2$ $\frac{dA}{dr} = 2\pi r$ $\frac{dr}{dt} = \frac{dr}{dA} \times \frac{dA}{dt}$ $\frac{dr}{dt} = \frac{1}{2\pi r} \times 3$ When $A = 36\pi$, $r = 6$ $\frac{dr}{dt} = \frac{1}{2\pi(6)} \times 3$ $= \frac{1}{4\pi}$ [metres/day] | B1 M1 A1ft B1 M1 A1 | Seen or used their value for r may be substituted in ft their $\frac{dA}{dr}$ ft their $\frac{dr}{dt}$ and their value of r CAO |
| | Question 3 Total | 6 | |

| Q | Answer | Marks | Comments |
|------|--|---|--|
| 4(a) | $2x - \frac{\pi}{2} = 2n\pi \pm \frac{2\pi}{3}$ $x = \frac{1}{2} \left(2n\pi \pm \frac{2\pi}{3} + \frac{\pi}{2} \right)$ $x = n\pi + \frac{\pi}{4} \pm \frac{\pi}{3}$ | <p>B1</p> <p>M1</p> <p>A1 A1</p> | <p>oe</p> <p>Rearranging to make x the subject</p> <p>going from $\left(2x - \frac{\pi}{2} \right)$ to x</p> <p>Allow one slip</p> <p>oe, eg $x = n\pi + \frac{7\pi}{12}$ or $x = n\pi - \frac{\pi}{12}$</p> |
| | | 4 | |

| Q | Answer | Marks | Comments |
|------|--|--|--|
| 4(b) | $k = 1 : \frac{7\pi}{12}, \frac{11\pi}{12}$ $k = 2 : \text{also } \frac{19\pi}{12}, \frac{23\pi}{12}, \frac{31\pi}{12}, \frac{35\pi}{12}$ $k = 1: 2 \text{ solutions}$ $k = 2: 6 \text{ solutions [etc]}$ $4k - 2$ | <p>M1</p> <p>M1</p> <p>A1</p> | <p>For investigating at least one positive value of k with their general solution from part (a)</p> <p>For finding the number of solutions for at least two values of k using their general solution from part (a)</p> <p>CAO</p> |
| | | 3 | |

| | | | |
|--|-------------------------|----------|--|
| | Question 4 Total | 7 | |
|--|-------------------------|----------|--|

| Q | Answer | Marks | Comments |
|------|-----------------------|-----------|----------|
| 5(a) | $\alpha + \beta = -5$ | B1 | |
| | $\alpha\beta = 9$ | B1 | |
| | | 2 | |

| Q | Answer | Marks | Comments |
|------|--|-----------|-----------------------|
| 5(b) | $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta = 25 - 18$ | M1 | or other valid method |
| | $\alpha^2 + \beta^2 = 7$ | A1 | |
| | | 2 | |

| Q | Answer | Marks | Comments |
|------|--|-----------|-----------------------|
| 5(c) | $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$ | M1 | or other valid method |
| | $= -125 - 3(9)(-5) = 10$ | A1 | |
| | | 2 | |

| Q | Answer | Marks | Comments |
|------|--|-------------|--|
| 5(d) | Sum of roots $= \alpha + \frac{\beta}{\alpha} + \beta + \frac{\alpha}{\beta}$ | M1 | |
| | $= \alpha + \beta + \frac{\beta^2 + \alpha^2}{\alpha\beta}$ | | |
| | $= -5 + \frac{7}{9} = -\frac{38}{9}$ | A1ft | ft their $\alpha^2 + \beta^2$ from part (b) |
| | Product of roots $= \left(\alpha + \frac{\beta}{\alpha}\right)\left(\beta + \frac{\alpha}{\beta}\right)$ | M1 | |
| | $= \alpha\beta + \frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha} + 1$ | | |
| | $= \alpha\beta + \frac{\alpha^3 + \beta^3}{\alpha\beta} + 1$ | m1 | Converts $\frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha}$ into $\frac{\alpha^3 + \beta^3}{\alpha\beta}$ |
| | $= 9 + \frac{10}{9} + 1 = \frac{100}{9}$ | A1 | |
| | $9x^2 + 38x + 100 = 0$ | A1 | Correct quadratic equation with integer coefficients |
| | | 6 | |


| | | | |
|--|-------------------------|-----------|--|
| | Question 5 Total | 12 | |
|--|-------------------------|-----------|--|

| Q | Answer | Marks | Comments |
|------|---------|-------|----------|
| 6(a) | $x = 0$ | B1 | |
| | $y = 0$ | B1 | |
| | | 2 | |

| Q | Answer | Marks | Comments |
|------|---|-------|----------------------------------|
| 6(b) | $k = \frac{2x+1}{x^2}$ | M1 | |
| | $kx^2 - 2x - 1 = 0$ | | |
| | $\Delta \geq 0$ so $(-2)^2 - 4k(-1) \geq 0$ | B1 | Explicit use of the discriminant |
| | $4 + 4k \geq 0$ so $k \geq -1$ | A1 | AG |
| | | 3 | |

| Q | Answer | Marks | Comments |
|------|---|-------|----------|
| 6(c) | When $k = -1$, $-x^2 - 2x - 1 = 0$ $(x+1)^2 = 0 \Rightarrow x = -1$ | M1 | |
| | Stationary point is $(-1, -1)$ | A1 | |
| | | 2 | |

| Q | Answer | Marks | Comments |
|------|-----------|-----------|---|
| 6(d) | See below | B1 | Correct general shape |
| | See below | B1 | Axis intercept correctly labelled (condone x -coordinate only) and stationary point correctly marked and labelled |
| | See below | B1 | Graph approaches all asymptotes |



| | | | |
|--|--|----------|--|
| | | 3 | |
|--|--|----------|--|

| Q | Answer | Marks | Comments |
|------|---|-----------|---|
| 6(e) | $\frac{2x+1}{x^2} > 3 \therefore 2x+1 > 3x^2$ | M1 | Allow equation if followed by attempt to solve inequality |
| | $3x^2 - 2x - 1 < 0$ $(3x+1)(x-1) < 0$ | M1 | Or for solving the corresponding equation |
| | $-\frac{1}{3} < x < 1$ | A1 | PI |
| | $-\frac{1}{3} < x < 0, 0 < x < 1$ | A1 | ACF , e.g. $-\frac{1}{3} < x < 1$ and $x \neq 0$ |
| | | 4 | |

| | | | |
|--|-------------------------|-----------|--|
| | Question 6 Total | 14 | |
|--|-------------------------|-----------|--|

| Q | Answer | Marks | Comments |
|------|---------------------------------------|-------|---|
| 7(a) | Because one of the limits is infinite | E1 | Or 'the range of integration is infinite' |
| | | 1 | |


| Q | Answer | Marks | Comments |
|------|---|-------|----------|
| 7(b) | Because the integrand is not defined at one of the limits of integration or Because the integrand is not defined when $x = 0$ | E1 | |
| | | 1 | |

| Q | Answer | Marks | Comments |
|------|--|-------------------------------|---|
| 7(c) | $I_2 = \lim_{h \rightarrow 0} \int_h^{64} \frac{1}{(\sqrt[3]{x})^2} dx$ $\left[= \lim_{h \rightarrow 0} \int_h^{64} x^{-\frac{2}{3}} dx \right]$ $= \lim_{h \rightarrow 0} \left[3x^{\frac{1}{3}} \right]_h^{64}$ $= 3(4) - 3(0)$ $= 12$ | <p>M1</p> <p>m1</p> <p>A1</p> | <p>Limiting process seen in the solution</p> <p>Condone 0 as lower limit if 1st M1 was awarded</p> <p>Correct answer with no limiting process shown is SC1</p> |
| | | 3 | |

| | | | |
|--|------------------|---|--|
| | Question 7 Total | 5 | |
|--|------------------|---|--|


| Q | Answer | Marks | Comments |
|------|----------|-------|----------|
| 8(a) | $4 + 6i$ | B1 | |
| | | 1 | |

| Q | Answer | Marks | Comments |
|------|----------------------|-------|--|
| 8(b) | L drawn correctly | B1 | Condone no indication that end point is not included, but must not be below the real axis |
| | OP drawn correctly | B1 | No need to extend beyond O or P |
| | T marked correctly | B1ft | The intersection of their line OP with the correct half-line L See artwork below |




| | | | |
|--|--|---|--|
| | | 3 | |
|--|--|---|--|

| Q | Answer | Marks | Comments |
|------|--|-------|--|
| 8(c) | $\tan \beta = \frac{3}{2}$ | B1 | See diagram below |
| | $\left[\sin \theta = \frac{2}{\sqrt{13}} \text{ so } \right] \tan \theta = \frac{2}{3}$ | B1 | |
| | $\arg z = \alpha = \beta - \theta$ | M1 | |
| | $\tan \alpha = \frac{5}{12}$ [or $\alpha = 0.39479\dots$] | A1 | |
| | $z = 4 + (4 \tan \alpha)i$ | M1 | |
| | $z = 4 + \frac{5}{3}i$ | A1 | Exact value for real and imaginary parts |



| | | | |
|--|------------------|----|--|
| | | 6 | |
| | Question 8 Total | 10 | |


| Q | Answer | Marks | Comments |
|-------------|---|--|--|
| 8(c) ALT | Equation of line OP is $y = mx$ $(x - 4)^2 + (mx - 6)^2 = 16$ $(m^2 + 1)x^2 - (8 + 12m)x + 36 = 0$ $\Delta = 0 \Rightarrow (8 + 12m)^2 - 4 \times (m^2 + 1) \times 36 = 0$ $m = \frac{5}{12}$ y-coordinate of T $y = \frac{5}{12} \times 4$ $z = 4 + \frac{5}{3}i$ | M1 A1 M1 A1 M1 A1 | ft their m Exact value for real and imaginary parts |



| Q | Answer | Marks | Comments |
|------|---|-------|----------|
| 9(a) | $(x-12)^2 + y^2 = (x+12)^2$ | M1 | |
| | $x^2 - 24x + 144 + y^2 = x^2 + 24x + 144$ | A1 | |
| | $y^2 = 48x$ | | |
| | | 2 | |

| Q | Answer | Marks | Comments |
|------|---------------------|-------|------------------------|
| 9(b) | $(y-4)^2 = 48(x-5)$ | B1 B1 | B1 for LHS, B1 for RHS |
| | | 2 | |

| Q | Answer | Marks | Comments |
|---------|--|-------|---|
| 9(c)(i) | Parabola with vertex facing left | B1 | |
| | Parabola with vertex in 1 st quadrant | B1 | ft their answer to part (b) if translation $\begin{bmatrix} 5 \\ -4 \end{bmatrix}$ is used, which leads to a parabola with a vertex in the 4 th quadrant |
| | Lines OQ and OR, Q and R marked correctly, R to the right of Q | B1 | See artwork below |



| | | | |
|--|--|---|--|
| | | 3 | |
|--|--|---|--|

| Q | Answer | Marks | Comments |
|----------|---|---|---|
| 9(c)(ii) | $y = mx$ is a tangent so $(mx - 4)^2 = 48(x - 5)$ $m^2x^2 - 8mx + 16 - 48x + 240 = 0$ $m^2x^2 - (8m + 48)x + 256 = 0$ $\Delta = 0 \Rightarrow (8m + 48)^2 - 4m^2(256) = 0$ $960m^2 - 768m - 2304 = 0$ $5m^2 - 4m - 12 = 0$ $m = 2$ or $m = -\frac{6}{5}$ $m = 2: 4x^2 - 64x + 256 = 0$ and $x = 8$ or $m = -\frac{6}{5}: \frac{36}{25}x^2 - \frac{192}{5}x + 256 = 0$ and $x = \frac{40}{3}$ $Q(8, 16)$ $R\left(\frac{40}{3}, -16\right)$ | <p>M1</p> <p>A1ft</p> <p>M1</p> <p>A1ft</p> <p>A1ft</p> <p>M1</p> <p>A1</p> <p>A1</p> | <p>ft their part (b)</p> <p>for either equation (oe) ft their part (b)</p> <p>ft their part (b)</p> |
| | | 8 | |
| | Question 9 Total | 15 | |