

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

I declare this is my own work.

INTERNATIONAL A-LEVEL PHYSICS

Unit 5 Physics in practice

Thursday 19 January 2023

07:00 GMT

Time allowed: 2 hours

Materials

For this paper you must have:

- a Data and Formulae Booklet as a loose insert
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate
- a protractor.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- All working must be shown.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
TOTAL	



Section A

Answer **all** questions in this section.

0	1
---	---


Fuse wire is designed to melt when the current in the wire reaches a certain value. The fusing current I_F is the minimum current that will cause a length of fuse wire to melt.

0	1	.	1
---	---	---	---

A student is provided with a length of fuse wire and normal electrical equipment.

Figure 1 shows the circuit symbol for a length of fuse wire.

Figure 1



Draw a suitable circuit diagram that the student can use to determine I_F .

[2 marks]



7


0 3

0 2

A student is investigating the cooling of oil. She puts 250 g of hot oil into a metal container and records the temperature θ of the oil for 300 s as it cools.

Figure 2 is a plot of θ against time t for the cooling oil.

Figure 2



Do not write
outside the
box



0 2 . 1

Explain **one** feature of the container that will ensure a maximum rate of heat transfer from the oil to the surroundings at any given temperature difference.

[1 mark]

0 2 . 2

Draw the line of best fit for the data plotted in **Figure 2**.

[1 mark]

0 2 . 3

Explain how **Figure 2**, together with other data, can be used to determine the total heat transfer from the oil during the first 300 s.

[2 marks]

0 2 . 4

Show that the rate of change of temperature of the oil at a temperature of 45 °C is approximately 0.1 K s⁻¹.

[2 marks]

0 2 . 5

At a temperature of 45 °C, the rate of heat transfer from the oil is 54 J s⁻¹.

Calculate the specific heat capacity of the oil.

[2 marks]

specific heat capacity = _____ J kg⁻¹ K⁻¹

8

Turn over ►



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



0 3

Figure 3 shows apparatus that a student uses to measure the wavelength of a horizontal beam of monochromatic light.


The light is initially reflected downwards by a glass plate towards a lens and a mirror.

The light travels through the lens.

The lower surface of the lens reflects some of the light upwards.

Light from the lower surface of the lens and light from the mirror surface then superpose to form a circular interference pattern that is photographed.

Figure 3



Question 3 continues on the next page

*Do not write
outside the
box*

Turn over ►



Figure 4 shows an actual-size photograph of the interference pattern. The pattern is a series of bright and dark circles called rings.


Figure 4



The diameter of the n th bright ring on this actual-size photograph is d_n . For example, the diameter of the 4th bright ring is d_4 .

The student makes a magnified image of the interference pattern that is **10 times larger** than the actual size. **Figure 5** shows the magnified image.

Figure 5



The diameter of the 4th bright ring, counting from the centre of the **magnified** image, is D_4 .

03.1

The student uses a millimetre scale to measure D_4 from the image in **Figure 5**.
The student measures D_4 as 44 mm.

Calculate the percentage uncertainty in D_4 .

[1 mark]

uncertainty in $D_4 = \pm$ _____ %

03.2

Calculate, using your answer to Question **03.1**, the absolute uncertainty in d_4^2 .

[2 marks]

absolute uncertainty in $d_4^2 = \pm$ _____ mm²

03.3

Determine, using **Figure 5**, an accurate value for d_9 , the actual diameter of the 9th bright ring.

[2 marks]

$d_9 =$ _____ mm

The wavelength λ of the monochromatic light is given by $\lambda = \frac{d_9^2 - d_4^2}{20r}$

where $r = (3.40 \pm 0.01)$ m = the radius of the curved surface of the lens in **Figure 3**.

03.4

Calculate λ .

[1 mark]

$\lambda =$ _____ m

Question 3 continues on the next page

Turn over ►



0 3 . 5

Calculate the percentage uncertainty in your value for λ .

Assume that the absolute uncertainty in d_9^2 is the same as the absolute uncertainty in d_4^2 calculated in Question **03.2**.

[3 marks]

uncertainty in λ = _____ %

0 3 . 6

Explain **one** advantage of using **Figure 5** rather than **Figure 4** to determine λ .**[1 mark]**

10



Turn over for the next question

*Do not write
outside the
box*


**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Turn over ►



0 4

Figure 6 shows a mass on the end of a string. The mass rotates freely in a horizontal circle at a constant speed.

Figure 6

0 4 . 1

State and explain the origin of the centripetal force acting on the mass.

[1 mark]

The period of rotation T of the mass is measured for a range of values of the length l of the string.

The radius r of the circle is kept constant as l is varied.

0 4 . 2

Table 1 shows measurements of l and T and some derived data from the experiment.

Table 1

l / m	T / s	$\sqrt{(l^2 - r^2)} / \text{m}$	T^2 / s^2
0.600	1.55	0.587	2.40
0.700	1.67	0.689	2.79
0.800	1.79	0.790	3.20
0.900	1.90	0.891	3.61
1.000	2.01	0.992	4.04




Draw, on **Figure 7**, a graph of T^2 against $\sqrt{l^2 - r^2}$.

Use false origins on both axes.

[4 marks]

Figure 7

T^2 / s^2



$\sqrt{l^2 - r^2} / \text{m}$

Question 4 continues on the next page

Turn over ►



0 4 . 3

Determine the gradient of the graph you have drawn on **Figure 7**.**[2 marks]**

gradient = _____

0 4 . 4

A student suggests that the relationship between T and l is given by

$$T^2 = \frac{4\pi^2}{k} \sqrt{(l^2 - r^2)}$$

Determine k .**[2 marks]** $k =$ _____ m s^{-2}

0 4 . 5

Determine whether or not the graph supports the idea that T^2 is directly proportional to $\sqrt{l^2 - r^2}$.**[2 marks]**

END OF SECTION A

Turn over for Section B

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Turn over ►




Section BAnswer **all** questions in this section.**0 5**

Figure 8 shows a belt-and-pulley system used by a car engine to drive an electrical generator.

The engine makes pulley **A** rotate and moves the belt.

The moving belt rotates pulley **B**. This drives the generator which is attached to **B**.

Figure 8

The car engine delivers a mean power of 930 W to the belt-and-pulley system.
Assume that the belt-and-pulley system is perfectly efficient.

0 5 . 1

The generator has an efficiency of 83%.

The generator has an rms output of 14 V.

Calculate the rms output current of the generator.

[3 marks]

rms output current = _____ A



0 5 . 2

The belt does not slip on the pulleys. The angular speed of pulley **B** is 480 rad s^{-1} .
The diameter of pulley **A** is 22 cm.
The diameter of pulley **B** is 12 cm.


Calculate, in revolutions per minute, the frequency of rotation of pulley **A**.

[3 marks]

frequency of rotation = _____ revolutions per minute

The tension F in the belt acts on pulley **B** at point **R** in the direction shown in **Figure 9**.

Figure 9



0 5 . 3

F exerts a torque on pulley **B**. The power delivered to pulley **B** is 930 W.

Calculate F .

[3 marks]

$F =$ _____ N

Question 5 continues on the next page

Turn over ►



0 5 . 4

The length of belt between **Q** and **R** is 0.25 m when the system is at rest and there is no tension in the belt.

Pulley **A** is made to rotate more quickly.

The tension in the belt between **Q** and **R** is now 55 N.

The belt between **Q** and **R** has an extension of 1.0 mm compared with its unstretched length.

The Young modulus of the belt material is 340 MPa.

Calculate the cross-sectional area of the belt.

[3 marks]

cross-sectional area = _____ m²

0 5 . 5

After the belt has been used for a long time, it becomes permanently stretched and slips on the pulleys.

Explain how the performance of the whole system is affected by this slipping.

[2 marks]



0 6

This question is about an attempt in 1798 to calculate the average density of the Earth.

Figure 10 shows a uniform rod, of negligible mass, suspended by a vertical wire from a fixed support.


Two small lead spheres are attached to the rod, one at each end. Each sphere acts as a point mass.

The mass m of each sphere is 0.730 kg.

P is the centre of the rod and the distance from **P** to each sphere is $\frac{L}{2}$.

The moment of inertia I of the spheres about **P** is 1.26 kg m^2 .

Figure 10



0 6

. 1

Show that L is approximately 1.9 m.

[3 marks]

Question 6 continues on the next page

Turn over ►



Figure 11 shows the rod viewed from above. The vertical wire is not shown.

The rod is turned through a small angle θ . This causes the vertical wire to twist and exert a torque on the rod.

Figure 11



When the rod is released it performs oscillations about **P** in a horizontal plane.

The frequency f of the oscillations is given by

$$f = \frac{1}{2\pi} \sqrt{\frac{k}{I}}$$

where k is a constant.

0 6 . 2 The time period of the oscillation is 424 s.

Suggest why this period is difficult to measure.

[2 marks]

0 6 . 3 Show that k is approximately $2.8 \times 10^{-4} \text{ N m rad}^{-1}$.

[2 marks]



The rod returns to rest at its equilibrium position.

A large lead sphere **S** of mass M is placed near each small sphere.

Gravitational forces F act between each **S** and its nearest small sphere, producing a torque on the rod. This twists the wire, leading to a torque T in the wire.


The rod turns horizontally through an angle α to reach a new equilibrium position.

The torque now exerted by the forces F is equal and opposite to T .

The rod does not oscillate.

Figure 12 shows the equilibrium position viewed from above.

Figure 12



At equilibrium, the distance between the centre of each **S** and the centre of its nearest small sphere is d .

0 6 . 4 Show that the gravitational constant G is given by:

$$G = \frac{Td^2}{MmL}$$

[2 marks]

Question 6 continues on the next page

Turn over ►



At equilibrium, T is given by:

$$T = k\alpha$$

0 6 . 5

In one experiment:

$$M = 158 \text{ kg}$$

$$m = 0.730 \text{ kg}$$

$$d = 0.225 \text{ m}$$

$$k = 2.8 \times 10^{-4} \text{ N m rad}^{-1}$$

$$\alpha = 9.30 \times 10^{-4} \text{ rad}$$

$$L = 1.90 \text{ m.}$$

Show that these data suggest a value for G of approximately $6 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$.

[2 marks]

0 6 . 6

Calculate, using the value for G in Question **06.5**, the average density of the Earth.

$$g = 9.81 \text{ N kg}^{-1}$$

[3 marks]


average density of the Earth = _____ kg m^{-3}

14



07

An ion drive is attached to a satellite and provides thrust to make small changes to the satellite's velocity. **Figure 13** shows the main features of the ion drive.

Figure 13

Xenon-131 atoms enter the chamber with negligible speed and pass through a positive grid. At the positive grid, a single electron is removed from each atom to form a xenon ion.

The positive grid is at a potential of +1100 V and the negative grid is at a potential of -200 V. The ions are accelerated through this potential difference.

A stream of ions is then ejected at high speed through the negative grid into space.

07.1

Explain why the ejection of a stream of ions in one direction causes the ion drive to accelerate in the opposite direction.

[3 marks]

Question 7 continues on the next page

Turn over ►



0 7 . 2

Show that the speed of the ions relative to the negative grid as they pass through it is approximately 44 km s^{-1} .

$$\text{mass of xenon-131 ion} = 2.18 \times 10^{-25} \text{ kg}$$

[2 marks]**0 7 . 3**

The ion drive produces a thrust of 0.049 N .

Calculate the mass of xenon ejected through the negative grid each second.

[1 mark]

mass ejected each second = _____ kg



One type of power supply used for ion drives is a radioisotope thermoelectric generator (RTG).

One RTG initially contains 7.80 kg of plutonium-238 that decays by alpha emission. The decay energy is 5.69 MeV and the decay constant is $2.50 \times 10^{-10} \text{ s}^{-1}$.

0 7 . 4

Explain why energy is released when plutonium-238 decays.

[2 marks]

0 7 . 5

Show that the RTG contains approximately 2.0×10^{25} nuclei of plutonium-238.

[2 marks]

Question 7 continues on the next page

Turn over ►



07.6

When 1.0 J of energy is transferred through nuclear decay, the RTG transfers 68 mJ into electrical energy.

The ion drive requires a power of approximately 1.0 kW.

Determine whether or not the RTG is able to deliver this power.

[4 marks]

07.7

Another type of RTG uses 7.80 kg of strontium-90 instead of plutonium-238. This strontium-based RTG has the same efficiency as the plutonium-based RTG. Strontium-90 has a decay constant of $7.63 \times 10^{-10} \text{ s}^{-1}$ and a decay energy of 0.546 MeV.

Suggest how this change of fuel type affects the performance of the RTG.

[2 marks]

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



[illegible]

*Do not write
outside the
box*

[illegible]

*Do not write
outside the
box*

[illegible]

*Do not write
outside the
box*

[illegible]

There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.oxfordaqaexams.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2023 Oxford International AQA Examinations and its licensors. All rights reserved.



3 2



2 3 1 X P H 0 5

IB/M/Jan23/PH05