Surname	Other nar	mes
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Statistics & Advanced/Advance	_	
Tuesday 10 June 2014 – Mo Time: 1 hour 30 minutes	orning	Paper Reference WST01/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Values from the statistical tables should be quoted in full. When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

P 4 3 1 4 4 A 0 1 2 4

Turn over ▶



1. A medical researcher is studying the relationship between age (x years) and volume of blood (y ml) pumped by each contraction of the heart. The researcher obtained the following data from a random sample of 8 patients.

Age (x)	20	25	30	45	55	60	65	70
Volume (y)	74	76	77	72	68	67	64	62

[You may use $\sum x = 370$, $S_{xx} = 2587.5$, $\sum y = 560$, $\sum y^2 = 39418$, $S_{xy} = -710$]

(a) Calculate S_{yy}

(2)

(b) Calculate the product moment correlation coefficient for these data.

(2)

(c) Interpret your value of the correlation coefficient.

(1)

The researcher believes that a linear regression model may be appropriate to describe these data.

(d) State, giving a reason, whether or not your value of the correlation coefficient supports the researcher's belief.

(1)

(e) Find the equation of the regression line of y on x, giving your answer in the form y = a + bx

(4)

Jack is a 40-year-old patient.

- (f) (i) Use your regression line to estimate the volume of blood pumped by each contraction of Jack's heart.
 - (ii) Comment, giving a reason, on the reliability of your estimate.

(2)





estion 1 continued		
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_



2. The table below shows the distances (to the nearest km) travelled to work by the 50 employees in an office.

Distance (km)	Frequency (f)	Distance midpoint (x)
0-2	16	1.25
3 – 5	12	4
6 – 10	10	8
11 – 20	8	15.5
21 – 40	4	30.5

[You may use
$$\sum fx = 394$$
, $\sum fx^2 = 6500$]

A histogram has been drawn to represent these data.

The bar representing the distance of 3-5 has a width of 1.5 cm and a height of 6 cm.

- (a) Calculate the width and height of the bar representing the distance of 6-10
- (b) Use linear interpolation to estimate the median distance travelled to work. (2)
- (c) (i) Show that an estimate of the mean distance travelled to work is 7.88 km.
 - (ii) Estimate the standard deviation of the distances travelled to work. (4)
- (d) Describe, giving a reason, the skewness of these data. (2)

Peng starts to work in this office as the 51st employee.

She travels a distance of 7.88 km to work.

- (e) Without carrying out any further calculations, state, giving a reason, what effect Peng's addition to the workforce would have on your estimates of the
 - (i) mean,
 - (ii) median,
 - (iii) standard deviation

of the distances travelled to work.

(3)



estion 2 continued	



3. A biased four-sided die has faces marked 1, 3, 5 and 7. The random variable X represents the score on the die when it is rolled. The cumulative distribution function of X, F(x), is given in the table below.

X	1	3	5	7
F(x)	0.2	0.5	0.9	1

(a) Find the probability distribution of X

(4)

(b) Find P($2 < X \le 6$)

(2)

(c) Write down the value of F(4)

(1)



4. The random variable $Y \sim N(\mu, \sigma^2)$

Given that P(Y < 17) = 0.6 find

(a) P(Y > 17)

(1)

(b) $P(\mu < Y < 17)$

(2)

(c) $P(Y < \mu \mid Y < 17)$

(2)

()



5. The discrete random variable *X* has the following probability distribution

x	-2	0	2	4
P(X=x)	а	b	а	С

where a, b and c are probabilities.

Given that E(X) = 0.8

(a) find the value of c.

(2)

Given also that $E(X^2) = 5$ find

(b) the value of a and the value of b,

(4)

(c) Var(X)

(2)

The random variable Y = 5 - 3X

Find

(d) E(*Y*)

(1)

(e) Var(Y)

(2)

(f) $P(Y \ge 0)$

(4)



nestion 5 continued	






6. The Venn diagram below shows the probabilities of customers having various combinations of a starter, main course or dessert at Polly's restaurant.

S = the event a customer has a starter.

M = the event a customer has a main course.

D = the event a customer has a dessert.



Given that the events S and D are statistically independent

(a) find the value of p.

(4)

(b) Hence find the value of q.

(2)

- (c) Find
 - (i) $P(D \mid M \cap S)$
 - (ii) $P(D \mid M \cap S')$

(4)

One evening 63 customers are booked into Polly's restaurant for an office party. Polly has asked for their starter and main course orders before they arrive.

Of these 63 customers

27 ordered a main course and a starter,

36 ordered a main course without a starter.

(d) Estimate the number of desserts that these 63 customers will have.

(2)







7.	One event at <i>Pentor</i> sports day is throwing a tennis ball. The distance a child throws a tennis ball is modelled by a normal distribution with mean 32 m and standard deviation 12 m. Any child who throws the tennis ball more than 50 m is awarded a gold certificate.			
	(a) Show that, to 3 significant figures, 6.68% of children are awarded a gold certificate. (3)			
	A silver certificate is awarded to any child who throws the tennis ball more than d metres but less than 50 m.			
	Given that 19.1% of the children are awarded a silver certificate,			
	(b) find the value of d . (4)			
	Three children are selected at random from those who take part in the throwing a tennis ball event.			
	(c) Find the probability that 1 is awarded a gold certificate and 2 are awarded silver certificates. Give your answer to 2 significant figures.			
	(3)			





Question 7 continued		Leav blan
		Q
	(Total 10 marks)	
	TOTAL FOR PAPER: 75 MARKS	
END		