



# Mark Scheme (Final)

October 2019

Pearson Edexcel IAL Mathematics

Mechanics 2 (WME02/01)

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

October 2019

Publications Code WME02\_01\_1910\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2019






**October 2019**  
**Mechanics 2 - WME02**  
**Mark Scheme - Final**

Q	Scheme	Marks	Notes
<b>1</b>	Moments about $y$ -axis	M1	Or a parallel axis. Require all terms and dimensionally correct. Condone sign errors $3m\bar{x}$ or $3\bar{x}$ is a method error
	$(k+4)m\bar{x} = 2m + 12m + km$	A1	Or equivalent
	Moments about $x$ -axis	M1	Or a parallel axis. Require all terms and dimensionally correct. Condone sign errors $3m\bar{y}$ or $3\bar{y}$ is a method error
	$(k+4)m\bar{y} = m + 9m + 5km$	A1	Or equivalent
			Allow the first 4 marks for a combined equation in vector format
	Use of $\bar{y} = 2\bar{x}$ to form equation in $k$	M1	
			Allow the first 5 marks if they go straight to an equation equating the right hand sides e.g. $m + 9m + 5km = 2(2m + 12m + km)$ with no incorrect statement about the total mass seen
	$10 + 5k = 2(14 + k) \Rightarrow k = 6$	A1	With no errors seen
		<b>(6)</b>	
		<b>[6]</b>	
<b>2</b>	Use of $F = \frac{16000}{U}$	M1	$\frac{16}{U}$ scores M0
	Equation of motion.	M1	All terms required. Dimensionally correct. Condone sign errors and sin/cos confusion.
	$F - 30U - 750g \sin \theta = 0$	A1	Correct unsimplified equation in $F$ or their $F$
	$\Rightarrow \frac{16000}{U} - 30U - 490 = 0$ $30U^2 + 490U - 16000 = 0$	DM1	Form and solve a quadratic equation in $U$ Horizontal form seen or implied. Dependent on the two preceding M marks Need to see working if solving an incorrect quadratic or obtain an incorrect answer.
	$U = 16.3$ or $U = 16$ only	A1	Max 3 sf
		<b>(5)</b>	
		<b>[5]</b>	





Q	Scheme			Marks	Notes
5a		Mass ratio	From $BC$	B1 B1	Mass ratios correct Vertical distances correct
	Large $\Delta$	60 (4)	$4a$		
	Small $\Delta$	15 (1)	$8a$		
	rectangle	15 (1)	$4.5a$		
	$T$	30 (2)	$\bar{y}$		
	Moments about $BC$			M1	Or a parallel axis. Signs correct for their split.
	$2\bar{y} = 16a - 8a - 4.5a = \frac{7}{2}a$			A1	Correct unsimplified equation
	$\bar{y} = \frac{7}{4}a$			A1	<b>Given Answer</b> Need to see a linear equation in $\bar{y}$
				(5)	
5b					
	$\bar{x} = 5a$ from $B$			B1	Or equivalent. Seen or implied Might be seen in (a) but needs to be used here.
	$\alpha = \tan^{-1} \left( \frac{\frac{5}{2}a}{6a - \bar{y}} \right)$			M1	Trig ratio of a relevant angle in a triangle involving the c of m
	$\theta = \alpha + \tan^{-1} \frac{5}{12}$			M1	Correct method to find the required angle e.g $\theta = 180^\circ - \tan^{-1} \frac{6}{2.5} - \tan^{-1} \frac{6a - \bar{y}}{2.5a}$
	$= \tan^{-1} \frac{10}{17} + \tan^{-1} \frac{5}{12}$			A1	Correct unsimplified expression for $\theta$ seen
					$\tan^{-1} \frac{10}{17} = 30.46\dots, \tan^{-1} \frac{17}{10} = 59.5\dots,$ $\tan^{-1} \frac{5}{12} = 22.6\dots, \tan^{-1} \frac{12}{5} = 67.38\dots$
	$\theta = 53$			A1	Q asks for a whole number
				(5)	
					See over for alternatives




<b>5balt 1</b>	$\bar{x} = 5a$ from $B$	B1	Or equivalent. Seen or implied Might be seen in (a) but needs to be used here.
	Correct method to find all 3 sides of a triangle containing $\theta$ (or two sides of a right angled triangle containing $\theta$ )	M1	e.g.: $BO^2 = (5a)^2 + \left(\frac{7a}{4}\right)^2 = \frac{449}{16}a^2, AB^2 = \frac{169}{4}a^2$ $AO^2 = \left(\frac{5a}{2}\right)^2 + \left(6a - \frac{7a}{4}\right)^2 = \frac{389}{16}a^2$
	Correct method to find the required angle	M1	e.g. $\cos \theta = \frac{AB^2 + AO^2 - OB^2}{2AB \cdot AO}$
	Correct unsimplified expression in $\theta$	A1	( $\cos \theta = 0.60062$ )
	$\theta = 53$	A1	Q asks for a whole number
		(5)	
<b>5balt 2</b>	$\bar{x} = 5a$ from $B$	B1	Or equivalent. Seen or implied Might be seen in (a) but needs to be used here.
	If $O$ is c of m, $\overrightarrow{AB} = \begin{pmatrix} -2.5a \\ -6a \end{pmatrix}, \overrightarrow{AO} = \begin{pmatrix} 2.5a \\ -4.25a \end{pmatrix}$	M1	Expressions for $\overrightarrow{AB}$ and $\overrightarrow{AO}$
	Use of scalar product to find $\cos \theta$	M1	
	$\cos \theta = \frac{\overrightarrow{AB} \cdot \overrightarrow{AO}}{ \overrightarrow{AB}  \cdot  \overrightarrow{AO} } = \frac{-6.25 + 25.5}{\frac{13}{2} \times \frac{\sqrt{389}}{4}}$	A1	Correct unsimplified expression for $\cos \theta$
	$\theta = 53$	A1	Q asks for a whole number
		(5)	
<b>5balt 3</b>	$\bar{x} = 5a$ from $B$	B1	Or equivalent. Seen or implied Might be seen in (a) but needs to be used here.
	If $O$ is c of m, find coordinates of point of intersection between the line $AO$ and the perpendicular line through $B$	M1	Relative to $B$ : $A(2.5a, 6a), O(5a, 1.75a)$ $\Rightarrow AO: y = -1.7x + 10.25a$ If perpendicular through $B$ intersects $AO$ at $L$ , $\Rightarrow BL: y = \frac{10}{17}x \Rightarrow L\left(\frac{3485}{778}, \frac{1025}{389}\right)$
	Correct method to find the required angle	M1	e.g. $\sin \theta = \frac{BL}{BA}$
	Correct unsimplified expression in $\theta$	A1	$= \frac{\sqrt{\left(\frac{3485}{778}\right)^2 + \left(\frac{1025}{389}\right)^2}}{6.5}$
	$\theta = 53$	A1	Q asks for a whole number
		(5)	
			See over for alternative

<b>5balt 4</b>	$\bar{x} = 5a$ from $B$	B1	Or equivalent. Seen or implied Might be seen in (a) but needs to be used here.
	If $O$ is c of m, find coordinates of point of intersection between the line $AB$ and the perpendicular line through $O$	M1	$AB: y = \frac{12}{5}x$ If perpendicular through $O$ intersects $AB$ at $H$ , $OH: y = -\frac{5}{12}x + \frac{23}{6}, H\left(\frac{230}{169}, \frac{552}{169}\right)$
	Correct method to find the required angle	M1	e.g. $\sin \theta = \frac{OH}{OA}$
	Correct unsimplified expression in $\theta$	A1	$= \frac{\frac{205}{52}}{\sqrt{\frac{389}{16}}}$
	$\theta = 53$	A1	Q asks for a whole number
		<b>[10]</b>	



<b>7a</b>			
	Use of CLM	M1	Need all terms. Dimensionally correct. Condone sign errors
	$2mku - 3m \times 4u = -2m \times 2u + 3m \times u$ $(2k - 12 = -4 + 3)$	A1	Correct unsimplified
	$\Rightarrow k = \frac{11}{2}$	M1	Solve for $k$
	$\text{KE lost} = \frac{1}{2} 2m(k^2 - 4)u^2 + \frac{1}{2} 3m(16 - 1)u^2$	M1	Complete expression. Dimensionally correct. Allow $\pm$ . In $k$ or their $k$
	$\left( = mu^2(k^2 - 4) + \frac{45mu^2}{2} \right)$		
	$= \frac{195mu^2}{4}, \lambda = \frac{195}{4}$	A1	Or equivalent (48.75). Accept 49, 48.8 ISW once correct answer seen.
		<b>(5)</b>	
<b>7b</b>	Use of CLM: $4mu = 8mw - 2mv$	M1	Dimensionally correct. Condone incorrect signs.
	Use of impact law:	M1	Used the right way round.
	$v + w = 2eu$	A1	Both equations correct unsimplified. Signs consistent.
	Solve for $v$ or a multiple of $v$	M1	$(v_B)$
	$(w = 2eu - v \Rightarrow 10v = 16eu - 4u)$		
	Inequality in $e$ : $10v > 10u \Rightarrow 16eu > 14u$	M1	Complete correct method with signs consistent with the direction of their $v$ Upper limit of 1 not necessary No extra incorrect inequality
	$\Rightarrow \frac{7}{8} < e \leq 1$	A1	Or equivalent. Want both limits. Must be $\leq 1$ .
			NB: The last two M marks might appear in reverse order. The inequality $v > u$ can be used with an expression for $w$ to form the inequality in $e$ .
		<b>(6)</b>	
		<b>[11]</b>	

8a			
	$P \text{ and } Q \text{ collide} \Rightarrow \text{at same vertical height}$ $\Rightarrow bt - \frac{1}{2}gt^2 = dt - \frac{1}{2}gt^2$ $\Rightarrow bt = dt, b = d$	B1	Obtain <b>given result</b> correctly No incorrect statements seen
		(1)	
8b	$10 = 3d - \frac{9}{2}g$	M1	Complete method using <i>suvat</i> to form an equation in $d$ e.g. use of $s = ut + \frac{1}{2}at^2$ with correct values substituted but condone sign error
	$\Rightarrow d = 18 \text{ (18.0)}$	A1	Follows use of 9.8. Not $\frac{541}{30}$ Must be a scalar
		(2)	
8c	$t = 3$ , vertical speed $v = d - 3g = -11.4$	M1	Condone sign errors
	$-11.4 = 11.4 - gt$	M1	Using symmetry with their 11.4
	$t = 2.3 \text{ (2.32)(s)}$	A1	Not $\frac{341}{147}$
		(3)	
8calt1	Vertical distance $10 = dt - \frac{g}{2}t^2$	M1	Condone sign errors
	Solve for $t$ and subtract $\left(3, \frac{200}{3 \times 98}\right)$	M1	
	$t = 2.3 \text{ (2.32)(s)}$	A1	
		(3)	
8calt2	Time to top: $0 = d - gt, (t = 1.84)$	M1	Condone sign errors
	Solve for required time: $T = 2(3 - 1.84)$	M1	
	$= 2.3 \text{ (s)}$	A1	
		(3)	
8d	Horizontal distance: $60 = 12 \times 3 + 3c$	M1	Allow if seen earlier and used here
	$(c = 8)$	A1	Correct unsimplified <b>equation</b> for horizontal distance. Allow $c = -8$ if used correctly later
	$\mathbf{v} = -8\mathbf{i} - 11.4\mathbf{j}$	M1	For their 8 and 11.4
	Direction $\tan^{-1} \frac{11.4}{8}$	DM1	Use of trig to find a relevant angle Dependent on the previous M1

	Downwards at $54.9^\circ$ to the horizontal ( $55^\circ$ ) ( $35.1^\circ$ to the vertical)	A1	Or equivalent. Direction must be stated or clearly implied by a diagram. 0.959 rads (0.96 rads) Max 3 sf
		(5)	
8dalt	Horizontal distance: $60 = 12 \times 3 + 3c$	M1	Allow if seen earlier and used here
	( $c = 8$ )	A1	Correct unsimplified <b>equation</b> for horizontal distance. Allow $c = -8$ if used correctly later
	$\mathbf{v} = -8\mathbf{i} + 18.0\mathbf{j}$ and conservation of energy $\frac{1}{2}mv^2 - \frac{1}{2}m(8^2 + 18^2) = -mg \times 10$ ( $v = 13.9$ )	M1	For their 8 and 18.0
	Direction $\cos^{-1} \frac{8}{13.9}$	DM1	Use of trig to find a relevant angle Dependent on the previous M1
	Downwards at $54.9^\circ$ to the horizontal ( $55^\circ$ ) ( $35.1^\circ$ to the vertical)	A1	Or equivalent. Direction must be stated or clearly implied by a diagram. 0.959 rads (0.96 rads) Max 3 sf
		(5)	
8e			
	$\frac{-x}{12} \times \frac{18}{12} = -1$ , $\left( \frac{-x}{12} \times \frac{d}{12} = -1 \right)$	M1	Complete method (with their $d$ ) using velocity to find $x$ Could be in scalar product format
	$x = 8$	A1	7.99 or better Allow +/-
	$-8 = 18 - 9.8T$ , ( $-x = d - 9.8T$ )	DM1	Form equation in $T$ . Signs correct Dependent on the previous M1
	$T = 2.7$ (2.65)	A1	Follows use of 9.8
		(4)	
8e alt	$\tan \alpha = \frac{d}{12}$ , $\beta = 90 - \alpha$	M1	Complete method (with their $d$ ) to find the direction of motion
	$= 33.64^\circ$	A1	
	$-\tan \beta = \frac{d - 9.8T}{12}$	DM1	Form equation in $T$ . Signs correct. Dependent on the previous M1
	$T = 2.7$ (2.65)	A1	CSO – need to be convinced by any changes of sign. Follows use of 9.8
		(4)	
		[15]	

