Please check the examination det	ails below	before ente	ring your candic	date information
Candidate surname			Other names	
Pearson Edexcel International Advanced Level	Centre	e Number		Candidate Number
Wednesday 6	5 No	oven	nber	2019
Morning (Time: 1 hour 30 minut	es)	Paper Re	eference W	WA12/01
Mathematics				
International Advanced Subsidiary/Advanced Level Pure Mathematics P2				
You must have: Mathematical Formulae and Sta	itistical 7	Tables (Lil	ac), calculato	Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use black ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶







(a) Find, using calculus, the x coordinates of the stationary points of C .	
(b) Hence find the values of x for which y is increasing.	

	Leave blank
Question 1 continued	
	21
(Total 6 marks)	



=• 1	The addit population of a town at the start of 2019 is 25 000
n	A model predicts that the adult population will increase by 2% each year, so that the number of adults in the population at the start of each year following 2019 will form a geometric sequence.

(a) Find, according to the model, the adult population of the town at the start of 2032

(3)

It is also modelled that every member of the adult population gives £5 to local charity at the start of each year.

(b) Find, according to these models, the total amount of money that would be given to local charity by the adult population of the town from 2019 to 2032 inclusive. Give your answer to the nearest £1 000

(3)	

Question 2 continued		blank
	Question 2 continued	
(Total 6 marks)		Q2
	(Total 6 marks)	



3. (a) Find the first 4 terms, in ascending powers of x, in the binomial expansion of

$$\left(1+\frac{x}{4}\right)^{12}$$

giving each coefficient in its simplest form.

(3)

(b) Find the term independent of x in the expansion of

$$\left(\frac{x^2+8}{x^5}\right)\left(1+\frac{x}{4}\right)^{12}$$

(3)

6

Question 3 continued	Leave blank
	Q3
(Total 6 marks)	



4.	f(x) = (x -	$3)(3x^2 + x -$	+ a) - 35	where a is a co	nstant

(a) State the remainder when f(x) is divided by (x-3).

(1)

Given (3x - 2) is a factor of f(x),

(b) show that a = -17

(2)

(c) Using algebra and showing each step of your working, fully factorise f(x).

(5)	

uestion 4 continued	



Question 4 continued	

Question 4 continued	Leave
	Q4
(Total 8 marks)	
, ,	



5. (a) Given 0 < a < 1, sketch the curve with equation

$$y = a^{\lambda}$$

showing the coordinates of the point at which the curve crosses the y-axis.

(2)

X	2	2.5	3	3.5	4
у	4.25	6.427	9.125	12.34	16.06

The table above shows corresponding values of x and y for $y = x^2 + \left(\frac{1}{2}\right)^x$

The values of y are given to 4 significant figures as appropriate.

Using the trapezium rule with all the values of y in the given table,

(b) obtain an estimate for
$$\int_{2}^{4} \left(x^{2} + \left(\frac{1}{2} \right)^{x} \right) dx$$
 (3)

Using your answer to part (b) and making your method clear, estimate

(c)
$$\int_{2}^{4} \left(x(x-3) + \left(\frac{1}{2}\right)^{x} \right) \mathrm{d}x$$
 (2)

nestion 5 continued	
	I



Question 5 continue	ed		

Question 5 continued	blank
	Q5
(Total 7 marks)	



6.




Figure 1

Figure 1 shows a sketch of a circle C with centre N(4, -1).

The line *l* with equation $y = \frac{1}{2}x$ is a tangent to *C* at the point *P*.

Find

(a) the equation of line PN in the form y = mx + c, where m and c are constants,

(b) the equation of C.

(5)

(2)

16

uestion 6 continued	




Question 6 continued		

Question 6 continued	Leave
	Q6
(Total 7 marks)	



7. Given $\log_a b = k$, find, in simplest form in terms of k,



(2)



(2)

(iii)
$$\sum_{n=1}^{50} (k + \log_a b^n)$$

(3)

Question 7 continued		Leavo
Question / continued		
		7
	<u>Q</u> ′	/
	(Total 7 marks)	



8. Solutions relying on calculator technology are not acceptable in this question.






Figure 2

Figure 2 shows a sketch of part of a curve with equation

$$y = \frac{8\sqrt{x} - 5}{2x^2} \qquad x > 0$$

The region R, shown shaded in Figure 2, is bounded by the curve, the line with equation x = 2, the x-axis and the line with equation x = 4

Find the exact area of R.

(5)

(ii) Find the value of the constant k such that

$$\int_{-3}^{6} \left(\frac{1}{2} x^2 + k \right) \mathrm{d}x = 55$$

(4)

	L
	b
uestion 8 continued	
	1



Question 8 continued	

Question 8 continued	blank
	Q8
/TT : 3.0	
(Total 9 r	narks)



- 9. Solutions based entirely on graphical or numerical methods are not acceptable in this question.
 - (i) Solve, for $0 \le \theta < 180^{\circ}$, the equation

$$3\sin(2\theta - 10^{\circ}) = 1$$

giving your answers to one decimal place.

(4)

(ii) The first three terms of an arithmetic sequence are

$$\sin \alpha$$
, $\frac{1}{\tan \alpha}$ and $2\sin \alpha$

where α is a constant.

(a) Show that $2\cos\alpha = 3\sin^2\alpha$

(3)

Given that $\pi < \alpha < 2\pi$,

(b) find, showing all working, the value of α to 3 decimal places.

(5)

uestion 9 continued	



Question 9 continued

	Leave
	blank
Question 9 continued	
	Q9
(Total 12 marks)	
(10tai 12 marks)	



10. The curve C has equation

$$y = ax^3 - 3x^2 + 3x + b$$

where a and b are constants.

Given that

- the point (2, 5) lies on C
- the gradient of the curve at (2, 5) is 7
- (a) find the value of a and the value of b.

(4)

(b) Prove that C has no turning points.

(3)

	Lea
	bla
Question 10 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	-
	_
	_
	_
	_



	(Total 7 ma	rke)