Candidate surname	ails below before	Other names
Pearson Edexcel nternational Advanced Level	Centre Num	Candidate Number
Monday 14 Ja	anuai	r y 2019
Afterna (Time - 1 le 20 - 1		
Afternoon (Time: 1 hour 30 minu	ites) Pap	per Reference WFM01/01
Further Pure M Advanced/Advanced S	athen	natics F1

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1.	The point A (12, 12) lies on the parabola with equation $y^2 = 12x$. The point S is the focus
	of this parabola. The line l passes through A and S .
	(a) Find an equation of the line <i>l</i> . (3)
	The line l meets the directrix of the parabola at the point B .
	(b) Find the coordinates of B . (3)

Question 1 continued		Leave
Zuranan z communen		
		01
		Q1
	(Total 6 marks)	



2.

$$f(z) = z^3 - 2z^2 + 16z - 32$$

(a) Show that f(2) = 0

(1)

Leave blank

(b) Use algebra to solve f(z) = 0 completely.

- **(3)**
- (c) Show, on a single Argand diagram, all three roots of the equation f(z) = 0

(2)

Question 2 continued		bla
		Q2
	(Total 6 marks)	



3. (a) Use the standard results for $\sum_{r=1}^{n} r$ and $\sum_{r=1}^{n} r^2$ to show that, for all positive integers n,

$$\sum_{r=1}^{n} (2r+5)^2 = \frac{n}{3} [(an+b)^2 + c]$$

where a, b and c are integers to be found.

(5)

(b) Use the answer to part (a) to evaluate $\sum_{r=0}^{100} (2r+5)^2$

(2)

	Leave
	blank
Question 3 continued	



Question 3 continued		

Question 3 continued		bl
	Q	23
	(Total 7 marks)	



4.

$$f(x) = 2x^3 - \frac{7}{x^2} + 16, \quad x \neq 0$$

The equation f(x) = 0 has a single root α between x = -2 and x = -1

(a) Starting with the interval [-2, -1], use interval bisection twice to find an interval of width 0.25 that contains α .

(3)

The equation f(x) = 0 also has a single root β in the interval [0.6, 0.7].

(b) Taking 0.65 as a first approximation to β , apply the Newton-Raphson procedure once to f(x) to obtain a second approximation to β , giving your answer to 4 decimal places.

(4)

	blank
Question 4 continued	
	_
	_
	_
	_
	_
	_
	Q4
(Total 7 mar	ks)
(=0002 + 20002	



- 5. The rectangular hyperbola H has equation xy = 16The point P, on H, has coordinates $\left(4p, \frac{4}{p}\right)$ where p is a non-zero constant.
 - (a) Show, using calculus, that the tangent to H at the point P has equation

$$x + p^2 y = 8p$$

Given that the tangent to H at the point P passes through the point (7, 1)

(b) use algebra to find the coordinates of the two possible positions of P.

<i>(</i> 1)	

(4)

	Leave
	blank
Question 5 continued	
	1



Question 5 continued		bla
		Q 5
	(Total 8 marks)	



6. It is given that α and β are roots of the equation

$$12x^2 - 3x + 4 = 0$$

Without solving the equation,

(a) find the exact value of $\frac{2}{\alpha} + \frac{2}{\beta}$



(b) find a quadratic equation that has roots $\frac{2}{\alpha} - \beta$ and $\frac{2}{\beta} - \alpha$ giving your answer in the form $ax^2 + bx + c = 0$, where a, b and c are integers to be found.



	Leave
	blank
Question 6 continued	



Question 6 continued		bla
		Q6
	(Total 9 marks)	



7.

$$\mathbf{P} = \begin{pmatrix} -1 & -\sqrt{3} \\ \sqrt{3} & -1 \end{pmatrix}$$

(a) Show that $P^3 = 8I$, where I is the 2 × 2 identity matrix.

(3)

(b) Describe fully the transformation represented by the matrix ${\bf P}$ as a combination of two simple geometrical transformations.

(4)

(c) Find the matrix P^{35} , giving your answer in the form

$$\mathbf{P}^{35} = 2^k \begin{pmatrix} -1 & a \\ b & -1 \end{pmatrix}$$

where k is an integer and a and b are surds to be found.

(2)

20

	Leave blank
Question 7 continued	biank



Question 7 continued

Question 7 continued		Lo bl
	Q	27
	(Total 9 marks)	



8. (i) Prove by induction that, for $n \in \mathbb{Z}^+$

$$\begin{pmatrix} 5 & -8 \\ 2 & -3 \end{pmatrix}^n = \begin{pmatrix} 1 + 4n & -8n \\ 2n & 1 - 4n \end{pmatrix}$$
 (5)

(ii) A sequence of positive numbers is defined by

$$u_1 = 8, \quad u_2 = 40$$

 $u_{n+2} = 8u_{n+1} - 12u_n \qquad n \geqslant 1$

Prove by induction that, for $n \in \mathbb{Z}^+$

$$u_n = 6^n + 2^n {(5)}$$

	Leave
	blank
Question 8 continued	



Question 8 continued	

Question 8 continued		blank
Question o continueu		
		Q8
	/T / 140	70
	(Total 10 marks)	



9. The complex numbers z_1 and z_2 are given by

$$z_1 = -1 - i$$
 and $z_2 = 3 - 4i$

- (a) Find the argument of the complex number $z_1 z_2$. Give your answer in radians to 3 decimal places.
 - **(3)**
- (b) Find the complex number $\frac{z_1}{z_2}$ in the form a + ib, where a and b are rational numbers.
 - **(3)**
- (c) Find the modulus of $\frac{z_1}{z_2}$, giving your answer as a simplified surd.
- **(2)**

(d) Find the values of the real constants p and q such that

$$\frac{p + \mathrm{i}q - 8z_1}{p - \mathrm{i}q - 8z_2} = 3\mathrm{i}$$

(5)

Duestion 9 continued	



Question 9 continued		

	Leave
	blank
Question 9 continued	
Question > continued	



uestion 9 continued	Leave blank
	Q9
(Total 13 marks) TOTAL FOR PAPER: 75 MARKS	
END	